

Positivity of Dim-6 Operators

Azatov, Aleksandr, Diptimoy Ghosh, and Amartya Harsh Singh. "Four-fermion operators at dimension 6: Dispersion relations and UV completions." *Physical Review D* 105.11 (2022): 115019.


Dim-8 operators \rightarrow Straightforward positivity bounds

Some useful applications (e.g. positivity cone)

But hard to measure!

How can we include dim-6 operators into our picture?

$$\begin{aligned}
\left. \frac{\partial}{\partial s} A_{ab \rightarrow ab}(s, 0) \right|_{s=\mu^2} &= \frac{1}{2\pi i} \oint ds \frac{A_{ab \rightarrow ab}(s, 0)}{(s - \mu^2)^2} - (\text{IR pole residues}) \quad C_\infty \\
&= \int \frac{ds}{\pi s} (\sigma_{ab \rightarrow ab} - \sigma_{a\bar{b} \rightarrow a\bar{b}}) + \boxed{\int_0^{2\pi} \frac{d\theta}{2\pi} \frac{A_{ab \rightarrow ab}(|s_\Lambda| e^{i\theta}, 0)}{(|s_\Lambda| e^{i\theta} - \mu^2)^2} |s_\Lambda| e^{i\theta}} \\
&\quad + O(m_{\text{IR}}^2/\mu^2, \mu^2/\Lambda^2) \quad \text{Not necessarily zero} \rightarrow \text{subtleties} \\
&\quad \Lambda^2 \gg \mu^2 \gg m_{\text{IR}}^2
\end{aligned}$$



Example: Four-fermi operators in SMEFT

$$c_{RR}(\bar{e}_R \gamma_\mu e_R)(\bar{e}_R \gamma^\mu e_R)$$

$$-8c_{RR} = \int \frac{ds}{\pi s} (\sigma_{e_R \bar{e}_R} - \sigma_{e_R e_R}) + C_\infty \quad \text{Sum rules}$$

Possible UV completions – negative sign (1)

$$\mathcal{L}_{Z'} = \lambda Z'_\mu \bar{e}_R \gamma^\mu e_R$$

$$c_{RR} = -\frac{\lambda^2}{2M_{Z'}^2} \quad (\text{tree-level})$$

$$A_{\bar{e}_R e_R \rightarrow \bar{e}_R e_R}(s, t) = -2\lambda^2 [23] \langle 14 \rangle \left(\frac{1}{s - M_{Z'}^2} + \frac{1}{t - M_{Z'}^2} \right)$$
$$\stackrel{t \rightarrow 0}{=} -2\lambda^2 s \left(\frac{1}{s - M_{Z'}^2} - \frac{1}{M_{Z'}^2} \right)$$

$$\Rightarrow C_\infty^{(Z')} = \frac{2\lambda^2}{M_{Z'}^2} = \left[\int \frac{ds}{\pi s} (\sigma_{e_R \bar{e}_R} - \sigma_{e_R e_R}) \right]^{(Z')}$$

Possible UV completions – negative sign (2)

$$\text{EFT} : c_{e\mu}(\bar{e}_R\gamma_\mu e_R)(\bar{\mu}_R\gamma^\mu\mu_R)$$

- $\mathcal{L}_{\text{UV}}^{(1)} = \lambda Z_{(1)}^\mu (\bar{e}_R\gamma_\mu\mu_R + h.c.)$

$$c_{e\mu} = -\frac{1}{2} \left[\int \frac{ds}{\pi s} (\sigma_{e_R\bar{\mu}_R} - \sigma_{e_R\mu_R}) \right] = -\frac{|\lambda|^2}{M_{(1)}^2}, \quad \boxed{C_\infty = 0}$$

- $\mathcal{L}_{\text{UV}}^{(2)} = (\lambda_1 Z_{(2)}^\mu \bar{e}_R\gamma_\mu e_R + \lambda_2 Z_{(2)}^\mu \bar{\mu}_R\gamma_\mu\mu_R)$

No t-channel Z(1)

$$\sigma_{e_R\bar{e}_R} = \sigma_{e_R e_R} = 0, \quad c_{e\mu} = -\frac{C_\infty}{2} = -\frac{\lambda_1\lambda_2}{M_{(2)}^2}$$

No UV cross-section,
but infinity pole saturates the dispersion relation

Possible UV completions – positive sign

$$\mathcal{L} = \kappa \phi e_R^T \mathcal{C} e_R + h.c.$$

$$c_{RR} = \frac{|\kappa|^2}{2M_\phi^2}, \quad C_\infty = 0$$

Both signs of the Wilson coefficient are possible, but they suggest different UV models.

Additional idea: utilising t-derivative together

Remmen, Grant N., and Nicholas L. Rodd. "Signs, spin, SMEFT: Sum rules at dimension six." *Physical Review D* 105.3 (2022): 036006.