

Uniform Random Variables

Sepehr Akbari

2024-03-01

Problem 1

Suppose the random variable X has a uniform distribution on the interval $[-1, 7]$.

(a) Compute $P(X > 0)$.

- All probabilities, excluding -1 to 0 . Therefore $\frac{7}{8}$.

(b) Compute $P(X < 0)$.

- Only probabilities from -1 to 0 . Therefore $\frac{1}{8}$.

(c) Briefly explain why $P(X = 0)$ is zero.

- A uniform distribution spreads the probability equally across the entire length. Since any single point (like 0) has no width or length on the ruler, the probability of landing exactly on that point is zero. We can make this more general and say an individual point has a probability of zero.

(d) Compute $P(X < 2 \vee X \geq 4)$.

- This basically means all probabilities, excluding the ones from 2 to 4 . Therefore $\frac{6}{8}$.

Calculating with the normal distribution


Sepehr Akbari

2024-03-01

Problem 1

Suppose X has a normal distribution with mean 24 and standard deviation 5.

(a) Sketch the distribution. Label at least 5 points on the x-axis.




(b) Fill in the blank: $P(24 - 5 < X < 24 + 5) = 68\%$. Use the empirical rule.

(c) Fill in the blank: $P(24 - (5 \times 2) < X < 24 + (5 \times 2)) = 95\%$. Use the empirical rule.

(d) Fill in the blank: $P(24 - (5 \times 3) < X < 24 + (5 \times 3)) = 99.7\%$. Use the empirical rule.

(e) Sketch $P(X \leq 28)$.



(f) Compute $P(X \leq 28)$. Include both R code and a numerical answer.

```
pnorm(28,24,5)
```

```
[1] 0.7881446
```

(g) Compute $P(X > 26)$. Include both R code and a numerical answer.

```
1 - pnorm(26,24,5)
```

```
[1] 0.3445783
```

(h) Compute $P(26 \leq X \leq 28)$. Include both R code and a numerical answer.

```
pnorm(28,24,5) - pnorm(26,24,5)
```

```
[1] 0.1327229
```