## Null space, Row space, Projectors Computational Intelligence, Lecture 2

by Sergei Savin

Spring 2022

### CONTENT

- Motivating questions
- Four Fundamental Subspaces
- Null space
  - Definition
  - Calculation
- Null space projection
- Closest element from a linear subspace
- Orthogonality, definition
- Projection
- Vectors in Null space, Row space
- Row and Null spaces in linear equations

## MOTIVATING QUESTIONS

You have a linear operator **A**. Try to answer the following questions:

- What are all vectors this operator can produce as outputs? How to find them?
- Are there two inputs that make it produce the same output?
- Are there inputs that produce zero as an output?
- Are there outputs that cannot be produced?
- What is the smallest vector  $\mathbf{x}$  that produces given output  $\mathbf{y}$ ?

These questions are directly related to the idea of fundamental subspaces of a linear operator.

## FOUR FUNDAMENTAL SUBSPACES

One of the key ideas in the linear algebra is that every linear operator has four fundamental subspaces:

- Null space
- Row space
- Column space
- Left null space

Our goal is to understand them. The usefulness of this understating is enormous.

# NULL SPACE Definition

Consider the following task: find all solutions to the system of equations  $\mathbf{A}\mathbf{x} = \mathbf{0}$ .

It can be re-formulated as follows: find all elements of the null space of A.

#### Definition 1

Null space of  $\bf A$  is the set of all vectors  $\bf x$  that  $\bf A$  maps to  $\bf 0$ 

We will denote null space as  $\mathcal{N}(\mathbf{A})$ . In the literature, it is often denoted as  $\ker(\mathbf{A})$  or  $\operatorname{null}(\mathbf{A})$ .

# NULL SPACE Calculation

Now we can find all solutions to the system of equations  $\mathbf{A}\mathbf{x} = \mathbf{0}$  by using functions that generate an orthonormal *basis* in the null space of  $\mathbf{A}$ . In MATLAB it is function null, in Python/Scipy - null\_space:

- $\blacksquare$  N = null(A).
- N = scipy.linalg.null\_space(A).

That is it! Space of solutions of  $\mathbf{A}\mathbf{x} = \mathbf{0}$  is the span of the columns of  $\mathbf{N}$ , and all solutions  $\mathbf{x}^*$  can be represented as  $\mathbf{x}^* = \mathbf{N}\mathbf{z}$ ; for any  $\mathbf{z}$  we get a unique solution, and for any solution - a unique  $\mathbf{z}$ .

### NULL SPACE PROJECTION

### Local coordinates

Let N be the orthonormal basis in the null space of matrix A. Then, if a vector  $\mathbf{x}$  lies in the null space of A, it can be represented as:

$$\mathbf{x} = \mathbf{N}\mathbf{z} \tag{1}$$

where  $\mathbf{z}$  are coordinates of  $\mathbf{x}$  in the basis  $\mathbf{N}$ .

However, there are vector which not only are not lying in the null space of  $\mathbf{A}$ , but the closest vector to them in the null space is zero vector.

## CLOSEST ELEMENT FROM A LINEAR SUBSPACE

Let 
$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$
. Its null space has orthonormal basis  $\mathbf{N} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ .

- $\begin{bmatrix} -2 \\ 0 \end{bmatrix} = -2\mathbf{N}, \begin{bmatrix} 10 \\ 0 \end{bmatrix} = 10\mathbf{N},$  both are in the null space.
- for  $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  the closest vector in the null space is  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ .

## ORTHOGONALITY, DEFINITION

### Definition

If for a vector  $\mathbf{x}$ , the closest vector to it from a linear subspace  $\mathcal{L}$  is zero vector,  $\mathbf{x}$  is called *orthogonal* to the subspace  $\mathcal{L}$ . We denote it as  $\mathbf{x} \in \mathcal{L}^{\perp}$ .

### Definition

The space of all vectors  $\mathbf{x}$ , for which closest vector to them from a linear subspace  $\mathcal{L}$  is zero vector, is called *subspace*, *orthogonal* to  $\mathcal{L}$  and is denoted as  $\mathcal{L}^{\perp}$ .

# PROJECTION Part 1

Let **L** be an orthonormal basis in a linear subspace  $\mathcal{L}$ . Take vector  $\mathbf{a} = \mathbf{x} + \mathbf{y}$ , where  $\mathbf{x}$  lies in the subspace  $\mathcal{L}$ , and  $\mathbf{y}$  is orthogonal to  $\mathcal{L}$ .

#### Definition

We call such vector  $\mathbf{x}$  a projection of  $\mathbf{a}$  onto subspace  $\mathcal{L}$ , and such vector  $\mathbf{y}$  a projection of  $\mathbf{a}$  onto subspace  $\mathcal{L}^{\perp}$ 

Projection of **a** onto  $\mathcal{L}$  can be found as:

$$\mathbf{x} = \mathbf{L}\mathbf{L}^{+}\mathbf{a} \tag{2}$$

Since L is orthonormal, this is the same as  $\mathbf{x} = \mathbf{L}\mathbf{L}^{\mathsf{T}}\mathbf{a}$ 

# PROJECTION Part 2

Since  $\mathbf{a} = \mathbf{x} + \mathbf{y}$ , and  $\mathbf{x} = \mathbf{L}\mathbf{L}^{+}\mathbf{a}$ , we can write:

$$\mathbf{a} = \mathbf{L}\mathbf{L}^{+}\mathbf{a} + \mathbf{y} \tag{3}$$

from which it follows that the projection of  $\mathbf{a}$  onto  $\mathcal{L}^{\perp}$  can be found as:

$$\mathbf{y} = (\mathbf{I} - \mathbf{L}\mathbf{L}^{+})\mathbf{a} \tag{4}$$

where  $\mathbf{I}$  is an identity matrix. Since  $\mathbf{L}$  is orthonormal, this is the same as  $\mathbf{y} = (\mathbf{I} - \mathbf{L} \mathbf{L}^{\top}) \mathbf{a}$ 

# ROW SPACE Definition

### Definition

Let  $\mathcal{N}$  be null space of  $\mathbf{A}$ . Then orthogonal subspace  $\mathcal{N}^{\perp}$  is called *row space* of  $\mathbf{A}$ .

### Definition

Row space of A is the space of all inputs to A that produce non-zero outputs (plus zero vector, which is included in all linear subspaces).

We will denote row space as  $\mathcal{R}$ .

## VECTORS IN NULL SPACE, ROW SPACE

Given vector  $\mathbf{x}$ , matrix  $\mathbf{A}$  and its nulls space basis  $\mathbf{N}$ , and we check if  $\mathbf{x}$  is in the null space of  $\mathbf{A}$ .

We do it by noticing that if it is in the null space of  $\mathbf{A}$ , it will have zero projection onto the row space of  $\mathbf{A}$ . So, the condition is as follows:

$$(\mathbf{I} - \mathbf{N}\mathbf{N}^{+})\mathbf{x} = 0 \tag{5}$$

By the same logic, condition for being in the row space is as follows:

$$\mathbf{N}\mathbf{N}^{+}\mathbf{x} = 0 \tag{6}$$

## ROW AND NULL SPACES IN LINEAR EQUATIONS Part 1

Consider another task: find all solutions to the system of equations  $\mathbf{A}\mathbf{x} = \mathbf{y}$ .

Assume we have two solutions to the system:  $\mathbf{x}_1$  and  $\mathbf{x}_2$ . We know that  $\mathbf{A}\mathbf{x}_1 = \mathbf{A}\mathbf{x}_2 = \mathbf{y}$ , hence  $\mathbf{A}(\mathbf{x}_1 - \mathbf{x}_2) = \mathbf{0}$ . In other words, the difference between any two solutions lies in the null space of  $\mathbf{A}$ .

On the other hand, let  $\mathbf{x}^*$  be a solution, and  $\mathbf{x}^N \in \mathcal{N}(\mathbf{A})$  be a vector in the null space of  $\mathbf{A}$ . Then  $\mathbf{x}^* + \mathbf{x}^N$  is also a solution, since  $\mathbf{A}(\mathbf{x}^* + \mathbf{x}^N) = \mathbf{A}\mathbf{x}^* + \mathbf{A}\mathbf{x}^N = \mathbf{A}\mathbf{x}^* = \mathbf{y}$ .

Therefore, the solution space is given by a single partial solution  $\mathbf{x}^p \notin \mathcal{N}(\mathbf{A})$  and the whole null space of  $\mathbf{A}$ .

## ROW AND NULL SPACES IN LINEAR EQUATIONS Part 2

There are infinitely many ways to chose  $\mathbf{x}^p$ , since if  $\mathbf{x}^p \notin \mathcal{N}(\mathbf{A})$ , then  $(\mathbf{x}^p + \mathbf{x}^N) \notin \mathcal{N}(\mathbf{A})$ , if  $\mathbf{x}^N \in \mathcal{N}(\mathbf{A})$ . However:

#### Statement 1

The smallest-norm  $\mathbf{x}^p$  will lie in the row space of  $\mathbf{A}$ .

We can prove it by observing that there can be only one  $\mathbf{x}^p \in \mathcal{R}(\mathbf{A})$  and adding to it any vector  $\mathbf{x}^N \in \mathcal{N}(\mathbf{A})$  can only increase its magnitude, as  $\mathbf{x}^p$  and  $\mathbf{x}^N$  are orthogonal.

## ROW AND NULL SPACES IN LINEAR EQUATIONS Part 3

If we have  $\mathbf{x}^*$ , which is a solution to  $\mathbf{A}\mathbf{x} = \mathbf{y}$ , we can find the particular solution  $\mathbf{x}^p \in \mathcal{R}(\mathbf{A})$  as a projection:

$$\mathbf{x}^p = (\mathbf{I} - \mathbf{N}\mathbf{N}^+)\mathbf{x}^* \tag{7}$$

where N is the null space basis for A. Alternatively, we can simply find it as:

$$\mathbf{x}^p = \mathbf{A}^+ \mathbf{y} \tag{8}$$

All solutions to Ax = y are then given as:

$$\mathbf{x}^* = \mathbf{A}^+ \mathbf{y} + \mathbf{N} \mathbf{z}, \ \forall \mathbf{z} \tag{9}$$

Lecture slides are available via Moodle.

 $You\ can\ help\ improve\ these\ slides\ at:$  github.com/SergeiSa/Computational-Intelligence-Slides-Spring-2022



Check Moodle for additional links, videos, textbook suggestions.