Synopsis for extractions of unpolarized TMDs: SV19

Alexey A. Vladimirov

These notes contains some information on TMD distributions extracted in the fit SV19, and its later updates. The collection is not complete!, and mainly concernce the latest updates.

Reference: Scimemi:2019cmh

I. Scimemi and A. Vladimirov,

"Non-perturbative structure of semi-inclusive deep-inelastic and Drell-Yan scattering at small transverse momentum,"

JHEP **06** (2020), 137 doi:10.1007/JHEP06(2020)137

Required artemide version: >2.04

For the detailed description of the non-perturbative model see sec.2 of reference. NP RAD (2.88):

$$\mathcal{D}(b,\mu) = \mathcal{D}_{\text{resum}}(b^*(b),\mu) + c_0 b b^*(b), \qquad b^*(b) = \frac{b}{\sqrt{1 + \frac{b^2}{B_{\text{NP}}^2}}}.$$
 (0.1)

NP part of unpolarized TMDPDF:

$$f_{NP}(x,b) = \exp\left(-\frac{\lambda_1 \bar{x} + \lambda_2 x + x \bar{x} \lambda_5}{\sqrt{1 + \lambda_3 x^{\lambda_4} \mathbf{b}^2}} \mathbf{b}^2\right). \tag{0.2}$$

NP part of unpolarized TMDFF:

$$D_{NP}(z,b) = \exp\left(-\frac{\eta_1 z + \eta_2 \bar{z}}{\sqrt{1 + \eta_3 (\mathbf{b}/z)^2}} \frac{\mathbf{b}^2}{z^2}\right) \left(1 + \eta_4 \frac{\mathbf{b}^2}{z^2}\right). \tag{0.3}$$

I. $SV19_NNLO_M=0$

Fit: DY+SIDIS (457+582=1039 points)

Perturbative order: NNLO Number of replicas: 300

Constants-file: const-DY+SIDIS_NNPDF31+DSS_nnlo_m=0


Replica-file: $SV19_nnlo_m=0.rep$


Distinctive feature: Target and production mass corrections in SIDIS kinematics are neglected.


Additional restrictions: Parameter fixed: $B_{NP} = 2 \text{GeV}$.


name	l N	chiL^2/N	chiD^2/N	chi^2/N	sys.shift%
CDF1	l 33	0.538	0.100	0.639	7.095 l
CDF2	l 39	l 1.335 l	0.100	1.336	1.334
D01	l 16	l 0.762 l	0.001	0.768	-1.287
D01	l 8	1.437 I	0.000	1.437	0.000
D02 D02m	l 3	0.421	0.300	0.721	0.293
A7-00 v10	l 5	1.452	0.002	1.454	-0.036 l
A7-10 y 20	l 5	1.432 I 5.522	0.002	5.533	-0.085
A7-20y24	l 5	1.367 I	0.001	1.367	-0.033 -0.010
A8-00 v04	l 5	1.307 2.397	0.000	2.496	2.019
A8-00 y 04 A8-04 y 08	l 5	l 2.586 l	0.464	3.050	2.019
A8-04y08 A8-08y12	l 5	0.826	0.300	1.126	2.016 2.164
A8-12 v16	l 5	1.515	0.268	1.782	2.750
•		l 0.765 l		1.762	1 2.750 I 1 3.456 I
A8-16y20	5		0.287		
A8-20y24	5 3	1.716 0.243	0.908 0.085	2.624	4.143
A8-46Q66	1 3 1 7	0.243 I 0.750 I	0.085	0.328	-0.265
A8-116Q150				0.781	0.278
CMS7	8	1.248	0.000	1.248	0.000
CMS8	8	0.772	0.000	0.772	0.000
LHCb7	8	1.612	0.569	2.181	4.535
LHCb8	7	2.773	1.052	3.825	4.453
LHCb13	9	0.607	0.185	0.792	5.044
PHE200] 3	0.284	0.000	0.284	-0.250
E228-200	43	1.034	0.043	1.077	33.958
E228-300	53	1.005	0.025	1.029	28.507
E228-400	76	0.834	0.008	0.842	19.414
E772	35	1.895	0.015	1.910	7.230
E605	53	0.513	0.034	0.547	20.057
					-
Total	457	1.096	0.079	1.175	

name	N	1	chiL^2/N		chiD^2/N		chi^2/N	sys.shift%	1
		- ا		- ا		· -			i –
hermes.p.vmsub.zxpt.pi+	24	1	2.371	1	0.237		2.607	1.983	1
hermes.p.vmsub.zxpt.pi-	24	1	1.277	1	0.044		1.321	0.853	ĺ
hermes.d.vmsub.zxpt.pi+	24	1	0.680	1	0.023		0.703	0.683	1
hermes.d.vmsub.zxpt.pi-	24	1	0.880	1	0.029		0.909	0.772	ĺ
hermes.p.vmsub.zxpt.k+	24	1	0.630	1	0.000		0.630	-0.039	ĺ
hermes.p.vmsub.zxpt.k-	24	1	0.627	1	0.000		0.627	0.061	ĺ
hermes.d.vmsub.zxpt.k+	24	1	0.474	1	0.000		0.474	0.038	ĺ
hermes.d.vmsub.zxpt.k-	24	1	1.214	1	0.001		1.215	0.159	ĺ
compass.d.h+	195	1	0.686	1	0.005		0.691	1.148	ĺ
compass.d.h-	195	1	0.817	1	0.064		0.881	-4.102	ĺ
		- ا		- ا		- -			I –
Total	582	1	0.840	1	0.037		0.877		









Parameter vectors:

$$\mathbf{Mean} = \{2., 0.0399, 0.186, 7.64, 492., 2.23, -2.49, 0., 0., 0.187, 0.476, 0.500, 0.428\} \tag{1.1}$$

$$\mathbf{std} = \{0., 0.0036, 0.0088, 0.52, 70., 0.087, 0.74, 0., 0.016, 0.015, 0.032, 0.027\} \tag{1.2}$$

Distribution of parameters over replicas

