

# **Part XI.**

# **Properties of Regular Languages**

# Pumping Lemma for RLs

**Gist: Pumping lemma demonstrates an infinite iteration of some substring in RLs.**

- Let  $L$  be a RL. Then, there is  $k \geq 1$  such that if  $z \in L$  and  $|z| \geq k$ , then there exist  $u, v, w: z = uvw$ ,  
1)  $v \neq \varepsilon$  2)  $|uv| \leq k$  3) for each  $m \geq 0$ ,  $uv^m w \in L$

**Example:** for RE  $r = ab^*c$ ,  $L(r)$  is *regular*.

There is  $k = 3$  such that 1), 2) and 3) holds.

# Pumping Lemma for RLs

**Gist: Pumping lemma demonstrates an infinite iteration of some substring in RLs.**

- Let  $L$  be a RL. Then, there is  $k \geq 1$  such that if  $z \in L$  and  $|z| \geq k$ , then there exist  $u, v, w: z = uvw$ ,  
 1)  $v \neq \varepsilon$  2)  $|uv| \leq k$  3) for each  $m \geq 0$ ,  $uv^m w \in L$

**Example:** for RE  $r = ab^*c$ ,  $L(r)$  is *regular*.

There is  $k = 3$  such that 1), 2) and 3) holds.

- for  $z = abc$ :  $z \in L(r)$  &  $|z| \geq 3$ :
 

$$\begin{array}{ccc} a & b & c \\ \downarrow & \downarrow & \downarrow \\ u & v & w \end{array}$$
 $v \neq \varepsilon, |uv| = 2 \leq 3$

$$\begin{aligned} uv^0w &= ab^0c = ac \in L(r) \\ uv^1w &= ab^1c = abc \in L(r) \\ uv^2w &= ab^2c = abbc \in L(r) \\ &\vdots \end{aligned}$$

# Pumping Lemma for RLs

**Gist: Pumping lemma demonstrates an infinite iteration of some substring in RLs.**

- Let  $L$  be a RL. Then, there is  $k \geq 1$  such that if  $z \in L$  and  $|z| \geq k$ , then there exist  $u, v, w: z = uvw$ ,  
 1)  $v \neq \varepsilon$  2)  $|uv| \leq k$  3) for each  $m \geq 0$ ,  $uv^m w \in L$

**Example:** for RE  $r = ab^*c$ ,  $L(r)$  is *regular*.

There is  $k = 3$  such that 1), 2) and 3) holds.

- for  $z = abc$ :  $z \in L(r)$  &  $|z| \geq 3$ :
 

$$\begin{array}{ccc} a & b & c \\ \downarrow & \downarrow & \downarrow \\ u & v & w \end{array}$$
 $v \neq \varepsilon, |uv| = 2 \leq 3$

$uv^0w = ab^0c = ac \in L(r)$   
 $uv^1w = ab^1c = abc \in L(r)$   
 $uv^2w = ab^2c = abbc \in L(r)$   
 $\vdots$
- for  $z = abbc$ :  $z \in L(r)$  &  $|z| \geq 3$ :
 

$$\begin{array}{ccc} a & b & b & c \\ \downarrow & \downarrow & \downarrow & \downarrow \\ u & v & w & \end{array}$$
 $\vdots$ 
 $v \neq \varepsilon, |uv| = 2 \leq 3$

$uv^0w = ab^0bc = abc \in L(r)$   
 $uv^1w = ab^1bc = abbc \in L(r)$   
 $uv^2w = ab^2bc = abbbc \in L(r)$   
 $\vdots$

# Pumping Lemma: Illustration

- $L$  = any regular language:
-

# Pumping Lemma: Illustration

- $L =$  any regular language:


---

  $z \in L$   $\rightarrow$  *nothing interesting*

---



# Pumping Lemma: Illustration

- $L =$  any regular language:



# Pumping Lemma: Illustration



- $L =$  any regular language:







# Pumping Lemma: Illustration

- $L =$  any regular language:





# Pumping Lemma: Illustration

- $L =$  any regular language:



# Pumping Lemma: Illustration

- $L =$  any regular language:




- 1)  $v \neq \epsilon$
- 2)  $|v| \leq k$



...

# Proof of Pumping Lemma 1/3

- Let  $L$  be a regular language. Then, there exists **DFA**  $M = (Q, \Sigma, R, s, F)$ , and  $L = L(M)$ .
- For  $z \in L(M)$ ,  $M$  makes  $|z|$  moves and  $M$  visits  $|z| + 1$  states:
- for  $z = a_1 a_2 \dots a_n$ :




## Proof of Pumping Lemma 2/3

- Let  $k = \text{card}(Q)$  (the number of states).

For each  $z \in L$  and  $|z| \geq k$ ,  $M$  visits  $k + 1$  or more states. As  $k + 1 > \text{card}(Q)$ , there exists a state  $q$  that  $M$  visits at least twice.

- For  $z$  exist  $u, v, w$  such that  $z = uvw$ :




# Proof of Pumping Lemma 2/3

- Let  $k = \text{card}(Q)$  (the number of states).

For each  $z \in L$  and  $|z| \geq k$ ,  $M$  visits  $k + 1$  or more states. As  $k + 1 > \text{card}(Q)$ , there exists a state  $q$  that  $M$  visits at least twice.

- For  $z$  exist  $u, v, w$  such that  $z = uvw$ :




# Proof of Pumping Lemma 2/3

- Let  $k = \text{card}(Q)$  (the number of states).

For each  $z \in L$  and  $|z| \geq k$ ,  $M$  visits  $k + 1$  or more states. As  $k + 1 > \text{card}(Q)$ , there exists a state  $q$  that  $M$  visits at least twice.

- For  $z$  exist  $u, v, w$  such that  $z = uvw$ :




# Proof of Pumping Lemma 2/3

- Let  $k = \text{card}(Q)$  (the number of states).

For each  $z \in L$  and  $|z| \geq k$ ,  $M$  visits  $k + 1$  or more states. As  $k + 1 > \text{card}(Q)$ , there exists a state  $q$  that  $M$  visits at least twice.

- For  $z$  exist  $u, v, w$  such that  $z = uvw$ :






# Proof of Pumping Lemma 2/3

- Let  $k = \text{card}(Q)$  (the number of states).

For each  $z \in L$  and  $|z| \geq k$ ,  $M$  visits  $k + 1$  or more states. As  $k + 1 > \text{card}(Q)$ , there exists a state  $q$  that  $M$  visits at least twice.

- For  $z$  exist  $u, v, w$  such that  $z = uvw$ :



Summary:

$$sz = suvw \vdash^i qvw \vdash^j qw \vdash^* f, f \in F$$

# Proof of Pumping Lemma 3/3

- There exist moves:

①  $su \vdash^i q$ ;    ②  $qv \vdash^j q$ ;    ③  $qw \vdash^* f, f \in F$ , so

# Proof of Pumping Lemma 3/3

- There exist moves:
  - ①  $su \vdash^i q$ ;    ②  $qv \vdash^j q$ ;    ③  $qw \vdash^* f, f \in F$ , so
- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

$suw$

# Proof of Pumping Lemma 3/3

- There exist moves:

①  $su \vdash^i q$ ;    ②  $qv \vdash^j q$ ;    ③  $qw \vdash^* f, f \in F$ , so

- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

①  
 $suw \vdash^i qw$

# Proof of Pumping Lemma 3/3

- There exist moves:

①.  $su \vdash^i q$ ;    ②.  $qv \vdash^j q$ ;    ③.  $qw \vdash^* f, f \in F$ , so

- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

①.  $suw \vdash^i qw \vdash^* f, f \in F$

# Proof of Pumping Lemma 3/3

- There exist moves:

①.  $su \vdash^i q$ ;    ②.  $qv \vdash^j q$ ;    ③.  $qw \vdash^* f, f \in F$ , so

- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

①.  $suw \vdash^i qw$     ③.  $qw \vdash^* f, f \in F$

- for each  $m > 0$ ,

$su v^m w$

# Proof of Pumping Lemma 3/3

- There exist moves:

①  $su \vdash^i q$ ;    ②  $qv \vdash^j q$ ;    ③  $qw \vdash^* f, f \in F$ , so

- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

①  $suw \vdash^i qw \vdash^* f, f \in F$

- for each  $m > 0$ ,

①  $su v^m w \vdash^i q v^m w$

# Proof of Pumping Lemma 3/3

- There exist moves:

①  $su \vdash^i q$ ;    ②  $qv \vdash^j q$ ;    ③  $qw \vdash^* f, f \in F$ , so

- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

①  $suw \vdash^i qw \vdash^* f, f \in F$

- for each  $m > 0$ ,

①  $su v^m w \vdash^i q v^m w \vdash^j q v^{m-1} w \vdash^j \dots \vdash^j q w$



# Proof of Pumping Lemma 3/3

- There exist moves:

①  $su \vdash^i q$ ;    ②  $qv \vdash^j q$ ;    ③  $qw \vdash^* f, f \in F$ , so

- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

①  $suw \vdash^i qw \vdash^* f, f \in F$

- for each  $m > 0$ ,

①  $su v^m w \vdash^i q v^m w \vdash^j q v^{m-1} w \vdash^j \dots \vdash^j qw \vdash^* f, f \in F$

# Proof of Pumping Lemma 3/3

- There exist moves:

$$\textcircled{1}. su \vdash^i q; \quad \textcircled{2}. qv \vdash^j q; \quad \textcircled{3}. qw \vdash^* f, f \in F, \text{ so}$$

- for  $m = 0$ ,  $uv^m w = uv^0 w = uw$ ,

$$\textcircled{1}. suw \vdash^i qw \vdash^* f, f \in F$$

- for each  $m > 0$ ,

$$\textcircled{1}. suv^m w \vdash^i \textcircled{2}. qv^m w \vdash^j \textcircled{2}. qv^{m-1} w \vdash^j \dots \vdash^j \textcircled{3}. qw \vdash^* f, f \in F$$

## Summary:

1)  $qv \vdash^j q, j \geq 1$ ; therefore,  $|v| \geq 1$ , so  $v \neq \varepsilon$

2)  $su \vdash^i qv \vdash^j q, i + j \leq k$ ; therefore,  $|uv| \leq k$

3) For each  $m \geq 0$ :  $su v^m w \vdash^* f, f \in F$ , therefore  $u v^m w \in L$

***QED***

# Pumping Lemma: Application I

- Based on the pumping lemma, we often make a proof by contradiction to demonstrate that a language is **not** regular

# Pumping Lemma: Application I

- Based on the pumping lemma, we often make a proof by contradiction to demonstrate that a language is **not** regular

Assume that  $L$  is regular

# Pumping Lemma: Application I

- Based on the pumping lemma, we often make a proof by contradiction to demonstrate that a language is **not** regular

Assume that  $L$  is regular



Consider the PL constant  $k$  and select  $z \in L$ , whose length depends on  $k$  so  $|z| \geq k$  is surely true.

# Pumping Lemma: Application I

- Based on the pumping lemma, we often make a proof by contradiction to demonstrate that a language is **not** regular

Assume that  $L$  is regular

Consider the PL constant  $k$  and select  $z \in L$ , whose length depends on  $k$  so  $|z| \geq k$  is surely true.

For all decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon$ ,  $|uv| \leq k$ , show:  
 there exists  $m \geq 0$  such that  $uv^m w \notin L$   
 from the pumping lemma,  $uv^m w \in L$  } **contradiction**

# Pumping Lemma: Application I

- Based on the pumping lemma, we often make a proof by contradiction to demonstrate that a language is **not** regular

Assume that  $L$  is regular

Consider the PL constant  $k$  and select  $z \in L$ , whose length depends on  $k$  so  $|z| \geq k$  is surely true.

For all decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon$ ,  $|uv| \leq k$ , show:  
 there exists  $m \geq 0$  such that  $uv^m w \notin L$  } **contradiction**  
 from the pumping lemma,  $uv^m w \in L$

**false assumption**

# Pumping Lemma: Application I

- Based on the pumping lemma, we often make a proof by contradiction to demonstrate that a language is **not** regular

Assume that  $L$  is regular

Consider the PL constant  $k$  and select  $z \in L$ , whose length depends on  $k$  so  $|z| \geq k$  is surely true.

For all decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon$ ,  $|uv| \leq k$ , show:  
 there exists  $m \geq 0$  such that  $uv^m w \notin L$  } **contradiction**  
 from the pumping lemma,  $uv^m w \in L$

**false assumption**




Therefore,  
 **$L$  is not regular**



# Pumping Lemma: Example

Prove that  $L = \{a^n b^n : n \geq 0\}$  is not regular:


- 1) Assume that  $L$  is regular. Let  $k \geq 1$  be the pumping lemma constant for  $L$ .
- 2) Let  $z = a^k b^k$ :  $a^k b^k \in L$ ,  $|z| = |a^k b^k| = 2k \geq k$
- 3) All decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon$ ,  $|uv| \leq k$ :



# Pumping Lemma: Example

Prove that  $L = \{a^n b^n : n \geq 0\}$  is not regular:

- 1) Assume that  $L$  is regular. Let  $k \geq 1$  be the pumping lemma constant for  $L$ .
- 2) Let  $z = a^k b^k$ :  $a^k b^k \in L$ ,  $|z| = |a^k b^k| = 2k \geq k$
- 3) All decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon$ ,  $|uv| \leq k$ :




• pumping lemma:  $uv^0w \in L$

# Pumping Lemma: Example

Prove that  $L = \{a^n b^n : n \geq 0\}$  is not regular:



- 1) Assume that  $L$  is regular. Let  $k \geq 1$  be the pumping lemma constant for  $L$ .
- 2) Let  $z = a^k b^k : a^k b^k \in L, |z| = |a^k b^k| = 2k \geq k$
- 3) All decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon, |uv| \leq k$ :



# Pumping Lemma: Example

Prove that  $L = \{a^n b^n : n \geq 0\}$  is not regular:

- 1) Assume that  $L$  is regular. Let  $k \geq 1$  be the pumping lemma constant for  $L$ .
- 2) Let  $z = a^k b^k : a^k b^k \in L, |z| = |a^k b^k| = 2k \geq k$
- 3) All decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon, |uv| \leq k$ :




**Contradiction!**

# Pumping Lemma: Example

Prove that  $L = \{a^n b^n : n \geq 0\}$  is not regular:

- 1) Assume that  $L$  is regular. Let  $k \geq 1$  be the pumping lemma constant for  $L$ .
- 2) Let  $z = a^k b^k : a^k b^k \in L, |z| = |a^k b^k| = 2k \geq k$
- 3) All decompositions of  $z$  into  $uvw$ ,  $v \neq \varepsilon, |uv| \leq k$ :



- 4) Therefore,  $L$  is not regular

# Note on Use of Pumping Lemma

- **Pumping lemma:**

if  $L$  is regular then  $\Rightarrow$  exist  $k \geq 1$  and ...

## Main application of the pumping lemma:

- proof by contradiction that  $L$  is **not** regular.
-

# Note on Use of Pumping Lemma

- **Pumping lemma:**

if  $L$  is regular  $\Rightarrow$  exist  $k \geq 1$  and ...

**Main application of the pumping lemma:**

- proof by contradiction that  $L$  is not regular.

- **However, the next implication is incorrect:**

~~if exist  $k \geq 1$  and ...  $\Rightarrow$   $L$  is regular~~

- We **cannot** use the pumping lemma to prove that  $L$  is regular.

## Pumping Lemma: Application II. 1/3

- We can use the pumping lemma to prove some other theorems.
- 

### Illustration:

- Let  $M$  be a DFA and  $k$  be the pumping lemma constant ( $k$  is the number of states in  $M$ ). Then,  $L(M)$  is infinite  $\Leftrightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

### Proof:

- 1) there exists  $z \in L(M)$ ,  $k \leq |z| < 2k \Rightarrow L(M)$  is infinite:



## Pumping Lemma: Application II. 1/3

- We can use the pumping lemma to prove some other theorems.

### Illustration:

- Let  $M$  be a DFA and  $k$  be the pumping lemma constant ( $k$  is the number of states in  $M$ ). Then,  $L(M)$  is infinite  $\Leftrightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

### Proof:

1) there exists  $z \in L(M)$ ,  $k \leq |z| < 2k \Rightarrow L(M)$  is infinite:

if  $z \in L(M)$ ,  $k \leq |z|$ , then by PL:

$z = uvw$ ,  $v \neq \varepsilon$ , and for each  $m \geq 0$ :  $uv^mw \in L(M)$

# Pumping Lemma: Application II. 1/3

- We can use the pumping lemma to prove some other theorems.

## Illustration:

- Let  $M$  be a DFA and  $k$  be the pumping lemma constant ( $k$  is the number of states in  $M$ ). Then,  $L(M)$  is infinite  $\Leftrightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

## Proof:

1) there exists  $z \in L(M)$ ,  $k \leq |z| < 2k \Rightarrow L(M)$  is infinite:

if  $z \in L(M)$ ,  $k \leq |z|$ , then by PL:

$z = uvw$ ,  $v \neq \varepsilon$ , and for each  $m \geq 0$ :  $uv^m w \in L(M)$

$L(M)$  is infinite

# Pumping Lemma: Application II. 2/3

2)  $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$ :

- We prove by contradiction, that

$L(M)$  is infinite  $\xrightarrow{\text{a)}}$  there exists  $z \in L(M)$ ,  $|z| \geq k$

$\downarrow \text{b)}$

there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

---

**a)** Prove by contradiction that

- **$L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $|z| \geq k$**

# Pumping Lemma: Application II. 2/3

2)  $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$ :

- We prove by contradiction, that

$L(M)$  is infinite  $\xrightarrow{\text{a)}}$  there exists  $z \in L(M)$ ,  $|z| \geq k$

$\downarrow \text{b)}$

there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

---

**a)** Prove by contradiction that

- **$L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $|z| \geq k$**

Assume that  **$L(M)$  is infinite** and there exists no  $z \in L(M)$ ,  $|z| \geq k$

# Pumping Lemma: Application II. 2/3

2)  $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$ :

- We prove by contradiction, that

$L(M)$  is infinite  $\xrightarrow{\text{a)}}$  there exists  $z \in L(M)$ ,  $|z| \geq k$

$\downarrow \text{b)}$

there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

---

**a)** Prove by contradiction that

- $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $|z| \geq k$

Assume that  $L(M)$  is infinite and there exists no  $z \in L(M)$ ,  $|z| \geq k$

for all  $z \in L(M)$  holds  $|z| < k$

# Pumping Lemma: Application II. 2/3

2)  $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$ :

- We prove by contradiction, that

$L(M)$  is infinite  $\xrightarrow{\text{a)}}$  there exists  $z \in L(M)$ ,  $|z| \geq k$

$\downarrow \text{b)}$

there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

---

**a)** Prove by contradiction that

- $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $|z| \geq k$

Assume that  $L(M)$  is infinite and there exists no  $z \in L(M)$ ,  $|z| \geq k$

for all  $z \in L(M)$  holds  $|z| < k$

$\downarrow$   
 $L(M)$  is finite

# Pumping Lemma: Application II. 2/3

2)  $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$ :

- We prove by contradiction, that

$L(M)$  is infinite  $\xrightarrow{\text{a)}}$  there exists  $z \in L(M)$ ,  $|z| \geq k$

$\downarrow \text{b)}$

there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

**a)** Prove by contradiction that

- $L(M)$  is infinite  $\Rightarrow$  there exists  $z \in L(M)$ ,  $|z| \geq k$

Assume that  $L(M)$  is infinite and there exists no  $z \in L(M)$ ,  $|z| \geq k$

for all  $z \in L(M)$  holds  $|z| < k$

**Contradiction !**

$L(M)$  is finite

# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**




# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**

Assume that **there is  $z \in L(M)$ ,  $|z| \geq k$**




# Pumping Lemma: Application II. 3/3

**b) Prove by contradiction**

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**

Assume that **there is**  $z \in L(M)$ ,  $|z| \geq k$   
and **there is no**  $z \in L(M)$ ,  $k \leq |z| < 2k$




# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists**  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
**there exists**  $z \in L(M)$ ,  $k \leq |z| < 2k$

Assume that **there is**  $z \in L(M)$ ,  $|z| \geq k$   
 and **there is no**  $z \in L(M)$ ,  $k \leq |z| < 2k$



Let  $z_0$  be **the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$


Because there exists no  $z \in L(M)$ ,  $k \leq |z| < 2k$ , so  $|z_0| \geq 2k$

# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**

Assume that **there is  $z \in L(M)$ ,  $|z| \geq k$**   
and **there is no  $z \in L(M)$ ,  $k \leq |z| < 2k$**



Let  $z_0$  be **the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$

Because there exists no  $z \in L(M)$ ,  $k \leq |z| < 2k$ , so  $|z_0| \geq 2k$

If  $z_0 \in L(M)$  and  $|z_0| \geq k$ , the PL implies:  $z_0 = uvw$ ,


$|uv| \leq k$ , and for each  $m \geq 0$ ,  $uv^m w \in L(M)$

# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**

Assume that **there is  $z \in L(M)$ ,  $|z| \geq k$**   
and **there is no  $z \in L(M)$ ,  $k \leq |z| < 2k$**



Let  $z_0$  be **the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$

Because there exists no  $z \in L(M)$ ,  $k \leq |z| < 2k$ , so  $|z_0| \geq 2k$

If  $z_0 \in L(M)$  and  $|z_0| \geq k$ , the PL implies:  $z_0 = uvw$ ,

$|uv| \leq k$ , and for each  $m \geq 0$ ,  $uv^m w \in L(M)$


$$|uw| = \underbrace{|z_0|}_{\geq 2k} - \underbrace{|v|}_{\leq k} \geq k$$

# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**

Assume that **there is  $z \in L(M)$ ,  $|z| \geq k$**   
and **there is no  $z \in L(M)$ ,  $k \leq |z| < 2k$**



Let  $z_0$  be **the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$

Because there exists no  $z \in L(M)$ ,  $k \leq |z| < 2k$ , so  $|z_0| \geq 2k$

If  $z_0 \in L(M)$  and  $|z_0| \geq k$ , the PL implies:  $z_0 = uvw$ ,

$|uv| \leq k$ , and for each  $m \geq 0$ ,  $uv^m w \in L(M)$

$$|uw| = \underbrace{|z_0|}_{\geq 2k} - \underbrace{|v|}_{\leq k} \geq k$$


↓  
for  $m = 0$ :  $uv^m w = uw \in L(M)$

# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**

Assume that **there is  $z \in L(M)$ ,  $|z| \geq k$**   
and **there is no  $z \in L(M)$ ,  $k \leq |z| < 2k$**



Let  $z_0$  be **the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$

Because there exists no  $z \in L(M)$ ,  $k \leq |z| < 2k$ , so  $|z_0| \geq 2k$

If  $z_0 \in L(M)$  and  $|z_0| \geq k$ , the PL implies:  $z_0 = uvw$ ,

$|uv| \leq k$ , and for each  $m \geq 0$ ,  $uv^m w \in L(M)$

$$|uw| = \underbrace{|z_0|}_{\geq 2k} - \underbrace{|v|}_{\leq k} \geq k \quad \text{for } m = 0: uv^m w = uw \in L(M)$$


**Summary:**  $uw \in L(M)$ ,  $|uw| \geq k$  and  $|uw| < |z_0|$ !

# Pumping Lemma: Application II. 3/3

b) Prove by contradiction

- there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$

Assume that **there is**  $z \in L(M)$ ,  $|z| \geq k$   
and **there is no**  $z \in L(M)$ ,  $k \leq |z| < 2k$



Let  $z_0$  be **the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$

Because there exists no  $z \in L(M)$ ,  $k \leq |z| < 2k$ , so  $|z_0| \geq 2k$

If  $z_0 \in L(M)$  and  $|z_0| \geq k$ , the PL implies:  $z_0 = uvw$ ,

$|uv| \leq k$ , and for each  $m \geq 0$ ,  $uv^m w \in L(M)$

$$|uw| = \underbrace{|z_0|}_{\geq 2k} - \underbrace{|v|}_{\leq k} \geq k \quad \downarrow \quad \text{for } m = 0: uv^m w = uw \in L(M)$$

**Summary:**  $uw \in L(M)$ ,  $|uw| \geq k$  and  $|uw| < |z_0|$ !

**$z_0$  is not the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$



# Pumping Lemma: Application II. 3/3

**b)** Prove by contradiction

- **there exists  $z \in L(M)$ ,  $|z| \geq k \Rightarrow$   
there exists  $z \in L(M)$ ,  $k \leq |z| < 2k$**

Assume that **there is  $z \in L(M)$ ,  $|z| \geq k$**   
and **there is no  $z \in L(M)$ ,  $k \leq |z| < 2k$**



Let  $z_0$  be **the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$

Because there exists no  $z \in L(M)$ ,  $k \leq |z| < 2k$ , so  $|z_0| \geq 2k$

If  $z_0 \in L(M)$  and  $|z_0| \geq k$ , the PL implies:  $z_0 = uvw$ ,

$|uv| \leq k$ , and for each  $m \geq 0$ ,  $uv^m w \in L(M)$

$$|uw| = \underbrace{|z_0|}_{\geq 2k} - \underbrace{|v|}_{\leq k} \geq k$$

for  $m = 0$ :  $uv^m w = uw \in L(M)$

**Summary:**  $uw \in L(M)$ ,  $|uw| \geq k$  and  $|uw| < |z_0|$ !

**$z_0$  is not the shortest string** satisfying  $z_0 \in L(M)$ ,  $|z_0| \geq k$

**Contradiction !**

## Closure properties 1/2


**Definition:** The family of regular languages is closed under an operation  $\circ$  if the language resulting from the application of  $\circ$  to **any** regular languages is also regular.

## Closure properties 1/2

**Definition:** The family of regular languages is closed under an operation  $\circ$  if the language resulting from the application of  $\circ$  to **any** regular languages is also regular.

### Illustration:

- The family of regular languages is closed under *union*.  
It means:




The family of  
regular languages

# Closure properties 1/2

**Definition:** The family of regular languages is closed under an operation  $\circ$  if the language resulting from the application of  $\circ$  to **any** regular languages is also regular.

## Illustration:

- The family of regular languages is closed under *union*.  
It means:




# Closure properties 1/2

**Definition:** The family of regular languages is closed under an operation  $\circ$  if the language resulting from the application of  $\circ$  to **any** regular languages is also regular.

## Illustration:

- The family of regular languages is closed under *union*.  
It means:



## Closure properties 2/2

**Theorem:** The family of regular languages is closed under **union**, **concatenation**, **iteration**.

### Proof:

- Let  $L_1, L_2$  be two **regular languages**
- Then, there exist two REs  $r_1, r_2$ :  $L(r_1) = L_1, L(r_2) = L_2$ ;
- By the definition of regular expressions:
  - $r_1.r_2$  is a RE denoting  $L_1 L_2$
  - $r_1 + r_2$  is a RE denoting  $L_1 \cup L_2$
  - $r_1^*$  is a RE denoting  $L_1^*$
- Every RE denotes regular language, so  $L_1 L_2, L_1 \cup L_2, L_1^*$  are a **regular languages**

# Algorithm: FA for Complement

- **Input:** Complete FA:  $M = (Q, \Sigma, R, s, F)$
- **Output:** Complete FA:  $M' = (Q, \Sigma, R, s, F')$ ,  


$$L(M') = \overline{L(M)}$$

---

## • Method:

- $F' := Q - F$
- 

**Example:**



# Algorithm: FA for Complement

- **Input:** Complete FA:  $M = (Q, \Sigma, R, s, F)$
- **Output:** Complete FA:  $M' = (Q, \Sigma, R, s, F')$ ,  


$$L(M') = \overline{L(M)}$$

---

## • Method:

- $F' := Q - F$
- 

**Example:**





# Algorithm: FA for Complement


- **Input:** Complete FA:  $M = (Q, \Sigma, R, s, F)$
- **Output:** Complete FA:  $M' = (Q, \Sigma, R, s, F')$ ,  

$$L(M') = \overline{L(M)}$$

## • Method:

- $F' := Q - F$

## Example:




$L(M) = \{x: ab \text{ is a substring of } x\}; L(M') = \{x: ab \text{ is no substring of } x\}$

# FA for Complement: Problem

- Previous algorithm requires a **complete** FA
- If  $M$  is incomplete FA, then  $M$  must be converted to a complete FA before we use the previous algorithm

## Example:

Incomplete DFA:




# FA for Complement: Problem

- Previous algorithm requires a **complete** FA
- If  $M$  is incomplete FA, then  $M$  must be converted to a complete FA before we use the previous algorithm

## Example:

Incomplete DFA:

$$L(M_1') \neq \overline{L(M)}! - c \notin L(M), c \notin L(M_1')$$




# FA for Complement: Problem


- Previous algorithm requires a **complete** FA
- If  $M$  is incomplete FA, then  $M$  must be converted to a complete FA before we use the previous algorithm

## Example:


Incomplete DFA:



$$L(M_1') \neq \overline{L(M)}! - c \notin L(M), c \notin L(M_1')$$



Complete DFA:






# FA for Complement: Problem

- Previous algorithm requires a **complete** FA
- If  $M$  is incomplete FA, then  $M$  must be converted to a complete FA before we use the previous algorithm

## Example:




Incomplete DFA:

$$L(M_1') \neq \overline{L(M)}! - c \notin L(M), c \notin L(M_1')$$



$$L(M_2') = \overline{L(M)}$$

Complete DFA:



# Closure properties: Complement

**Theorem:** The family of regular languages is closed under **complement**.

## Proof:

- Let  $L$  be a **regular language**
- Then, there exists a complete DFA  $M$ :  $L(M) = L$
- We can construct a complete DFA  $M'$ :  $L(M') = \overline{L}$  by using the previous algorithm
- Every FA defines a regular language, so  $\overline{L}$  is a **regular language**

# Closure properties: Intersection

**Theorem:** The family of regular languages is closed under **intersection**.

## Proof:

- Let  $L_1, L_2$  be two **regular languages**
- $\overline{L_1}, \overline{L_2}$  are **regular languages**  
(the family of regular languages is closed under complement)
- $\overline{L_1} \cup \overline{L_2}$  is a **regular language**  
(the family of regular languages is closed under union)
- $\overline{\overline{L_1} \cup \overline{L_2}}$  is a **regular language**  
(the family of regular languages is closed under complement)
- $L_1 \cap L_2 = \overline{\overline{L_1} \cup \overline{L_2}}$  is a **regular language** (DeMorgan's law)

# Boolean Algebra of Languages

**Definition:** Let a family of languages be closed under union, intersection, and complement. Then, this family represents a *Boolean algebra of languages*.

**Theorem:** The family of regular languages is a Boolean algebra of languages.

**Proof:**

- The family of regular languages is closed under union, intersection, and complement.




## Minimization: Distinguishable States


**Gist:** String  $w$  *distinguishes* states  $p$  and  $q$  if WSFA reaches a final state from precisely one of configurations  $pw$  and  $qw$ .

**Definition:** Let  $M = (Q, \Sigma, R, s, F)$  be a WSFA, and let  $p, q \in Q, p \neq q$ . States  $p$  and  $q$  are *distinguishable* if there exists  $w \in \Sigma^*$  such that:  $pw \vdash^* p'$  and  $qw \vdash^* q'$ , where  $p', q' \in Q$  and  $((p' \in F \text{ and } q' \notin F) \text{ or } (p' \notin F \text{ and } q' \in F))$ ; otherwise, states  $p$  and  $q$  are *indistinguishable*.

# Distinguishable States: Example



# Distinguishable States: Example




- $s$  and  $q_1$  are distinguishable, because for  $w = a$ :

$$sa \models s, s \notin F$$

$$q_1a \models q_2, q_2 \in F$$

# Distinguishable States: Example




- $s$  and  $q_1$  are **distinguishable**, because for  $w = a$ :

$$\begin{aligned} sa &\vdash s, s \notin F \\ q_1 a &\vdash q_2, q_2 \in F \end{aligned}$$

- $q_2$  and  $q_3$  are **indistinguishable**, because for each  $w \in \Sigma^*$ :

$$\begin{aligned} q_2 w &\vdash^* q_2, q_2 \in F \\ q_3 w &\vdash^* q_3, q_3 \in F \end{aligned}$$

# Distinguishable States: Example



- $s$  and  $q_1$  are **distinguishable**, because for  $w = a$ :

$$\begin{aligned} sa &\vdash s, s \notin F \\ q_1 a &\vdash q_2, q_2 \in F \end{aligned}$$

- $q_2$  and  $q_3$  are **indistinguishable**, because for each  $w \in \Sigma^*$ :

$$\begin{aligned} q_2 w &\vdash^* q_2, q_2 \in F \\ q_3 w &\vdash^* q_3, q_3 \in F \end{aligned}$$

- Other pairs of states are trivially **distinguishable** for  $w = \varepsilon$ .

## Minimum-State FA

**Definition:** Let  $M$  be a WSFA. Then,  $M$  is *minimum-state FA* if  $M$  contains only distinguishable states.

**Theorem:** For every WSFA  $M$ , there is an equivalent minimum-state FA  $M_m$

**Proof:** Use the next algorithm.


# Algorithm: WSFA to Min-State FA

- **Input:** WSFA  $M = (Q, \Sigma, R, s, F)$
- **Output:** Minimum-State FA  $M_m = (Q_m, \Sigma, R_m, s_m, F_m)$
- **Method:**
  - $Q_m = \{\{p: p \in F\}, \{q: q \in Q - F\}\};$
  - **repeat**
    - if there exist**  $X \in Q_m, d \in \Sigma, X_1, X_2 \subset X$  such that
 
$$X = X_1 \cup X_2, X_1 \cap X_2 = \emptyset \text{ and}$$

$$\{q_1: p_1 \in X_1, p_1 d \rightarrow q_1 \in R\} \subseteq Q_1, Q_1 \in Q_m,$$


$$\{q_2: p_2 \in X_2, p_2 d \rightarrow q_2 \in R\} \cap Q_1 = \emptyset$$
    - then** divide  $X$  into  $X_1$  and  $X_2$  in  $Q_m$
  - until** no division is possible;
  - $R_m = \{Xa \rightarrow Y: X, Y \in Q_m, pa \rightarrow q \in R, p \in X, q \in Y, a \in \Sigma\};$
  - $s_m = X$  with  $s \in X; F_m := \{X: X \in Q_m, X \cap F \neq \emptyset\}.$

# Minimization: Example 1/4






# Minimization: Example 1/4




# Minimization: Example 1/4



1)  $X = \{s, f\}$ :

$d = a$ :  $sa \rightarrow f$   
 $fa \rightarrow s$

# Minimization: Example 1/4




1)  $X = \{s, f\}$ : From one set

$d = a$ :

$sa$	$\rightarrow$	$f$
$fa$	$\rightarrow$	$s$


# Minimization: Example 1/4



1)  $X = \{s, f\}$ : From one set

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$  (From one set)  
 $d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

# Minimization: Example 1/4




1)  $X = \{s, f\}$ : **From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

**From one set**

# Minimization: Example 1/4



1)  $X = \{s, f\}$ : **From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$


**From one set**

2)  $X = \{q_1, q_2, q_3, q_4\}$ :


$d = a$ :

- $q_1 a \rightarrow q_1$
- $q_2 a \rightarrow q_2$
- $q_3 a \rightarrow q_4$
- $q_4 a \rightarrow q_3$


# Minimization: Example 1/4




1)  $X = \{s, f\}$ :      **From one set**      **From one set**




2)  $X = \{q_1, q_2, q_3, q_4\}$ :      **From one set**




# Minimization: Example 1/4



1)  $X = \{s, f\}$ :      **From one set**      **From one set**




2)  $X = \{q_1, q_2, q_3, q_4\}$ :      **From one set**







# Minimization: Example 1/4




1)  $X = \{s, f\}$ :      **From one set**      **From one set**




2)  $X = \{q_1, q_2, q_3, q_4\}$ :      **From one set**




# Minimization: Example 1/4




1)  $X = \{s, f\}$ :      **From one set**      **From one set**




2)  $X = \{q_1, q_2, q_3, q_4\}$ :      **From one set**




# Minimization: Example 1/4



1)  $X = \{s, f\}$ : From one set



2)  $X = \{q_1, q_2, q_3, q_4\}$ : From one set



**Division:**  $\{q_1, q_2, q_3, q_4\} \Rightarrow \underbrace{\{q_1, q_2\}}_{X_1}, \underbrace{\{q_3, q_4\}}_{X_2}$

$\{q_1, q_2\} \cap Q_1 = \emptyset$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

---

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

---


1)  $X = \{s, f\}$ :

$$d = a: \begin{array}{l} sa \rightarrow f \\ fa \rightarrow s \end{array}$$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$


1)  $X = \{s, f\}$ : **From one set**



# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ : **From one set**

$$\begin{array}{ll}
 d = a: & \begin{array}{l} sa \rightarrow f \\ fa \rightarrow s \end{array} \\
 d = b: & \begin{array}{l} sb \rightarrow q_3 \\ fb \rightarrow q_4 \end{array}
 \end{array}$$


# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$   
 $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$   
 $fb \rightarrow q_4$



# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$   
 $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$   
 $fb \rightarrow q_4$

2)  $X = \{q_1, q_2\}$ :

$d = a$ :  $q_1 a \rightarrow q_1$   
 $q_2 a \rightarrow q_2$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ : **From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

**From one set**

2)  $X = \{q_1, q_2\}$ : **From one set**

$d = a$ :  $q_1a \rightarrow q_1$ ,  $q_2a \rightarrow q_2$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

2)  $X = \{q_1, q_2\}$ :

**From one set**

$d = a$ :  $q_1a \rightarrow q_1$ ,  $q_2a \rightarrow q_2$

$d = b$ :  $q_1b \rightarrow s$ ,  $q_2b \rightarrow f$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

2)  $X = \{q_1, q_2\}$ :

**From one set**

$d = a$ :  $q_1a \rightarrow q_1$ ,  $q_2a \rightarrow q_2$

**From one set**

$d = b$ :  $q_1b \rightarrow s$ ,  $q_2b \rightarrow f$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

2)  $X = \{q_1, q_2\}$ :

**From one set**

$d = a$ :  $q_1a \rightarrow q_1$ ,  $q_2a \rightarrow q_2$

**From one set**

$d = b$ :  $q_1b \rightarrow s$ ,  $q_2b \rightarrow f$

3)  $X = \{q_3, q_4\}$ :

$d = a$ :  $q_3a \rightarrow q_3$ ,  $q_4a \rightarrow q_4$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

2)  $X = \{q_1, q_2\}$ :

**From one set**

$d = a$ :  $q_1a \rightarrow q_1$ ,  $q_2a \rightarrow q_2$

**From one set**

$d = b$ :  $q_1b \rightarrow s$ ,  $q_2b \rightarrow f$

3)  $X = \{q_3, q_4\}$ :

**From one set**

$d = a$ :  $q_3a \rightarrow q_3$ ,  $q_4a \rightarrow q_4$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

2)  $X = \{q_1, q_2\}$ :

**From one set**

$d = a$ :  $q_1a \rightarrow q_1$ ,  $q_2a \rightarrow q_2$

**From one set**

$d = b$ :  $q_1b \rightarrow s$ ,  $q_2b \rightarrow f$

3)  $X = \{q_3, q_4\}$ :

**From one set**

$d = a$ :  $q_3a \rightarrow q_3$ ,  $q_4a \rightarrow q_4$

$d = b$ :  $q_3b \rightarrow q_1$ ,  $q_4b \rightarrow q_2$

# Minimization: Example 2/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

1)  $X = \{s, f\}$ :

**From one set**

$d = a$ :  $sa \rightarrow f$ ,  $fa \rightarrow s$

**From one set**

$d = b$ :  $sb \rightarrow q_3$ ,  $fb \rightarrow q_4$

2)  $X = \{q_1, q_2\}$ :

**From one set**

$d = a$ :  $q_1a \rightarrow q_1$ ,  $q_2a \rightarrow q_2$

**From one set**

$d = b$ :  $q_1b \rightarrow s$ ,  $q_2b \rightarrow f$

3)  $X = \{q_3, q_4\}$ :

**From one set**

$d = a$ :  $q_3a \rightarrow q_3$ ,  $q_4a \rightarrow q_4$

**From one set**

$d = b$ :  $q_3b \rightarrow q_1$ ,  $q_4b \rightarrow q_2$

**No next divisions !!!**



# Minimization: Example 3/4

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$$

$$\begin{array}{l} sa \rightarrow f \in R: \\ fa \rightarrow s \in R: \end{array} \} \Rightarrow \{s, f\}a \rightarrow \{s, f\} \in R_m$$

$$\begin{array}{l} sb \rightarrow q_3 \in R: \\ fb \rightarrow q_4 \in R: \end{array} \} \Rightarrow \{s, f\}b \rightarrow \{q_3, q_4\} \in R_m$$

$$\begin{array}{l} q_1a \rightarrow q_1 \in R: \\ q_2a \rightarrow q_2 \in R: \end{array} \} \Rightarrow \{q_1, q_2\}a \rightarrow \{q_1, q_2\} \in R_m$$

$$\begin{array}{l} q_1b \rightarrow s \in R: \\ q_2b \rightarrow f \in R: \end{array} \} \Rightarrow \{q_1, q_2\}b \rightarrow \{s, f\} \in R_m$$

$$\begin{array}{l} q_3a \rightarrow q_3 \in R: \\ q_4a \rightarrow q_4 \in R: \end{array} \} \Rightarrow \{q_3, q_4\}a \rightarrow \{q_3, q_4\} \in R_m$$

$$\begin{array}{l} q_3b \rightarrow q_1 \in R: \\ q_4b \rightarrow q_2 \in R: \end{array} \} \Rightarrow \{q_3, q_4\}b \rightarrow \{q_1, q_2\} \in R_m$$

# Minimization: Example 4/4

$$s \in \{s, f\} \Rightarrow s_m := \{s, f\}$$


---

$$\begin{matrix} s \in F: \\ f \in F: \end{matrix} \Rightarrow \{s, f\} \in F_m$$


---

$$M_m = (Q_m, \Sigma, R_m, s_m, F_m), \text{ where: } \Sigma = \{a, b\}, s_m = \{s, f\}$$

$$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}, F_m = \{\{s, f\}\}$$

$$R_m = \{\{s, f\}a \rightarrow \{s, f\}, \{s, f\}b \rightarrow \{q_3, q_4\}, \{q_1, q_2\}a \rightarrow \{q_1, q_2\}, \\ \{q_1, q_2\}b \rightarrow \{s, f\}, \{q_3, q_4\}a \rightarrow \{q_3, q_4\}, \{q_3, q_4\}b \rightarrow \{q_1, q_2\}\}$$


---

# Minimization: Example 4/4

$$s \in \{s, f\} \Longrightarrow s_m := \{s, f\}$$


$$\left. \begin{array}{l} s \in F: \\ f \in F: \end{array} \right\} \Longrightarrow \{s, f\} \in F_m$$

$M_m = (Q_m, \Sigma, R_m, s_m, F_m)$ , where:  $\Sigma = \{a, b\}$ ,  $s_m = \{s, f\}$

$Q_m = \{\{s, f\}, \{q_1, q_2\}, \{q_3, q_4\}\}$ ,  $F_m = \{\{s, f\}\}$

$R_m = \{\{s, f\}a \rightarrow \{s, f\}, \{s, f\}b \rightarrow \{q_3, q_4\}, \{q_1, q_2\}a \rightarrow \{q_1, q_2\},$   
 $\{q_1, q_2\}b \rightarrow \{s, f\}, \{q_3, q_4\}a \rightarrow \{q_3, q_4\}, \{q_3, q_4\}b \rightarrow \{q_1, q_2\}\}$

Summary:



# Variants of FA: Summary

	FA	$\epsilon$ -free FA	DFA	Complete FA	WSFA	Min-State FA
Number of rules of the form $p \rightarrow q$ , where $p, q \in Q$	<b>0-<math>n</math></b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
Number of rules of the form $pa \rightarrow q$ , for any $p \in Q, a \in \Sigma$	<b>0-<math>n</math></b>	<b>0-<math>n</math></b>	<b>0-1</b>	<b>1</b>	<b>1</b>	<b>1</b>
Number of inaccessible states	<b>0-<math>n</math></b>	<b>0-<math>n</math></b>	<b>0-<math>n</math></b>	<b>0-<math>n</math></b>	<b>0</b>	<b>0</b>
Number of nonterminating states	<b>0-<math>n</math></b>	<b>0-<math>n</math></b>	<b>0-<math>n</math></b>	<b>0-<math>n</math></b>	<b>0-1</b>	<b>0-1</b>
Number of this FAs for any regular language.	$\infty$	$\infty$	$\infty$	$\infty$	$\infty$	<b>1</b>

# Main Decidable Problems

## 1. Membership problem:

- **Instance:** FA  $M$ ,  $w \in \Sigma^*$ ; **Question:**  $w \in L(M)$ ?

## 2. Emptiness problem:

- **Instance:** FA  $M$ ; **Question:**  $L(M) = \emptyset$ ?

## 3. Finiteness problem:

- **Instance:** FA  $M$ ; **Question:** Is  $L(M)$  finite?

## 4. Equivalence problem:

- **Instance:** FA  $M_1, M_2$ ; **Question:**  $L(M_1) = L(M_2)$ ?

## Algorithm: Membership Problem

- **Input:** DFA  $M = (Q, \Sigma, R, s, F)$ ;  $w \in \Sigma^*$
  - **Output:** **YES** if  $w \in L(M)$   
**NO** if  $w \notin L(M)$
- 

- **Method:**
  - **if**  $sw \vdash^* f$ ,  $f \in F$  **then** write (**YES**)  
**else** write (**NO**)
- 

**Summary:**

The membership problem for FAs is decidable

## Algorithm: Emptiness Problem

- **Input:** FA  $M = (Q, \Sigma, R, s, F)$ ;
  - **Output:** **YES** if  $L(M) = \emptyset$   
**NO** if  $L(M) \neq \emptyset$
- 

- **Method:**
  - **if**  $s$  is nonterminating **then** write ('**YES**')  
**else** write ('**NO**')
- 

**Summary:**

The emptiness problem for FAs is decidable

# Algorithm: Finiteness Problem

- **Input:** DFA  $M = (Q, \Sigma, R, s, F)$ ;
  - **Output:** **YES** if  $L(M)$  is finite  
**NO** if  $L(M)$  is infinite
- 
- **Method:**
  - Let  $k = \text{card}(Q)$
  - **if** there exist  $z \in L(M)$ ,  $k \leq |z| < 2k$  **then** write ('**NO**')  
**else** write ('**YES**')
- 

**Note:** This algorithm is based on

$L(M)$  is infinite  $\Leftrightarrow$  there exists  $z: z \in L(M), k \leq |z| < 2k$


---

**Summary:**

The finiteness problem for FAs is decidable




# Decidable Problems: Example



---

Question:  $ab \in L(M)$  ?


# Decidable Problems: Example



**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

# Decidable Problems: Example




**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

**Answer:** **YES** because  $sab \vdash^* f, f \in F$

# Decidable Problems: Example




---

**Question:**  $ab \in L(M)$  ?


$sab \vdash sb \vdash f, f \in F$

**Answer: YES** because  $sab \vdash^* f, f \in F$

---

**Question:**  $L(M) = \emptyset$  ?

# Decidable Problems: Example




---

**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$


**Answer:** **YES** because  $sab \vdash^* f, f \in F$

---

**Question:**  $L(M) = \emptyset$  ?

$Q_0 = \{f\}$

# Decidable Problems: Example



**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

**Answer: YES** because  $sab \vdash^* f, f \in F$


**Question:**  $L(M) = \emptyset$  ?

$Q_0 = \{f\}$

1.  $qa' \rightarrow f; q \in Q; a' \in \Sigma: sb \rightarrow f, fa \rightarrow f$

$Q_1 = \{f\} \cup \{s, f\} = \{f, s\} \dots s$  is terminating

# Decidable Problems: Example



**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

**Answer:** **YES** because  $sab \vdash^* f, f \in F$

**Question:**  $L(M) = \emptyset$  ?


$Q_0 = \{f\}$

1.  $qa' \rightarrow f; q \in Q; a' \in \Sigma: sb \rightarrow f, fa \rightarrow f$

$Q_1 = \{f\} \cup \{s, f\} = \{f, s\} \dots s$  is terminating

**Answer:** **NO** because  $s$  is terminating

# Decidable Problems: Example



**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

**Answer:** **YES** because  $sab \vdash^* f, f \in F$

**Question:**  $L(M) = \emptyset$  ?

$Q_0 = \{f\}$

1.  $qa' \rightarrow f; q \in Q; a' \in \Sigma: sb \rightarrow f, fa \rightarrow f$


$Q_1 = \{f\} \cup \{s, f\} = \{f, s\} \dots s$  is terminating

**Answer:** **NO** because  $s$  is terminating

**Question:** Is  $L(M)$  finite?



# Decidable Problems: Example



**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

**Answer: YES** because  $sab \vdash^* f, f \in F$

**Question:**  $L(M) = \emptyset$  ?

$Q_0 = \{f\}$

1.  $qa' \rightarrow f; q \in Q; a' \in \Sigma: sb \rightarrow f, fa \rightarrow f$


$Q_1 = \{f\} \cup \{s, f\} = \{f, s\} \dots s$  is terminating

**Answer: NO** because  $s$  is terminating

**Question:** Is  $L(M)$  finite?  $k = \text{card}(Q) = 2$

All strings  $z \in \Sigma^*: 2 \leq |z| < 4: aa, bb, ab, \dots$

# Decidable Problems: Example



**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

**Answer: YES** because  $sab \vdash^* f, f \in F$

**Question:**  $L(M) = \emptyset$  ?

$Q_0 = \{f\}$

1.  $qa' \rightarrow f; q \in Q; a' \in \Sigma: sb \rightarrow f, fa \rightarrow f$

$Q_1 = \{f\} \cup \{s, f\} = \{f, s\} \dots s$  is terminating


**Answer: NO** because  $s$  is terminating

**Question:** Is  $L(M)$  finite?

$k = \text{card}(Q) = 2$

All strings  $z \in \Sigma^*: 2 \leq |z| < 4: aa, bb, ab \in L(M), \dots$

# Decidable Problems: Example



**Question:**  $ab \in L(M)$  ?

$sab \vdash sb \vdash f, f \in F$

**Answer:** **YES** because  $sab \vdash^* f, f \in F$

**Question:**  $L(M) = \emptyset$  ?

$Q_0 = \{f\}$

1.  $qa' \rightarrow f; q \in Q; a' \in \Sigma: sb \rightarrow f, fa \rightarrow f$

$Q_1 = \{f\} \cup \{s, f\} = \{f, s\} \dots s$  is terminating

**Answer:** **NO** because  $s$  is terminating

**Question:** Is  $L(M)$  finite?

$k = \text{card}(Q) = 2$

All strings  $z \in \Sigma^*: 2 \leq |z| < 4: aa, bb, ab \in L(M), \dots$

**Answer:** **NO** because there exist  $z \in L(M), k \leq |z| < 2k$

## Algorithm: Equivalence Problem

- **Input:** Two minimum state FA,  $M_1$  and  $M_2$
  - **Output:** **YES** if  $L(M_1) = L(M_2)$   
**NO** if  $L(M_1) \neq L(M_2)$
- 
- **Method:**
  - if  $M_1$  coincides with  $M_2$  except for the name of states  
then write (**YES**)  
else write (**NO**)
- 


### Summary:

The equivalence problem for FA is decidable


# Equivalence Problem: Example

**Question:**  $L(M_1) = L(M_2)$ ?

$M_1$ :




$M_2$ :




# Equivalence Problem: Example

Question:  $L(M_1) = L(M_2)$ ?


$M_1$ :



$M_{min1}$ :




$M_2$ :




# Equivalence Problem: Example

Question:  $L(M_1) = L(M_2)$ ?


$M_1$ :




$M_{min1}$ :



$M_2$ :




$M_{min2}$ :



A minimum state FA

# Equivalence Problem: Example

Question:  $L(M_1) = L(M_2)$ ?



Answer: **YES** because  $M_{min1}$  coincides with  $M_{min2}$