

Contents

1	Constants, Basic definitions	4
2	Existing systems	5
2.1	Octave	5
3	Elementary functions	7
3.1	Polynomials	7
3.1.1	sqrt	7
3.1.2	cbrt	8
3.1.3	pow	8
3.1.4	polynomial roots	8
3.1.5	polynomial_value	9
3.2	Exponentials	9
3.3	Logarithms	10
3.4	Trigonometric functions	11
3.4.1	Circular	11
3.4.2	Hyperbolic	12
3.4.3	Inverse Circular	12
3.4.4	Inverse Hyperbolic	12
3.4.5	Misc	12
3.5	Miscellaneous	13
3.5.1	AGM	13
4	Gamma and related functions	13
4.1	Gamma functions	13
4.2	Factorial, Pochhammer symbol	15
4.3	Incomplete gamma functions	15
4.4	Digamma functions and related	16
4.5	Beta functions	17
4.6	Incomplete beta functions	17
5	Error and related functions	17
5.1	Faddeeva and Dawson integrals	18
5.2	Fresnel integrals	19
5.2.1	Arctangent integral	21
5.3	Iterated error functions	22
6	Bessel, Hankel, Airy functions	22
6.1	Bessel functions	22
6.2	Modified Bessel functions	22
6.3	Incomplete Bessel functions	22
6.4	Spherical Bessel functions	23
6.5	Integral Bessel functions	24
6.6	Hankel, (Whittaker?)	24
6.7	Airy functions	24
6.7.1	Scorer functions	25
6.8	Kelvin functions	27
6.9	Struve functions	27
7	Spherical harmonics, Legendre functions	29

8 Exponential integrals	29
9 Wave equation solutions	30
10 Theta functions	30
10.1 Jacobian theta functions	30
11 Elliptic integrals	31
11.1 Elliptic integrals	32
11.2 Complete elliptic integrals	32
11.3 Complementary complete elliptic integrals	32
11.4 Carlson's symmetric elliptic integrals	33
11.5 Jacobian elliptic functions	33
12 Parabolic cylinder functions	34
12.1 Lebedev	34
12.2 NIST handbook	34
13 Lambert functions	36
14 Hypergeometric, Meijer G functions	37
14.1 Whittaker functions	37
15 Zeta functions	38
15.1 Riemann zeta function	38
15.2 Hurwitz' zeta function	40
15.3 Lerch's transcendent	40
15.4 Dirichlet L-function	40
15.5 Polylogarithms and related	41
15.5.1 Dilogarithm	41
15.5.2 Polylogarithm	43
15.5.3 Fermi-Dirac and Bose-Einstein integrals	43
16 Coulomb wave functions	44
17 Orthogonal polynomials	45
17.1 Chebyshev polynomials	46
17.1.1 First kind	46
17.1.2 Second kind	47
17.2 Legendre polynomials	47
17.3 Laguerre (generalized) polynomials	48
17.4 Gegenbauer (ultraspherical) polynomials	48
17.5 Jacobi polynomials	49
17.6 Hermite polynomials	49
17.6.1 Physicists'	49
17.6.2 Probabilists'	50
18 Misc functions	50
18.1 Anger, Weber functions	50
18.1.1 Lommel functions	50
18.1.2 Anger function	50
18.1.3 Weber function	51
18.2 Howland integrals	51
18.3 Mathieu functions	51
18.3.1 Modified Mathieu functions	52
18.4 Debye functions	52

18.5	Sievert integral	53
18.6	Abramowitz functions	53
18.7	Abramowitz 2 functions	54
18.8	Clausen integral	54
18.9	Voigt (plasma dispersion) function	55
18.10	Spence integral	55
18.11	Angular momentum coupling coefficients	56
18.11.13	$-j$ coefficients	56
18.11.26	$-j$ coefficients	56
18.11.39	$-j$ coefficients	56
19	Combinatorial functions and numbers	57
19.1	Bernoulli numbers, polynomials	57
19.2	Euler numbers, polynomials	57
19.3	Tangent numbers, polynomials	57
19.4	Genocchi numbers, polynomials	58
19.5	Number theoretic functions	58
20	Mathematical finance	58
20.1	Brownian motion	58
20.2	Bachelier model	58
20.3	Black-Scholes-Merton (BS) model	58
20.3.1	Vanilla	58
20.4	Heston stochastic-volatility model	59
20.5	Variance-gamma (VG) model	59
20.6	Constant-elasticity-of-variance (CEV) model	59
21	Probability functions	59
22	Useful techniques	59
22.1	Misc	59
22.1.1	Euler-Maclaurin summation	59
22.1.2	Boole summation	60
22.1.3	Borel summation	60
22.1.4	Kahan's compensated summation trick	60
22.2	Continued fractions	61
22.2.1	Pincherle's theorem	61
22.2.2	Modified Lentz algorithm	61
22.2.3	Quotient-difference algorithm	61
22.3	Sequence acceleration	62
22.3.1	Aitken acceleration	62
22.3.2	Cohen-Villegas-Zagier convergence acceleration of alternating series	62
22.4	Root finding	62
22.4.1	Newton's method	62
22.4.2	Halley's method	62
23	Notes on implementations	63
24	Notes on testing	63
25	General mathematical library	64

1 Constants, Basic definitions

INDEX GLOSSARY SYMBOLS (MARGIN NOTES) EQUATION NUMBERING?

[Generally need *zeros* as well as *derivatives* of all functions... and possibly *integrals*!]

- Mathematics
 - definitions
 - relations / expansions / extensions / etc.
 - discussion of context, usage, other notations, etc.
 - graphics
- Computer science
 - routines (API)
 - implementation notes
 - pseudo-code / code
 - example usage
 - error plots, etc.

Coding style: CamelCase as basic function names, but underscore to separate modifications? For example, Exp, Exp_m1, StruveH, StruveH_ddx; double underscore for internal variations (e.g. different methods, etc.), e.g. Exp_m1_naive, Exp_m1_series.

- $\mathbb{N} = \{1, 2, \dots\}$
- $\mathbb{N}_0 = \{0, 1, 2, \dots\} = \mathbb{N} \cup \{0\}$
- $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$
- Pi

π

$$\pi = 3.1415926535897932384626433\dots$$

$$\frac{1}{\pi} = 12 \sum_{k=0}^{\infty} (-)^k \frac{(6k)!(13591409 + 545140134k)}{(3k)!(k!)^3 640320^{3k+3/2}} \quad \text{Chudnovsky algorithm}$$

- Euler's gamma

γ

$$\gamma = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k} - \log n \right) = - \int_0^{\infty} e^{-t} \ln t \, dt = 0.57721566490153286060\dots$$

- Khintchine constant

K_0

$$K_0 = \prod_{k=1}^{\infty} \left[1 + \frac{1}{k(k+2)} \right]^{\log_2 k} = 2.6854520010\dots$$

- misc

$$\log(2) = \dots$$

$$\sqrt{2} = \dots$$

$$\zeta(2) = \frac{\pi^2}{6} = \dots$$

$$\Gamma(1/3) = 2.6789385347077\dots$$

Basic operations expected of any numeric type: $+, -, *, /, + =, - =, * =, / =, ++, --, ==, !=, <, >, <=, >=$, pow, pow_{int}, sqrt, cbrt, ln, exp, abs, arg, signum, floor, ceil, truncate, mod

Use double-doubles just for more accurate summations (and range-reductions?)

From [Precise]: given a number x and an uncertainty in x of δ_x , we define the *accuracy* as $\text{Accuracy}(x, \delta_x) = (-\log_{10} |\delta_x|)$ (corresponding to the absolute error), the *precision* as $\text{Precision}(x, \delta_x) = -\log_{10} |\frac{\delta_x}{x}|$ (corresponding to the relative error), and the *scale* as $\text{Scale} = \text{Precision} - \text{Accuracy}$. We can visualize different cases as:

If *accuracy*, *precision*, and *scale* are all positive, we have:

$$\overbrace{x_1 x_2 \cdots x_s \cdot x_{s+1} x_{s+2} \cdots x_{s+m}}^{\text{Precision}}$$

$\underbrace{\hspace{1.5cm}}_{\text{Scale}} \quad \underbrace{\hspace{1.5cm}}_{\text{Accuracy}}$

If *accuracy* and *precision* are positive and *scale* negative, we have:

$$\overbrace{.00 \cdots 0 x_1 x_2 \cdots x_p}^{\text{Accuracy}}$$

$\underbrace{\hspace{1.5cm}}_{\text{Scale}} \quad \underbrace{\hspace{1.5cm}}_{\text{Precision}}$

If *precision* and *scale* are positive and *accuracy* negative, we have:

$$\overbrace{x_1 x_2 \cdots x_p 00 \cdots 0.}^{\text{Scale}}$$

$\underbrace{\hspace{1.5cm}}_{\text{Precision}} \quad \underbrace{\hspace{1.5cm}}_{\text{Accuracy}}$

error diagrams; diagrams for techniques (colored nicely) used in 2-parameters, e.g. test of continued fraction typesetting:

$$\frac{1}{x + \frac{2}{x^2 + \frac{3}{x^3 + \cdots}}}$$

$$(1 - \cos x) / \sin x = \sin x / (1 + \cos x)$$

$$e^{a+b} = e^a e^b = e^a (1 + (e^b - 1)) = e^a + e^a (e^b - 1)$$

2 Existing systems

[Check Matlab special function support!]

2.1 Octave

IDEA: build out the special function support for Octave!? Issue is lack of long double type to help with getting full precision. (Can create objects + overloading to maybe create double-double types?)

Note that Octave includes a gsl package for GSL bindings (but GSL generally has poor support for complex numbers...) And, the statistics package includes various probability distributions.

- Built-in version 3.6.2:

- `[a,ierr] = airy(k,z,opt)` — $\text{Ai}(z), \text{Ai}'(z), \text{Bi}(z), \text{Bi}'(z)$ with optional scaling; z can be real or complex
- `[r,ierr] = besselj(alpha,x,opt)/bessely/besseli/besselk/besselh` — Bessel functions $J_\alpha(x), Y_\alpha(x), I_\alpha(x), K_\alpha(x), H_\alpha(x)$ with optional scaling; α must be real, x can be complex

- **beta(a,b)**, **betaln(a,b)** — Beta function $B(a,b)$ or natural logarithm of Beta function $\ln(B(a,b))$; with a,b real
- **betainc(x,a,b)** — incomplete Beta function with x,a,b real
- **bincoeff(n,k)** — binomial coefficient; n,k integers; [accepts some non-integers but returns suspicious values]
- **erf(z)**, **erfc(z)**, **erfcx(z)** — error function $\operatorname{erf}(z)$, complementary error function $\operatorname{erfc}(z) = 1 - \operatorname{erf}(z)$, scaled complementary error function $e^{z^2} \operatorname{erfc}(z)$; z real
- **erfinv(x)** — inverse error function; x real
- **gamma(z)**, **lgamma(z)/gammaln(z)** — gamma function $\Gamma(z)$ or natural logarithm of gamma function $\ln \Gamma(z)$; z real; [looks like iffy/low precision implementation: **gamma(6)**=119.99999999999997]
- **gammainc(x,a,opt)** — normalized incomplete gamma function $P(a,x)$ or complementary version; x,a real
- **legendre(n,x,opt)** — Legendre functions of degree n and order $m = 0 \dots n$, $P_n^m(x)$ optionally normalized
- elementary functions; x,y real, z real or complex
 - * **exp(z)**, **expm1(z)**, **log(z)**, **reallog(z)**, **log1p(z)**, **log10(z)**, **log2(z)**
 - * **pow2(z)**, **pow2(f,e)**, **nextpow2(x)**, **realpow(x,y)**, **sqrt(z)**, **cbrt(x)**, **nthroot(x,n)**
 - * **cos(z)**, **sin(z)**, **tan(z)**, **cosh(z)**, **sinh(z)**, **tanh(z)**, **sec(z)**, **csc(z)**, **cot(z)**, **sech(z)**, **csch(z)**, **coth(z)**
 - * **acos(z)**, **asin(z)**, **atan(z)**, **acosh(z)**, **asinh(z)**, **atanh(z)**, **asec(z)**, **acsc(z)**, **acot(z)**, **asech(z)**, **acsch(z)**, **acoth(z)**
 - * **atan2(y,x)**, **abs(z)**, **arg(z)**, **conj(z)**, **real(z)**, **imag(z)**
 - * **sind(z)**, **cosd(z)**, **tand(z)**, **secd(z)**, **cscd(z)**, **cotd(z)**, **asind(z)**, **acosd(z)**, **atand(z)**, **asecd(z)**, **acscd(z)**, **acotd(z)**
 - * **ceil(z)**, **fix(z)**, **floor(z)**, **round(z)**, **roundb(z)**, **max(z1...zn)**, **min(z1...zn)**, **hypot(z1...zn)**

• Package **specfun-1.1.0**

- **cosint(z) = Ci(z)** — cosine integral function $\operatorname{Ci}(z) = \int_x^\infty \frac{\cos(t)}{t} dt$; z real or complex; [uses expression in terms of exponential integrals E_1 or Ei] [iffy for some cases: **Ci(0) = NaN - NaN*i**]
- **sinint(z) = Si(z)** — sine integral function $\operatorname{Si}(z) = \int_0^x \frac{\sin(t)}{t} dt$; z real or complex; [uses summation of Bessel J functions] [iffy for some cases: **Si(Inf) = Si(1e10) = NaN + NaN*i**]
- **heaviside(x,threshold)** — Heaviside function; z real or complex (but poor behaviour for complex (all non-real give 1) — this raises the question of what *should* be the extension of the heaviside to \mathbb{C} ?)
- **dirac(z)** — Dirac delta function: **Inf** iff $x = 0$ else 0; z real or complex
- **zeta(z)** — Riemann zeta function; z real or complex; [uses quadrature $\int_0^\infty \frac{x^{z-1}}{\Gamma(z)(e^x-1)} dx$ for real $z > 1$; summation $\frac{1}{1-2^{1-z}} \sum_{k=1}^\infty \frac{(-)^{k-1}}{k^z}$ for values with $\Re z \geq 0$; and otherwise recursively computes $2^z \pi^{z-1} \sin(\pi z/2) \Gamma(1-z) \zeta(1-z)$]
- **lambertw(b,z)** — computes Lambert W function $W(z)$ for branch b ; z real or complex, b integer; [uses series around $-1/e$ and asymptotic expansion at 0 and ∞ to get initial guess, then Halley iteration to refine]
- **psi(x)** — computes $\frac{d}{dx} \ln \Gamma(x)$; real x ; [does simple finite centered difference approximation]
- **expint(z)=expint.E1(z)**, — computes exponential integral $\int_x^\infty e^t/t dt$; z real or complex; [uses **expint.Ei(-z)+C**]
- **expint.Ei(z)** — computes exponential integral $-\int_{-x}^\infty e^t/t dt$; z real or complex; [uses quadrature for real $z > 2$ or $z < 0$; if $|z| \geq 10$, $\Im z \leq 0$ uses asymptotic expansion (with conjugation for $\Im z > 0$); otherwise uses series expansion]

- `erfcinv(z)` — inverse complementary error function; z real or complex; [simply returns `erfinv(1-x)` — which will have precision issues for $x \sim 1$]
- `[y,p] = laguerre(z,n)` — computes the Laguerre polynomial of order n at z ; z real or complex, $n > 0$ integer; [computes polynomial coefficients via recursion and then evaluates resulting polynomial at z ; sounds iffy to me...]
- `[k,e] = ellipke(m)` — returns complete elliptic integrals of first $K(m)$ and second $E(m)$ kind; real $m \leq 1$; [uses AGM algorithm and a transform for $m < 0$]

- Package `miscellaneous-1.1.0`

- `chebyshevpoly(kind,order,z)` — computes first $T_n(x)$ or second $U_n(x)$ Chebyshev polynomials (coefficients or value at z); z real or complex; [uses recursion for coefficients; value is evaluated from polynomial; seems iffy to me]
- `hermitepoly(order,z)` — computes Hermite polynomials $H_n(z)$ (coefficients or value at z); z real or complex; [uses low-precision (8-digits) table for $0 \leq \text{order} \leq 50$; otherwise uses recursion to compute polynomial coefficients; value is evaluated from polynomial coefficients; seems iffy]
- `laguerrepoly(order,z)` — computes Laguerre polynomials $L_n(z)$ (coefficients or value at z); z real or complex; [uses recursion to compute polynomial coefficients; value is evaluated from polynomial coefficients; seems iffy]
- `legendrepoly(order,z)` — computes Legendre polynomials $L_n(z)$ (coefficients or value at z); z real or complex; [uses recursion to compute polynomial coefficients; value is evaluated from polynomial coefficients; seems iffy]

3 Elementary functions

`ldexp`

`frexp`

`mod`

`abs`

`hypot`

`hypot3`

Also an `fmod(z,a,b)` which reduces z with high-accuracy modulo $a + b$ (a “pseudo-quad-double” functionality) which is useful for trigonometric range-reduction, for example.

A handy function is a multi-precision constant factory: takes decimal string, say, and returns n doubles whose sum is equal to the constant. (Case $n = 2$ is just the constructor for `qdouble...`)

3.1 Polynomials

3.1.1 `sqrt`

Assume that $a < 0$ or $b \neq 0$, then we have:

$$\sqrt{a + bi} = \frac{b}{2d} + di \quad d = \sqrt{\frac{-a \pm \sqrt{a^2 + b^2}}{2}}$$

We could also use a trigonometric approach: $\sqrt{a + bi} = \sqrt{\sqrt{a^2 + b^2}} \left(\cos(\text{atan}(\frac{b}{a})/2) + i \sin(\text{atan}(\frac{b}{a})/2) \right)$.

Implementation notes for \sqrt{z} for $z \geq 0$:

- Use the Newton iteration $x_{n+1} = \frac{1}{2}(x_n + \frac{z}{x_n})$
- This converges (for doubles) in only a couple more iterations than Halley's method (and each iteration is cheaper)
- Gives exact answers (for gcc) in 4 or 5 iterations
- For starting value, let $z = 2^N \cdot f$ with N even and $f \in [1/2, 2)$ (use **frexp**), and use the starting guess of $2^{N/2} \hat{f}$ where \hat{f} is the linearly interpolated value of $\sqrt{\cdot}$ on $[1/2, 2)$ for the value f .
- Brent claims that it is better (optimal?) to use $\sqrt{a} = a/\sqrt{a}$ and compute $a^{-1/2}$ directly via the 3rd-order iteration $x_{n+1} = x_n - \frac{x_n}{2}(\varepsilon_n - \frac{3}{4}\varepsilon_n^2)$ where $\varepsilon_n = ax_n^2 - 1$. But for the case of a double, this converges no faster than Newton.

3.1.2 cbrt

For complex values, use trigonometric approach. (Use a newton-step to clean digits?) Also, note that if $x = z^{1/3}$ is a cube-root, then so are ζx , $\zeta^2 x$, where $\zeta = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ is a cube root of unity.

Implementation notes for $\sqrt[3]{z}$ for $z \in \mathbb{R}$:

- Use the Newton iteration $x_{n+1} = \frac{x_n^4 + 2zx_n}{2x_n^3 + z}$
- This converges (for doubles) in only a couple more iterations than Halley's method (and each iteration is cheaper)
- For starting value, let $z = 2^N \cdot f$ with $N = 3k$ and $f \in [1/2, 4)$ (use **frexp**), and use the starting guess of $2^{N/3} \hat{f}$ where \hat{f} is the linearly interpolated value of $\sqrt[3]{\cdot}$ on $[1/2, 4)$ for the value f .
- Gives exact answers (for gcc) in 5 or 6 iterations (Halley takes about 4 iterations)

3.1.3 pow

$$\text{pow}(z, r) = z^r \quad z \in \mathbb{R}, r \in \mathbb{R}$$

$$\text{pow.int}(z, n) = z^n \quad z \in \mathbb{R}, n \in \mathbb{Z}$$

integer, real, complex versions (branches?), fully general (integer) $x^2, x^3, x^y - 1, \sqrt{1+x} - 1, \sqrt{1+x} - x$

Implementation notes for $z^n, n \in \mathbb{Z}$:

- Assume $n > 0$ (compute z^{-n} and use $z^n = 1/z^{-n}$)
- Use repeated doubling algorithm - takes $\log n$ steps
- Gives exact answers

3.1.4 polynomial roots

$$\text{solve.linear}(a, b) = \rho \quad \text{s.t.} \quad a\rho + b = 0 \quad a, b \in \mathbb{R}$$

$$\text{solve.quadratic}(a, b, c) = \rho \quad \text{s.t.} \quad a\rho^2 + b\rho + c = 0 \quad a, b, c \in \mathbb{R}$$

$$\text{solve_cubic}(a, b, c, d) = \rho \quad \text{s.t.} \quad a\rho^3 + b\rho^2 + c\rho + d = 0 \quad a, b, c, d \in \mathbb{R}$$

$$\text{solve_quartic}(a, b, c, d, e) = \rho \quad \text{s.t.} \quad a\rho^4 + b\rho^3 + c\rho^2 + d\rho + e = 0 \quad a, b, c, d, e \in \mathbb{R}$$

$$\text{polynomial_root}(\vec{a}, n) = \rho \quad \text{s.t.} \quad \sum_{k=0}^n a^{n-k} \rho^k = 0 \quad \vec{a} \in \mathbb{R}^{n+1}$$

Complex versions also...

For n -th root of r , newton looks like $x_{k+1} = \frac{n-1}{n}x_k + \frac{r}{nx_k^{n-1}}$.

There are direct methods here, trigonometric (using complex numbers), etc.

For general polynomial root-finding, NR recommends “Laguerre” method + deflation. (Which works decently as long as there are not repeated roots...)

Remark: to transform a general monic quadratic $x^4 + bx^3 + cx^2 + dx + e$ to a “depressed” quadratic we take $x = y - \frac{b}{4}$ and get

$$y^4 + (c - \frac{3b^2}{8})y^2 + (\frac{b^3}{8} - \frac{bc}{2} + d)y + (e - \frac{3b^4}{256} + \frac{b^2c}{16} - \frac{bd}{4})$$

3.1.5 polynomial_value

$$\text{polynomial_value}(\vec{a}, n, z) = \sum_{k=0}^n a^{n-k} z^k \quad \vec{a} \in \mathbb{R}^{n+1}, z \in \mathbb{R}$$

3.2 Exponentials

Exponential function:

$$\exp(z) = e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!} \quad z \in \mathbb{C}$$

$\exp(z), e^z$

$$\text{exp}(z) = e^z$$

$$\text{exp_m1}(z) = e^z - 1$$

$$\text{exp_m1_dx}(z) = \frac{e^z - 1}{z}$$

$$2^x, 10^x, e^z - 1 - z, \frac{e^z - 1 - z}{z^2}, e_N(x) = \sum_{n=0}^N \frac{x^n}{n!}, e^z - e_N(z), \frac{e^z - e_N(z)}{z^{N+1}}$$

Remark: $e^{x+yi} = e^x \cos y + ie^x \sin y$ for $x, y \in \mathbb{R}$.

Implementation notes for $\exp(z)$ for $z \in \mathbb{R}$:

- For $z < 0$, use $\exp(z) = \frac{1}{\exp(-z)}$ and compute $\exp(-z)$ (to avoid negative cancellation in series)
- For $|z| > \frac{1}{2}$, use $\exp(2^N \cdot x) = \exp(x)^{2^N}$ and compute $\exp(z \cdot 2^N)$ with $z \cdot 2^N \leq \frac{1}{2}$
- Finally, use standard power-series $\exp(z) = {}_0F_0(|z|) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$ for the argument $0 < z \leq \frac{1}{2}$ to compute.
- In tests, this gives results which are within 1 ulp of the (gcc math library computed) actual values, many values are exactly the same. (Note that this is actually depending on intermediate values being computed with higher precision! using the `-fexcess-precision=standard` option in gcc gives much higher error rates! Explicitly using `long double` variables (20-byte) brings us back to the low error-rate.)

- Luke claims that convergence is better for $\Re(z) > -1$ computing:

$$e^{-z} = (1+z)^{-1} {}_1F_1\left(\begin{matrix} z \\ 2+z \end{matrix} \middle| -z\right) = \frac{1}{1+z} \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} \frac{(z)_k}{(2+z)_k}$$

(If α_n is the n th term in the sum, then $\alpha_{n+1} = -\alpha_n \frac{z(z+n)}{(n+1)(z+n+2)}$.)

- Another idea: write $z = (\ln 2)^N \xi$ so that $e^z = 2^N e^\xi$

Implementation notes for $\exp(z) - 1$ for $z \in \mathbb{R}$:

- If $|z| \geq 1/2$, just compute $\exp(z) - 1$ directly
- Otherwise use a power series $\sum_{n=1}^{\infty} \frac{z^n}{n!}$
- This takes up to 18 terms in the series to converge for double precision
- There are sporadic errors of 1 or 2 ulps (vs gcc) in both branches of the above
- for the series, this can be ameliorated a lot by using long double
- can we use the trick $e^x - 1 = (e^{x/2} + 1)(e^{x/4} + 1)(e^{x/8} + 1)(e^{x/8} - 1)$ (etc.) somehow?

Note that for $z \in \mathbb{C}$ there can be cancellation of the real-part away from $z = 1$ also (in particular when $(\Re z) \cos(\Im z) \sim 1 \dots$)

3.3 Logarithms

$$\log(z) = \rho \quad \text{s.t.} \quad e^\rho = z$$

$$\log_b(z) = \log(z) / \log(b) = \rho \quad \text{s.t.} \quad b^\rho = z$$

$$\mathbf{log}(z) = \log(z)$$

$$\mathbf{Log}(z, w) = \mathbf{Log}_{[w]}(z) \quad \text{with branch}$$

$$\mathbf{log_b}(z, b) = \log_b(z)$$

$$\mathbf{log_p1}(z) = \log(1+z)$$

$$\mathbf{log_p1_mx}(z) = \log(1+z) - z$$

$$\mathbf{log_2}(z) = \log_2(z)$$

$$\mathbf{log_10}(z) = \log_{10}(z)$$

$z \ln z, \frac{\ln(1+z)}{z}, \frac{\ln(1+z)-z}{-z^2/2}$, etc. partial series, etc.

Remark: $\ln x + y\hat{i} = \ln r + \vartheta\hat{i}$, where $x, y \in \mathbb{R}$, $x + y\hat{i} = re^{\vartheta\hat{i}}$, thus $r = |x + y\hat{i}| = \sqrt{x^2 + y^2}$ and $\vartheta = \arg x + y\hat{i} = \text{atan2}(y, x)$.

Implementation notes for $\log(z)$ for $z > 0$:

- For $z > 1$, use $\log(z) = -\log \frac{1}{z}$ and compute $\log \frac{1}{z}$ (to get $z \leq 1$)
- Next, use $\log(2^N x) = N \log 2 + \log x$, so can reduce to get $\frac{1}{2} \leq z \leq \frac{3}{2}$
- Finally, use series $\log(1+z) = -\sum_{n=1}^{\infty} (-)^n \frac{z^n}{n}$

In tests, this gives results which are within 3 ulp of the (gcc math library computed) actual values.

3.4 Trigonometric functions

polar to/from rectangular, angle, radians to/from degrees to/from gradians, etc.

`sin`, `cos`, `tan`, `sec`, `csc`, `cot`, `sinh`, `cosh`, `tanh`, `seca`, `csca`, `cota`, `asin`, `acos`, `atan`, `asec`, `acsc`, `acot`, `asinh`, `acosh`, `atanh`, `aseca`, `acsca`, `acota`, `sind`, `cosd`, ..., `atan2`, `gudermannian`, `versine`, `haversine`, `coversine`, `hacoversine`, `exsecant`, `excosecant`, `sinc`, `sinc_a`

$$\text{sinc}(x) = \frac{\sin \pi x}{\pi x}$$

$$\sin_{\pi}(x) = \sin(\pi x)$$

(and etc. Allows for reduced rounding error in many usages of `sin`.)

3.4.1 Circular

$$\cos(z) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{(2n)!}$$

$$\sin(z) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!}$$

$$\tan(z) = \frac{\sin z}{\cos z} =$$

$$\sec(z) = \frac{1}{\cos z}$$

$$\cot(z) = \frac{1}{\tan z} = \frac{\cos z}{\sin z}$$

$$\csc(z) = \frac{1}{\sin z}$$

Remark: $\cos x + y\hat{i} = \cos x \cosh y - \hat{i} \sin x \sinh y$, and $\sin x + y\hat{i} = \sin x \cosh y + \hat{i} \cos x \sinh y$, when $x, y \in \mathbb{R}$.

- continued fraction $\tan z = \frac{z}{1 - \frac{z}{3 - \frac{z}{5 - \dots}}}$
- $\cot z = \frac{1}{2} + 2z \sum_{k=1}^{\infty} \frac{1}{z^2 - k^2 \pi^2}$
- $\text{asin } z = z + \frac{1}{2} \frac{z^3}{3} + \frac{1 \cdot 3}{2 \cdot 4} \frac{z^5}{5} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \frac{z^7}{7} + \dots$
- continued fraction $\frac{\text{asin } z}{\sqrt{1-z^2}} = \frac{z}{1 - \frac{1 \cdot 2 z^2}{3 - \frac{1 \cdot 2 z^2}{5 - \frac{3 \cdot 4 z^2}{7 - \frac{3 \cdot 4 z^2}{9 - \dots}}}}}$
- Expansions for $\text{atan } z$
 - continued fraction $\text{atan } z = \frac{z}{1 + \frac{z^2}{3 + \frac{4z^2}{5 + \frac{9z^2}{7 + \dots}}}}$
 - for $|z| \leq 1$, $\text{atan } z = z - \frac{z^3}{3} + \frac{z^5}{5} - \frac{z^7}{7} \dots$
 - for $|z| > 1$, $\text{atan } z = \frac{\pi}{2} - \frac{1}{z} + \frac{1}{3z^2} - \frac{1}{5z^4} \dots$
 - for any $z \neq \pm \hat{i}$, $\text{atan } z = \frac{z}{1+z^2} \left(1 + \frac{2}{3} \frac{z^2}{1+z^2} + \frac{2 \cdot 4}{3 \cdot 5} \left(\frac{z^2}{1+z^2} \right)^2 + \dots \right)$

3.4.2 Hyperbolic

$$\begin{aligned}\cosh(z) &= \cos(\hat{i}z) = \sum_{n=0}^{\infty} \frac{z^{2n}}{(2n)!} = \frac{e^z + e^{-z}}{2} \\ \sinh(z) &= -\hat{i} \sin(\hat{i}z) = \sum_{n=0}^{\infty} \frac{z^{2n+1}}{(2n+1)!} = \frac{e^z - e^{-z}}{2} \\ \tanh(z) &= -\hat{i} \tan(\hat{i}z) = \frac{\sinh z}{\cosh z} = \frac{e^z - e^{-z}}{e^z + e^{-z}} \\ \operatorname{sech}(z) &= \frac{1}{\cosh z} \\ \operatorname{coth}(z) &= \frac{1}{\tanh z} = \frac{\cosh z}{\sinh z} \\ \operatorname{csch}(z) &= \frac{1}{\sinh z}\end{aligned}$$

3.4.3 Inverse Circular

$$\begin{aligned}\operatorname{acos}(z) &= \rho \quad \text{s.t.} \quad \cos \rho = z \\ \operatorname{asin}(z) &= \rho \quad \text{s.t.} \quad \sin \rho = z\end{aligned}$$

$$\operatorname{atan}(z) = \rho \quad \text{s.t.} \quad \tan \rho = z \quad \rho \in (-\pi/2, +\pi/2)$$

$$\operatorname{atan2}(y, x) = \operatorname{atan}(y/x) \quad \text{s.t.} \quad \text{correct for } x \sim 0 \text{ and signs}$$

Remark: $\operatorname{acos} x + y\hat{i} = \operatorname{acos} \beta + \hat{i} \operatorname{sign}_+(y) \ln(\alpha + \sqrt{\alpha^2 - 1})$ and $\operatorname{asin} x + y\hat{i} = \operatorname{asin} \beta + \hat{i} \operatorname{sign}_+(y) \ln(\alpha + \sqrt{\alpha^2 - 1})$; where $x, y \in \mathbb{R}$, $\alpha = \frac{\sqrt{(x+1)^2 + y^2} + \sqrt{(x-1)^2 + y^2}}{2}$, $\beta = \frac{\sqrt{(x+1)^2 + y^2} - \sqrt{(x-1)^2 + y^2}}{2}$, and

$$\operatorname{sign}_+(y) = \begin{cases} +1 & y \geq 0 \\ -1 & y < 0 \end{cases}$$

3.4.4 Inverse Hyperbolic

$$\begin{aligned}\operatorname{acosh}(z) &= \rho \quad \text{s.t.} \quad \cosh \rho = z \\ \operatorname{asinh}(z) &= \rho \quad \text{s.t.} \quad \sinh \rho = z \\ \operatorname{atanh}(z) &= \rho \quad \text{s.t.} \quad \tanh \rho = z\end{aligned}$$

3.4.5 Misc

$$\operatorname{gud}(z) \quad \operatorname{gud}(z) = 2 \operatorname{atan} e^z - \frac{\pi}{2} \quad \text{Gudermannian}$$

Note: for a function $F(z)$, we define $\operatorname{ver}F(z) = 2F(z/2)^2$, $\operatorname{co}F(z) = F(\frac{\pi}{2} - z)$, $\operatorname{ha}F(z) = F(z)/2$, $\operatorname{ex}F(z) = F(z) - 1$, giving us

$$\operatorname{vsin}(z) = \operatorname{versine}(z) = 2 \sin(z/2)^2 = 1 - \cos(z) = \operatorname{exc cosine}(z)$$

$$\operatorname{hvsin}(z) = \operatorname{haversine}(z) = \frac{1}{2} \operatorname{versine}(z) = \frac{1 - \cos(z)}{2} = \sin(z/2)^2$$

$$\operatorname{cvsin}(z) = \operatorname{coversine}(z) = 1 - \sin(z) = \operatorname{versine}(\frac{\pi}{2} - z) = 2 \sin(\frac{\pi}{4} - \frac{z}{2})$$

$$\operatorname{chvsin}(z) = \operatorname{hvc sin}(z) = \operatorname{hacoversine}(z) = \operatorname{cohaversine}(z) = \frac{1}{2} \operatorname{coversine}(z) = \frac{1 - \sin(z)}{2}$$

$$\operatorname{vcos}(z) = \operatorname{vercosine}(z) = 2 \cos(z/2)^2 = 1 + \cos(z)$$

$$\operatorname{hvcos}(z) = \operatorname{havercosine}(z) = \frac{1}{2} \operatorname{vercosine}(z) = \frac{1 + \cos(z)}{2}$$

$$\text{cvcos}(z) = \text{covercosine}(z) = 1 + \sin(z) = \text{vercosine}(\frac{\pi}{2} - z) = 2 \sin(\frac{\pi}{4} + \frac{z}{2})$$

$$\text{hcvcos}(z) = \text{chvcos}(z) = \text{hacovercosine}(z) = \text{cohavercosine}(z) = \frac{1}{2} \text{covercosine}(z) = \frac{1 + \sin(z)}{2}$$

$$\text{exsec}(z) = \text{exsecant}(z) = \sec(z) - 1 = \frac{1 - \cos(z)}{\cos(z)} = \frac{\text{versine}(z)}{\cos(z)} = 2 \sin(z/2)^2 \sec(z)$$

$$\text{excsc}(z) = \text{excosecant}(z) = \text{exsecant}(\frac{\pi}{2} - z) = \csc(z) - 1$$

[The equalities for excsc need to be verified... they appear to be incorrect]

3.5 Miscellaneous

3.5.1 AGM

`agm(a,b)`, `agm(a,b,out c)`

Given $a_0 = \alpha$, $b_0 = \beta$, let

$$\begin{aligned} a_{n+1} &= (a_n + b_n)/2 \\ b_{n+1} &= (a_n \cdot b_n)^{1/2} \\ c_{n+1} &= (a_n - b_n)/2 \end{aligned}$$

then $\text{agm}(\alpha, \beta) = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n$. For complex values, we choose the square-root such that $\text{Ph } b_{n+1} = (\text{Ph } a_n + \text{Ph } b_n)/2$ with the phases of a_n , b_n chosen such that the difference is less than π .

4 Gamma and related functions

What about computation of *ratios* of Gamma functions? (To handle cancellation of poles, etc. nicely; avoid scaling issues, etc.) `gamma_ratio([a1,...,an],[b1,...,bm])`

Note that

$$\lim_{\varepsilon \rightarrow 0} \frac{\Gamma(-n + \varepsilon)}{\Gamma(-m + \varepsilon)} = (-)^{m+n} \frac{m!}{n!} \quad m, n \in \mathbb{N}$$

4.1 Gamma functions

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt \quad z \in \mathbb{C} \setminus \mathbb{N}_0 \quad \Gamma(z)$$

Note that that integral definition only applies for $\Re z > 0$, but the function can be extended to a meromorphic function with simple poles at $z = 0, -1, -2, \dots$

$$\Gamma^*(z) = \frac{x^{-x+1/2} e^x}{\sqrt{2\pi}} \Gamma(z) \quad \Gamma^*(z)$$

$$\frac{1}{\Gamma(z+1)} = e^{\gamma z} \prod_{n=1}^{\infty} e^{-z/n} (1 + z/n)$$

Binet function:

$$J(z) = \ln \Gamma(z) + z - (z - \frac{1}{2}) \ln z - \frac{1}{2} \ln 2\pi \quad J(z)$$

$$\text{gamma}(z) = \Gamma(z)$$

$$\text{gamma_p1_m1}(z) = \Gamma(1+z) - 1$$

$$\text{log_gamma}(z) = \log(\Gamma(z))$$

$$\text{log_abs_gamma}(z) = \log(\Gamma(|z|))$$

$$\text{gammastar}(z) = \Gamma^*(z)$$

$$\text{gamma_inv}(z) = 1/\Gamma(z)$$

log-gamma+sign also

Amusing fact: $\zeta(3) = \lim_{\varepsilon \rightarrow 0} \frac{1}{2\varepsilon^3} \frac{\Gamma^3(1+\varepsilon)\Gamma(1-\varepsilon)}{\Gamma(1+\varepsilon)}$ with error $\sim 2\varepsilon$.

Spouge's approximation (a modification of Stirling's):

$$\Gamma(z+1) = (z+a)^{z+1/2} e^{-(z+a)} \left(c_0 + \sum_{k=1}^{a-1} \frac{c_k}{z+k} + \varepsilon_a(z) \right)$$

where a is arbitrary positive integer and $c_0 = \sqrt{2\pi}$ with (for $k = 1, 2, \dots, a-1$):

$$c_k = \frac{(-)^{k-1}}{(k-1)!} (-k+a)^{k-1/2} e^{-k+a}$$

If $\Re z > 0$ and $a > 2$, then the relative error in discarding $\varepsilon_a(z)$ is bounded by $a^{-1/2}(2\pi)^{-(a+1/2)}$. (Note that this has controllable error but the large coefficients have cancellative properties...) This is a convenient “work-horse” algorithm — it's conveniently modified to directly compute Γ^* nicely.

The Lanczos approximation (from Wikipedia):

$$\Gamma(z+1) = \sqrt{2\pi} (z+g+\frac{1}{2})^{z+1/2} e^{-(z+g+1/2)} A_g(z)$$

with g an arbitrary constant (with $\Re(z+g+1/2) > 0$) and

$$A_g(z) = \frac{1}{2}p_0(g) + \frac{z}{z+1}p_1(g) + \frac{z(z-1)}{(z+1)(z+2)}p_2(g) + \dots$$

... valid only for $\Re z > 0$... (test of implementation from wikipedia page gives bad results, not clear why)

A few useful relations

(A) $\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin \pi z}$; equivalently, $\ln |\Gamma(z)| + \ln |\Gamma(1-z)| = \ln \pi - \ln |\sin \pi z|$

(B) $\Gamma(z+1) = z\Gamma(z)$; equivalently $\ln |\Gamma(z+1)| = \ln |z| + \ln |\Gamma(z)|$

(C) Stirling's approximation for large n :

$$n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \left(1 + \frac{1}{12n} + \frac{1}{288n^2} - \frac{139}{51840n^3} - \frac{571}{2488320n^4} \dots\right)$$

(D) For $|z| < 1$ (< 2 ?),

$$\ln \Gamma(1+z) = -\ln(1+z) + z(1-\gamma) + \sum_{k=2}^{\infty} (-)^k (\zeta(k) - 1) \frac{z^k}{k}$$

Use this for tiny z , especially to directly compute $\Gamma(1+z) - 1$ — it works beautifully.

(E) A power-series at zero, $\frac{1}{\Gamma(z)} = \sum_{k=1}^{\infty} c_k z^k$, where $c_1 = 1$, $c_2 = \gamma$, and $(k-1)c_k = \gamma c_{k-1} - \zeta(2)c_{k-2} + \zeta(3)c_{k-3} - \dots + (-)^k \zeta(k-1)c_1$ for $k \geq 3$

For implementation, we use **(C)** for $x > 90$ and **(D)** for $x < 2$ and otherwise use **(B)** repeatedly to bring $x < 2$. This seems to give 15 digits precision for all $x > 0$. Values of $\zeta(k) - 1$ are precomputed at high-precision and stored in a table — 100 values are enough. For $x < 0$ we use **(A)** to transform to a positive value, but naive implementation gives poor results for $-90 < x < 0$.

Note that the above relations apply for complex z also, so can be the basis for complex Γ implementation.

4.2 Factorial, Pochhammer symbol

$$n! = \Gamma(n+1)$$

$$(\alpha)_\nu = \alpha^{\uparrow\nu} = \frac{\Gamma(\alpha+\nu)}{\Gamma(\alpha)}$$

$$\alpha^{\downarrow\nu} = \frac{\Gamma(\alpha)}{\Gamma(\alpha-\nu)}$$

Thus $(\alpha)_0 = 1$ and $(\alpha)_{n+1} = (\alpha+n) \cdot (\alpha)_n$. $n!!$, $n!!!$, \dots (+logs), log-factorial, rising/falling_factorial, $x^n/n!$, ratio of pochhammers $\frac{(a_1)_n \dots (a_p)_n}{(b_1)_n \dots (b_q)_n}$, integer/general pochhammers, etc.

4.3 Incomplete gamma functions

$$\gamma(z, \xi) = \int_0^\xi t^{z-1} e^{-t} dt \quad \gamma(z, \xi)$$

$$\gamma^*(z, \xi) = z^{\cdots} \gamma(z, \xi)$$

$$P(z, \xi) = \frac{\gamma(z, \xi)}{\Gamma(z)} \quad P(z, \xi)$$

$$Q(z, \xi) = 1 - P(z, \xi) = \frac{\Gamma(z, \xi)}{\Gamma(z)} \quad Q(z, \xi)$$

$$\Gamma(z, \xi) = \Gamma(z) - \gamma(z, \xi) = \int_\xi^\infty t^{z-1} e^{-t} dt \quad \Gamma(z, \xi)$$

For a fixed ξ , $\gamma(z, \xi)$ is a meromorphic function of z with simple poles at $z = 0, -1, -2, \dots$, while $\Gamma(z, \xi)$ is an entire function of z .

Inverse functions ...

Various potentially useful relations

(A) For $a \neq -1, -2, \dots$,

$$\Gamma(a, z) = \Gamma(a) - \sum_{n=0}^{\infty} \frac{\Gamma(a) e^{-z} z^{n+a}}{\Gamma(a+n+1)} = \Gamma(a) - \sum_{n=0}^{\infty} \frac{(-)^n z^{n+a}}{(a+n)n!}$$

(B) As $z \rightarrow \infty$,

$$\Gamma(a, z) \sim z^{a-1} e^{-z} \sum_{n=0}^{\infty} \frac{\Gamma(a)}{\Gamma(a-n) z^n}$$

(C) Continued fraction expansion: (best when $a \gg z$)

$$\Gamma(a, z) = \frac{e^{-z} z^a}{z+} \frac{1-a}{1+} \frac{1}{z+} \frac{2-a}{1+} \frac{2}{z+} \dots$$

(C') Equivalent fraction (converges for $\Re z > 0$), where $v = z^{-1}$

$$e^z z^{1-a} \Gamma(a, z) = \frac{1}{1+} \frac{(1-a)v}{1+} \frac{v}{1+} \frac{(2-a)v}{1+} \frac{2v}{1+} \dots$$

(D) For $n \geq 0$; (though claimed to be of limited use unless $a \sim -n$)

$$\Gamma(a+n+1, z) = \Gamma(a+n, z) + \frac{\Gamma(a) \exp(-z) z^{n+a}}{\Gamma(a+n+1)}$$

4.4 Digamma functions and related

The digamma function is defined via

$$\psi(z) = \frac{d}{dz} \log \Gamma(z) = \frac{\Gamma'(z)}{\Gamma(z)}$$

meromorphic with simple poles at $z = 0, -1, -2, \dots$

And the polygamma function (for $n = 1, 2, 3, \dots$)

$$\psi^{(n)}(z) = \frac{d^n}{dz^n} \psi(z) = \frac{d^{n+1}}{dz^{n+1}} \ln \Gamma(z) = (-)^{n+1} \int_0^\infty \frac{t^n e^{-zt}}{1 - e^{-t}} dt$$

which is meromorphic and single-valued with poles of order $n + 1$ at $z = -m$ ($m = 0, 1, 2, \dots$).

$$\psi^{(n)}(z + 1) = \psi^{(n)}(z) + (-)^n n! z^{-n-1}$$

$$\psi^{(n)}(1 - z) - (-)^n \psi^{(n)}(z) = (-)^n \pi \frac{d^n}{dz^n} \cot(\pi z)$$

$$\psi^{(n)}(1 + z) = (-)^{n+1} \sum_{k=n}^{\infty} (-)^k k! \zeta(k + 1) \frac{z^{k-n}}{(k - n)!}$$

$$\psi^{(n)}(z) = (-)^{n+1} n! \sum_{k=0}^{\infty} (z + k)^{-n-1} \quad (z \neq 0, -1, -2, \dots)$$

Using Euler-Maclaurin summation with this last series works quite well.

This approach for digamma works pretty well, but get loss of precision as $x \rightarrow^- 2$ or $x \rightarrow^+ 0$.

(A) Use the following to reduce the (real) argument to get $0.5 < z \leq 1.5$

$$\psi(z + 1) = \frac{1}{z} + \psi(z)$$

(B) Then use this series for the resulting z .

$$\psi(1 + z) = -\gamma + \sum_{k=2}^{\infty} (-)^k \zeta(k) z^{k-1}$$

Remark: for reflection of polygamma function, we need $\frac{d^n}{dz^n} \cot(\pi z)$. From Mathematica we have the following table: perhaps some pattern can be gleaned for arbitrary n ... [TODO: solve this puzzle]

n	$\frac{(-)^n}{\csc(x)^{n+1}} \frac{d^n}{dx^n} \cot(x)$
0	$\cos(x)$
1	1
2	$2 \cos(x)$
3	$2 \cos(2x) + 4$
4	$22 \cos(x) + 2 \cos(3x)$
5	$52 \cos(2x) + 2 \cos(4x) + 66$
6	$604 \cos(x) + 114 \cos(3x) + 2 \cos(5x)$
7	$2382 \cos(2x) + 240 \cos(4x) + 2 \cos(6x) + 2416$
8	$31238 \cos(x) + 8586 \cos(3x) + 494 \cos(5x) + 2 \cos(7x)$
9	$176468 \cos(2x) + 29216 \cos(4x) + 1004 \cos(6x) + 2 \cos(8x) + 156190$
10	$2620708 \cos(x) + 910384 \cos(3x) + 95680 \cos(5x) + 2026 \cos(7x) + 2 \cos(9x)$
11	$19476228 \cos(2x) + 4406976 \cos(4x) + 305274 \cos(6x) + 4072 \cos(8x) + 2 \cos(10x) + 15724248$
12	$325024572 \cos(x) + 132636948 \cos(3x) + 20375370 \cos(5x) + 956542 \cos(7x) + 8166 \cos(9x) + 2 \cos(11x)$

4.5 Beta functions

$$B(a, b) = \int_0^1 t^{a-1} (1-t)^{b-1} dt = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} \quad B(a, b)$$

beta

4.6 Incomplete beta functions

$$B_\xi(a, b) = \int_0^\xi t^{a-1} (1-t)^{b-1} dt \quad B_\xi(a, b)$$

$$I_\xi(a, b) = \frac{B_\xi(a, b)}{B(a, b)} \quad I_\xi(a, b)$$

+complementary incomplete beta (& scaled)

Gil-Segura-Temme book gives the continued fraction

$$B_x(p, q) = \frac{x^p(1-x)^q}{p} \left(\frac{1}{1+} \frac{d_1}{1+} \frac{d_2}{1+} \frac{d_3}{1+} \dots \right)$$

where

$$d_{2n+1} = -x \frac{(p+n)(p+q+n)}{(p+2n)(p+2n+1)} \quad d_{2n+2} = x \frac{(n+1)(q-n-1)}{(p+2n+1)(p+2n+2)}$$

with the note that when $p, q > 1$ the best numerical results are obtained when $x \leq x_0 = (p-1)/(p+q-2)$ and for $x_0 < x \leq 1$ you should use the reflection $B_x(p, q) = B(p, q) - B_{1-x}(q, p)$. Note also that for this continued fraction, the convergents C_{4n} and C_{4n+1} are less than and the convergents C_{4n+2} and C_{4n+3} are greater than the value of the continued fraction. [This works quite well (with a few domain issues for complex/negative parameters).]

5 Error and related functions

Dawson's integral, Fadeeva function (zeros?) [normal cdf, etc.] Black-Scholes/Implied-Volatility core: $e^\alpha \Phi[\alpha/s + s/2] - \Phi[\alpha/s - s/2]$, **erf**, **erfc**, **inverse_erf**, **inverse_erfc**, **erfc_star**, **Dawson**, **Fadeeva**,

Iterated error functions $i^n = \int i^{n-1} \operatorname{erfc}$

$$\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt \quad z \in \mathbb{C}$$

$$\operatorname{erfc}(z) = 1 - \operatorname{erf}(z) \quad z \in \mathbb{C}$$

$$\operatorname{erfc}^*(z) = e^{z^2} \operatorname{erfc}(z)$$

$$\operatorname{erfi}(z) = -i \operatorname{erf}(iz)$$

(useful for $z \rightarrow \infty$, $z > 0$)

$$\mathbf{erf}(z) = \operatorname{erf}(z)$$

$$\mathbf{erfc}(z) = \operatorname{erfc}(z)$$

An approximation from A&S is

$$\operatorname{inverf}(x) = t + \frac{1}{3}t^3 + \frac{7}{30}t^5 + \frac{127}{630}t^7 + \dots$$

where $t = x\sqrt{\pi}/2$. This can give the first guess to an iterative root-finder (such as Halley's method).

Note that Lebedev defines $\Phi(z)$ = our erf and $\operatorname{Erf}(z) = \int_0^z e^{-t^2} dt$. He also defines $F(z) = e^{-z^2} \int_0^z e^{u^2} du$.

$$\operatorname{erf}(ix) = \frac{2x\hat{i}}{\sqrt{\pi}} \sum_{n=0}^{\infty} (-1)^n \left(i_{2n}^{(1)}(x^2) + i_{2n+1}^{(1)}(x^2) \right) \quad x \in \mathbb{R}$$

- A series that works reasonably well for small z is the following; note that we ensure $z \geq 0$ by using the fact that $\operatorname{erf}(-z) = -\operatorname{erf}(z)$ (to avoid cancellation)

$$\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} e^{-z^2} \sum_{n=0}^{\infty} \frac{2^n z^{2n+1}}{1 \cdot 3 \cdot 5 \cdots (2n+1)}$$

- A continued fraction that works well the following (evaluated with, say, the Wallis recurrence)

$$\operatorname{erfc}(z) = \frac{e^{-z^2}}{\sqrt{\pi}} \frac{2z}{z^2+1} - \frac{1 \cdot 2}{z^2+5} \frac{3 \cdot 4}{z^2+9} \frac{5 \cdot 6}{z^2+13} \cdots$$

- An asymptotic expansion that generally underperforms the continued fraction is

$$\operatorname{erfc}(z) \sim \frac{e^{-z^2}}{z\sqrt{\pi}} \sum_{m=0}^{\infty} (-)^m \frac{1 \cdot 3 \cdot 5 \cdots (2m-1)}{(2z^2)^m}$$

- Marsaglia [Marsaglia] points out (for the normal cdf) that given the value of $\operatorname{erfc}(x_0)$ at a point x_0 , there is a simple recurrence relation to generate the coefficients of the Taylor series around that point, that is:

$$\operatorname{erf}^{(n+2)}(z) = -2z \operatorname{erf}^{(n+1)}(z) - 2n \operatorname{erf}^{(n)}(z) \quad n \geq 0$$

- A reasonable approach is to use the continued fraction for $|x| > 3$, the series for $|x| < 1$, and the local Taylor expansions centered at $\pm 1\frac{1}{2}, \pm 2, \pm 2\frac{1}{2}$ for other points. (Using a table to store the “seeds” for those select points.)

Other notes:

$$\operatorname{erf}(x + iy) = \operatorname{erf}(x) + \frac{e^{-x^2}}{2\pi x} [(1 - \cos(2xy)) + i \sin(2xy)] + \frac{2}{\pi} e^{-x^2} \sum_{n=1}^{\infty} \frac{e^{-n^2/4}}{n^2 + 4x^2} [f_n(x, y) + i g_n(x, y)] + \varepsilon$$

with $\varepsilon \sim 10^{-16} \cdot |\operatorname{erf}(z)|$ where $f_n(x, y) = 2x - 2x \cosh(ny) \cos(2xy) + n \sinh(ny) \sin(2xy)$ and $g_n(x, y) = 2x \cosh(ny) \sin(2xy) + n \sinh(ny) \cos(2xy)$. This seems to work pretty well for small(ish) z , but gives NaN for, say, $z = 17(1 + i)$. (Note that $x = 0$ is bad, though this case reduces to...)

5.1 Fadeeva and Dawson integrals

Dawson’s integral:

$$\operatorname{F}(x) = e^{-x^2} \int_0^x e^{t^2} dt = -i \frac{\sqrt{\pi}}{2} e^{-z^2} \operatorname{erf}(iz) = i \frac{\sqrt{\pi}}{2} (e^{-z^2} - \operatorname{w}(z))$$

Generalized Dawson’s integral:

$$\operatorname{F}_p(x) = e^{-x^p} \int_0^x e^{t^p} dt$$

Fadeeva’s integral:

$$\operatorname{w}(z) = e^{-z^2} \operatorname{erfc}(-iz) = e^{-z^2} (1 + \frac{2i}{\sqrt{\pi}} \int_0^z e^{t^2} dt)$$

Series:

$$\begin{aligned}
 F(x) &= x \frac{\sqrt{\pi}}{2} \sum_{k=0}^{\infty} (-)^k \frac{x^{2k}}{\Gamma(k + 3/2)} \\
 &= x \sum_{k=0}^{\infty} (-)^k \frac{x^{2k}}{(3/2)_k} \\
 &= \sum_{n=0}^{\infty} (-)^n \frac{2^n x^{2n+1}}{(2n+1)!!}
 \end{aligned}$$

Series of spherical Bessel functions

$$\begin{aligned}
 F(x) &= e^{-x^2} x \sum_{n=0}^{\infty} (-)^n \left(i_{2n}^{(1)}(x^2) + i_{2n+1}^{(1)}(x^2) \right) \\
 &= e^{-x^2} x (i_0 + i_1 - i_2 - i_3 + i_4 + i_5 - \dots) \\
 &= e^{-x^2} x \left(\frac{3}{z} i_1 + \frac{5}{z} i_2 + \frac{11}{z} i_5 + \frac{13}{z} i_6 + \frac{19}{z} i_9 + \frac{21}{z} i_{10} + \dots \right)
 \end{aligned}$$

where we use the recurrence relation $i_{n-1} - i_{n+1} = \frac{2n+1}{z} i_n$ for the last equation. Unfortunately, this approach is not numerically stable and we lose much accuracy from bad cancellation.

Continued fractions:

$$F(z) = \frac{z}{1+} \frac{2z^2}{3-} \frac{4z^2}{5+} \frac{6z^2}{7-} \frac{8z^2}{9+} \dots$$

which is good for $|z|$ smaller.

$$F(z) = \frac{z}{1+2z^2-} \frac{4z^2}{3+2z^2-} \frac{8z^2}{5+2z^2-} \frac{12z^2}{7+2z^2-} \dots$$

which is good for $|z|$ larger.

5.2 Fresnel integrals

Need *zeros* also

$$\begin{aligned}
 C(z) &= \int_0^z \cos(t^2 \pi/2) dt \\
 S(z) &= \int_0^z \sin(t^2 \pi/2) dt
 \end{aligned}$$

`Fresnel_C`, `Fresnel_S`

We have the series expansions

$$\begin{aligned}
 C(z) &= \sum_{n=0}^{\infty} (-)^n \frac{(\pi/2)^{2n}}{(2n)!(4n+1)} z^{4n+1} = \sum_{n=0}^{\infty} j_{2n}(z^2 \pi/2) \\
 S(z) &= \sum_{n=0}^{\infty} (-)^n \frac{(\pi/2)^{2n+1}}{(2n+1)!(4n+3)} z^{4n+3} = \sum_{n=0}^{\infty} j_{2n+1}(z^2 \pi/2)
 \end{aligned}$$

(for using the spherical bessel expansions, should extract the whole sequence of needed values for efficiency...) Note that DLMF and A&S include a z multiplier on the bessel expansions but numeric tests seem to indicate that that is wrong!

Use the two series above as well as asymptotic expansion...

“modified Fresnel? (see Z & J)”

Note: in Lebedev’s notation, we have $\Phi(\sqrt{i}x)/\sqrt{i} = \frac{2}{\sqrt{\pi}} \int_0^x \cos u^2 du - i \frac{2}{\sqrt{\pi}} \int_0^x \sin u^2 du$

$$\begin{aligned}\text{Ci}(z) &= \int_{\infty}^z \frac{\cos t}{t} dt = \gamma + \log(z) + \int_0^z \frac{\cos t - 1}{t} dt \\ \text{Si}(z) &= \int_0^z \frac{\sin t}{t} dt\end{aligned}$$

$\text{Si}(z)$ is an entire function, while $\text{Ci}(z)$ is analytic in the split plane (deleting the negative real axis). Note that $\text{Ei}(\hat{i}x) = \text{Ci}(x) - \hat{i}(\frac{\pi}{2} - \text{Si}(x))$ for $x > 0$.

When $|\text{ph } z| < \pi/2$, we have

$$\begin{aligned}\text{Ci}(z) &= -\frac{1}{2} (\text{E}_1(\hat{i}z) + \text{E}_1(-\hat{i}z)) \\ \text{Si}(z) &= \frac{\hat{i}}{2} (\text{E}_1(-\hat{i}z) - \text{E}_1(\hat{i}z)) + \frac{\pi}{2}\end{aligned}$$

Using these with continued-fraction expansion for $\text{E}_1(z) = e^{-z}/(z + 1/(1 + 1/(z + 2/(1 + 2/(z + \dots))))))$ works well. (Note that for real x , we have $\text{Ci}(x) = -\Re \text{E}_1(\hat{i}x)$.)

[Move much of this to Exponential Integrals section]

$$\text{Cin}(z) = \int_0^z \frac{1 - \cos t}{t} dt$$

thus $\text{Ci}(z) = -\text{Cin}(z) + \ln z + \gamma$

Series expansions:

$$\begin{aligned}\text{Ci}(z) &= \gamma + \ln(z) + \sum_{n=1}^{\infty} (-)^n \frac{z^{2n}}{(2n)!(2n)} \\ \text{Si}(z) &= \sum_{n=0}^{\infty} (-)^n \frac{z^{2n+1}}{(2n+1)!(2n+1)} \\ \text{Si}(z) &= z \sum_{n=0}^{\infty} j_n^2(z/2)\end{aligned}$$

which works well (and can use recurrence for fast computation...)

$$\text{Ci}(z) = \sum_{n=0}^{\infty} a_n j_n^2(z/2)$$

where $a_n = (2n+1)(1 - (-)^n + \psi(n+1) - \psi(1))$

$$\text{Ei}(x) = \gamma + \ln|x| + \sum_{n=0}^{\infty} (-)^n (x - a_n) i^{(1)}(x/2)^2 \quad (x \neq 0)$$

where $a_n = (2n+1)(1 - (-)^n + \psi(n+1) - \psi(1))$

$$\text{Ein}(x) = ze^{-z/2} \left(i_0^{(1)}(z/2) + \sum_{n=1}^{\infty} \frac{2n+1}{n(n+1)} i_n^{(1)}(z/2) \right)$$

Also, $\text{E}_1(z) = \Gamma(0, z)$ and $\text{E}_1(z) = e^{-z}U(1, 1, z)$

Asymptotic expansions ($|z| = 25$ seems to be a reasonable transition point from the series to asymptotic)

$$\begin{aligned}\text{Si}(z) &= \frac{\pi}{2} - \cos(z)f(z) - \sin(z)g(z) \\ \text{Ci}(z) &= \sin(z)f(z) - \cos(z)g(z) \\ f(z) &\sim \frac{1}{z} \left(1 - \frac{2!}{z^2} + \frac{4!}{z^4} - \frac{6!}{z^6} + \dots \right) \\ g(z) &\sim \frac{1}{z^2} \left(1 - \frac{3!}{z^2} + \frac{5!}{z^4} - \frac{7!}{z^6} + \dots \right)\end{aligned}$$

For $x > 0$ we have the following (actually for any z in the slit plane $\mathbb{C} \setminus (-\infty, 0]$):

$$\text{Shi}(x) = (\text{Ei}(z) + \text{E}_1(x))/2$$

$$\text{Chi}(x) = (\text{Ei}(z) - \text{E}_1(x))/2$$

We have

$$\text{li}(z) = \text{Ei}(\ln z)$$

Asymptotic expansion for Fresnel integrals

$$\begin{aligned} C(x) &= \text{sgn}(x) \frac{1}{2} + \frac{1}{\pi x} (f(x) \sin(x^2 \pi/2) - g(x) \cos(x^2 \pi/2)) \\ S(x) &= \text{sgn}(x) \frac{1}{2} - \frac{1}{\pi x} (f(x) \cos(x^2 \pi/2) + g(x) \sin(x^2 \pi/2)) \end{aligned}$$

where

$$\begin{aligned} f(x) &\sim 1 + \sum_{n=1}^{\infty} (-1)^n \frac{1 \cdot 3 \cdot 5 \cdots (4n-1)}{(\pi x^2)^{2n}} \\ g(x) &\sim \sum_{n=0}^{\infty} (-1)^n \frac{1 \cdot 3 \cdot 5 \cdots (4n+1)}{(\pi x^2)^{2n+1}} \end{aligned}$$

[quality of this needs to be validated...]

5.2.1 Arctangent integral

$$\text{atanint}(z) = \int_0^z \frac{\text{atan}(t)}{t} dt$$

then since $\text{atan}(z) = z - z^3/3 + z^5/5 - \cdots$ (for $|z| \leq 1$, $z \neq \pm i$), we have

$$\text{atanint}(z) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)^2}$$

(compare dilogarithm...)

In terms of the Lerch Φ function we have

$$\text{atanint}(z) = \frac{z}{4} \sum_{n=0}^{\infty} \frac{(-z^2)^n}{(n+1/2)^2} = \frac{z}{4} \Phi[-z^2, 2, 1/2]$$

We have the asymptotic series

$$\text{atanint}(z) = \frac{\pi}{2} \ln z + \sum_{n=0}^{\infty} \frac{(-1)^n}{z^{2n+1} (2n+1)^2} \quad |z| \geq 1, \Re z \neq 0$$

(from similar series for atan)

We can expand in Taylor series at 1. We get

$$\begin{aligned} \text{ataint}(1+x) &= C + \frac{\pi}{4}x + \left(\frac{1}{2} - \frac{\pi}{4}\right) \frac{x^2}{2!} + \left(\frac{\pi}{4} - \frac{3}{2}\right) \frac{x^3}{3!} + \left(5 - \frac{3\pi}{2}\right) \frac{x^4}{4!} \\ &\quad + (6\pi - 20) \frac{x^5}{5!} + (97 - 30\pi) \frac{x^6}{6!} + (180\pi - 567) \frac{x^7}{7!} + (3924 - 1260\pi) \frac{x^8}{8!} + \cdots \end{aligned}$$

where $C = 0.9159655942 \dots$ is Catalan's constant.

One can attempt Euler-Maclaurin expansion in the series, but you end up with $\Gamma(-1, x)$ term for the integral (which can be written in terms of $\text{Ei}(x)$), but the computation becomes more expensive... (integral term in EM is $\frac{-\ln(z)}{4z} \Gamma(-1, (4n+1) \ln z)$ and we have $\Gamma(-1, x) = \text{Ei}(-x) + \frac{e^{-x}}{x} + \frac{1}{2}(\ln(-1/x) - \ln(-x)) + \ln x$.)

To implement for real x , use series, asymptotic series, reflection, and local series at 1.

5.3 Iterated error functions

Define

$$\begin{aligned} i^{-1} \operatorname{erfc}(z) &= \frac{2}{\sqrt{\pi}} e^{-z^2} \\ i^0 \operatorname{erfc}(z) &= \operatorname{erfc}(z) \\ i^n \operatorname{erfc}(z) &= \int_z^\infty i^{n-1} \operatorname{erfc}(t) dt \end{aligned}$$

We have the recurrence

$$i^n \operatorname{erfc}(z) = -\frac{z}{n} i^{n-1} \operatorname{erfc}(z) + \frac{1}{2n} i^{n-2} \operatorname{erfc}(z)$$

6 Bessel, Hankel, Airy functions

Need *zeros* also

`Jn`, `Jnu`, `Yn`, `Ynu`, `In`, `Inu`, `Kn`, `Knu`, `dJn`, ..., `Integrate_Jn`, ... `jn`, `yn`, `in`, `kn`, ..., (`array versions...`) — real / imaginary / complex arguments, integer / real / imaginary / complex parameters
 $\Lambda_\nu(x) = ?$, “Riccati Bessel functions”?

6.1 Bessel functions

Bessel functions are also known as “cylinder functions”

Bessel’s differential equation $z^2 y'' + zy' + (z^2 - \nu^2)y = 0$.

Bessel function of the first kind, regular at 0; if $\nu \in \mathbb{Z}$ then we can have any $z \in \mathbb{C}$, but for arbitrary $\nu \in \mathbb{C}$ we need $|\arg z| < \pi$.

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k+\nu}}{k!(k+\nu)!}$$

Note that

$$J_\nu(z) = \frac{1}{\pi} \int_0^\pi \cos(z \sin \vartheta - \nu \vartheta) d\vartheta - \frac{\sin \nu \pi}{\pi} \int_0^\infty e^{-z \sinh t - \nu t} dt$$

Bessel function of the second kind, irregular at 0:

$$Y_\nu(z) = \frac{J_\nu(z) \cos \nu \pi - J_{-\nu}(z)}{\sin \nu \pi}$$

Some people use the notation $N_\nu(z) = Y_\nu(z)$. For $\nu = n \in \mathbb{Z}$, define $Y_n(z) = \lim_{\nu \rightarrow n} Y_\nu(z)$.

6.2 Modified Bessel functions

Hyperbolic or modified Bessel function (of the first kind), regular at 0:

$$I_\nu(z) = e^{-\nu \pi i/2} J_\nu(\hat{i}z) \quad z \in \mathbb{C}, \nu \in \mathbb{C}$$

Hyperbolic or modified Bessel function (of the third kind), irregular at 0, also called Bassett’s function or Macdonald’s function:

$$K_\nu(z) = xxx \quad z \in \mathbb{C} \setminus \mathbb{R}_{\leq 0}, \nu \in \mathbb{C}$$

6.3 Incomplete Bessel functions

$$I_\nu(z; \xi) = \dots$$

6.4 Spherical Bessel functions

$$j_n(z) = \sqrt{\frac{\pi}{2z}} J_{n+1/2}(z)$$

$$y_n(z) = \sqrt{\frac{\pi}{2z}} Y_{n+1/2}(z)$$

$$i_n^{(1)}(z) =$$

$$i_n^{(2)}(z) =$$

$$k_n(z) =$$

$$h_n^{(1)}(z) =$$

$$h_n^{(2)}(z) =$$

Some explicit formulæ for spherical Bessel functions: let $a_k(n + \frac{1}{2}) = \begin{cases} \frac{(n+k)!}{2^k k! (n-k)!} & k = 0, 1, \dots, n \\ 0 & k > n \end{cases}$

$$j_n(z) = \sin(z - n\frac{\pi}{2}) \sum_{k=0}^{\lfloor n/2 \rfloor} (-)^k \frac{a_{2k}(n + \frac{1}{2})}{z^{2k+1}} + \cos(z - n\frac{\pi}{2}) \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} (-)^k \frac{a_{2k+1}(n + \frac{1}{2})}{z^{2k+2}}$$

(The Bessel polynomial of degree n is given by $\sum_{k=0}^n a_k(n + \frac{1}{2}) z^{n-k}$.)

$$j_0(z) = \frac{\sin z}{z}$$

$$j_1(z) = \frac{\sin z}{z^2} - \frac{\cos z}{z} = \sum_{n=0}^{\infty} (-)^{n+1} \frac{z^{2n+1}}{(2n+1)!(2n+3)}$$

$$y_0(z) = -\frac{\cos z}{z}$$

$$y_1(z) = -\frac{\cos z}{z^2} - \frac{\sin z}{z}$$

$$k_0(z) = \frac{\pi}{2} e^{-z} \frac{1}{z}$$

$$k_1(z) = \frac{\pi}{2} e^{-z} \left(\frac{1}{z} + \frac{1}{z^2} \right)$$

$$i_0^{(1)}(z) = \frac{\sinh z}{z} = \sum_{n=0}^{\infty} \frac{z^{2n}}{(2n+1)!}$$

$$i_1^{(1)}(z) = -\frac{\sinh z}{z^2} + \frac{\cosh z}{z} = \frac{1}{z} \sum_{n=0}^{\infty} \frac{2n}{(2n+1)!} z^{2n}$$

$$i_0^{(2)}(z) = \frac{\cosh z}{z}$$

$$i_1^{(2)}(z) = -\frac{\cosh z}{z^2} + \frac{\sinh z}{z}$$

Reflection relations:

$$j_n(-z) = (-)^n j_n(z)$$

$$y_n(-z) = (-)^{n+1} y_n(z)$$

$$i_n^{(1)}(-z) = (-)^n i_n^{(1)}(z)$$

$$i_n^{(2)}(-z) = (-)^{n+1} i_n^{(2)}(z)$$

$$k_n(-z) = -\frac{\pi}{2} \left(i_n^{(1)}(-z) i_n^{(2)}(-z) \right)$$

Recurrence relations: Let $f_n(z)$ denote $j_n(z)$, $y_n(z)$, or $h_n^{(j)}(z)$, then

$$f_{n-1}(z) + f_{n+1}(z) = \frac{2n+1}{z} f_n(z)$$

and

$$f'_n(z) = f_{n-1}(z) - \frac{n+1}{z} f_n(z)$$

For y_n , use forward. For j_n , use forward when $|z| > n$ and backward when $|z| < n$.

Let $g_n(z)$ denote $(-)^n k_n(z)$, $i_n^{(j)}(z)$, then

$$g_{n-1}(z) - g_{n+1}(z) = \frac{2n+1}{z} g_n(z)$$

and

$$g'_n(z) = g_{n-1}(z) - \frac{n+1}{z} g_n(z)$$

For k_n , use forward. For $i_n^{(2)}$, use forward. For $i_n^{(1)}$, use forward when $|z| > n$ and backward when $|z| < n$.

For scaling in recurrence, can use:

$$\sum_{n=0}^{\infty} (2n+1) j_n^2(z) = 1$$

(or can use, for example, $j_0(z) = \sin z/z$.)

6.5 Integral Bessel functions

$$\text{Ji}_\nu(z) = \int_0^z J_\nu(t) dt$$

$$\text{Ii}_\nu(z) = \int_0^z I_\nu(t) dt$$

etc.

Note that Lebedev defines $\text{Ji}_\nu(z) = \int_0^z \frac{J_\nu(t)}{t} dt$.

6.6 Hankel, (Whittaker?)

Hankel functions are also known as Bessel functions of the third kind:

$$H_\nu^{(1)}(z) = J_\nu(z) + i Y_\nu(z)$$

$$H_\nu^{(2)}(z) = J_\nu(z) - i Y_\nu(z)$$

6.7 Airy functions

Solutions of Airy DE $y'' - zy = 0$. [Need derivatives also]

$$\text{Ai}(z) \quad \text{Ai}(z) = \frac{1}{\pi} \int_0^\infty \cos(t^3/3 + xt) dt = \frac{\sqrt{z/3}}{\pi} K_{1/3}(\zeta) = \frac{\sqrt{z}}{3} (I_{-1/3}(\zeta) - I_{1/3}(\zeta))$$

$$\text{Bi}(z) \quad \text{Bi}(z) = \frac{1}{\pi} \int_0^\infty e^{-t^3/3+xt} \sin(t^3/3 + xt) dt = \frac{\sqrt{z}}{3} (I_{-1/3}(\zeta) + I_{1/3}(\zeta))$$

where $\zeta = (2/3)z^{3/2}$.

[For positive z use continued fraction from power series??]

We also have

$$\text{Ai}(-z) = \frac{\sqrt{z}}{3} (J_{1/3}(\zeta) - J_{-1/3}(\zeta))$$

$$\text{Bi}(-z) = \sqrt{\frac{z}{3}} (J_{-1/3}(\zeta) - J_{1/3}(\zeta))$$

Plus scaled versions, derivatives, integrals, ... (joint versions)

Remark: one could attempt to do "Taylor stepping" (repeated expansion of power-series as analytic continuation...) One can derive a recurrence for the derivatives at an arbitrary point from the ordinary differential equation satisfied by Airy functions $w'' - zw = 0$ gives $w^{(n+3)} = z \cdot w^{(n+1)} + n \cdot w^{(n)}$. However, in practice this works poorly: backwards is unstable, forwards gives low accuracy, relative error is $O(h^3)$ for step-sizes of h , etc.

Atlas [ACMF] recommends to compute these as $\text{Ai}(z) = c_1 f(x) - c_2 g(x)$, $\text{Bi}(z) = \sqrt{3}[c_1 f(x) + c_2 g(x)]$, where $c_1 = (3^{2/3}\Gamma(2/3))^{-1}$, $c_2 = (3^{1/3}\Gamma(1/3))^{-1}$, and

$$f(x) = \sum_{k=0}^{\infty} \frac{1 \cdot 4 \cdot 7 \cdots (3k-2)}{(3k)!} x^{3k} = 1 + \frac{x^3}{3!} + 4 \frac{x^6}{6!} + \cdots$$

$$g(x) = \sum_{k=0}^{\infty} \frac{2 \cdot 5 \cdot 8 \cdots (3k-1)}{(3k+1)!} x^{3k} = x(1 + 2 \frac{x^3}{4!} + 2 \cdot 5 \frac{x^6}{7!} + \cdots)$$

For $x \gg 0$, use the asymptotics

$$\text{Ai}(x) \sim \frac{e^{-\zeta}}{2\sqrt{\pi x^{1/2}}} \left(\sum_{k=0}^{\infty} (-)^k \frac{t_k}{\zeta^k} \right)$$

$$\text{Bi}(x) \sim \frac{e^{\zeta}}{2\sqrt{\pi x^{1/2}}} \left(\sum_{k=0}^{\infty} \frac{t_k}{\zeta^k} \right)$$

where $t_k = \frac{(2k+1)(2k+3)\cdots(6k-1)}{216^k k!}$ (for $k > 0$), $t_0 = 1$.

For $x \ll 0$, use the asymptotics

$$\text{Ai}(x) \sim \frac{1}{\sqrt{\pi x^{1/2}}} [\sin(\zeta + \pi/4)F(\zeta) - \cos(\zeta + \pi/4)G(\zeta)]$$

$$\text{Bi}(x) \sim \frac{1}{\sqrt{\pi x^{1/2}}} [\cos(\zeta + \pi/4)F(\zeta) + \sin(\zeta + \pi/4)G(\zeta)]$$

where $F(\zeta) = \sum_{k=0}^{\infty} (-)^k c_{2k}/\zeta^{2k}$ and $G(\zeta) = \sum_{k=0}^{\infty} (-)^k c_{2k+1}/\zeta^{2k+1}$. (In practice, he solves for F, G in terms of Ai, Bi and computes a rational approximation to them.)

6.7.1 Scorer functions

$\text{Gi}(z)$ is solution of inhomogeneous Airy DE $y'' - zy = -\frac{1}{\pi}$ with $\text{Gi}(0) = \text{Bi}(0)/3$, $\text{Gi}'(0) = \text{Bi}'(0)/3$, $\text{Gi}(z)$

$\text{Hi}(z)$ is solution of inhomogeneous Airy DE $y'' - zy = +\frac{1}{\pi}$ with $\text{Hi}(0) = 2\text{Bi}(0)/3$, $\text{Hi}'(0) = 2\text{Bi}'(0)/3$, $\text{Hi}(z)$

$$\text{Gi}(z) = \text{Bi}(z) \int_z^{\infty} \text{Ai}(t) dt + \text{Ai}(z) \int_0^z \text{Bi}(t) dt$$

$$\text{Hi}(z) = \text{Bi}(z) \int_{-\infty}^z \text{Ai}(t) dt - \text{Ai}(z) \int_{-\infty}^z \text{Bi}(t) dt$$

For small $|z|$ we can use series

$$\text{Gi}(z) = \frac{3^{-2/3}}{\pi} \sum_{k=0}^{\infty} \cos(\pi \frac{2k-1}{3}) \Gamma(\frac{k+1}{3}) \frac{(3^{1/3}z)^k}{k!}$$

$$\text{Hi}(z) = \frac{3^{-2/3}}{\pi} \sum_{k=0}^{\infty} \Gamma\left(\frac{k+1}{3}\right) \frac{(3^{1/3}z)^k}{k!}$$

Note that from periodicity of \cos , we don't need to compute for each term. Similarly, $\Gamma(n/3)$ can be computed via recurrence from $\Gamma(1/3)$ and $\Gamma(2/3)$.

For Mathematica use, these integral forms were useful:

$$\text{Gi}(z) = \frac{1}{\pi} \int_0^{\infty} \sin\left(\frac{t^3}{3} + xt\right) dt \quad x \in \mathbb{R}$$

$$\text{Hi}(z) = \frac{1}{\pi} \int_0^{\infty} e^{-t^3/3+zt} dt$$

Another integral form (useful for intermediate $|z|$?) is

$$\text{Gi}(z) = -\frac{1}{\pi} \int_0^{\infty} e^{-t^3/3-zt/2} \cos\left(\frac{\sqrt{3}}{2}zt + \frac{2}{3}\pi\right) dt$$

For large $|z| \rightarrow \infty$ we have asymptotic forms

$$\text{Gi}(z) \sim \frac{1}{\pi z} \sum_{k=0}^{\infty} \frac{(3k)!}{k!(3z^3)^k} \quad |\text{ph } z| \leq \frac{\pi}{3} - \delta$$

$$\text{Hi}(z) \sim -\frac{1}{\pi z} \sum_{k=0}^{\infty} \frac{(3k)!}{k!(3z^3)^k} \quad |\text{ph } -z| \leq \frac{2}{3}\pi - \delta$$

For other phase ranges, use the connection formulas

$$\text{Gi}(z) + \text{Hi}(z) = \text{Bi}(z)$$

$$\text{Gi}(z) = \frac{1}{2} e^{\pi i/3} \text{Hi}(ze^{-2\pi i/3}) + \frac{1}{2} e^{-\pi i/3} \text{Hi}(ze^{2\pi i/3})$$

$$\text{Gi}(z) = e^{\mp \pi i/3} \text{Hi}(ze^{\pm 2\pi i/3}) \pm i \text{Ai}(z)$$

$$\text{Hi}(z) = e^{\pm 2\pi i/3} \text{Hi}(ze^{\pm 2\pi i/3}) + 2e^{\mp \pi i/6} \text{Ai}(ze^{\mp 2\pi i/3})$$

Finally a few other integral representations:

$$\text{Gi}(x) = \frac{4x^2}{3^{3/2}\pi} \int_0^{\infty} \frac{K_{1/3}(t)}{\zeta^2 - t^2} dt \quad x > 0, \text{CHECK!}$$

$$\text{Hi}(-z) = \frac{4z^2}{3^{3/2}\pi^2} \int_0^{\infty} \frac{K_{1/3}(t)}{\zeta^2 + t^2} dt \quad |\text{ph } z| < \frac{\pi}{3}$$

where $\zeta = \frac{2}{3}z^{3/2}$

$$\text{Hi } z = \frac{3^{-2/3}}{2\pi^2 i} \int_{-i\infty}^{+i\infty} \Gamma\left(\frac{1}{3} + \frac{1}{3}t\right) \Gamma(-t) (3^{1/3}e^{\pi i}z)^t dt$$

where the integration contour separates the poles of $\Gamma(\frac{1}{3} + \frac{1}{3}t)$ (which are $-1, -4, -7, -10, \dots$) from the poles of $\Gamma(-t)$ (which are $0, 1, 2, 3, \dots$).

6.8 Kelvin functions

Regular at 0

$$\text{ber}_\nu(z) = \sum_{n=0}^{\infty} \frac{(z/2)^{2n+\nu}}{n!(n+\nu)!} \cos((3\nu/4 + n/2)\pi)$$

$$\text{bei}_\nu(z) = \sum_{n=0}^{\infty} \frac{(z/2)^{2n+\nu}}{n!(n+\nu)!} \sin((3\nu/4 + n/2)\pi)$$

Note $\text{ber}_\nu(z) - \hat{i} \text{bei}_\nu(z) = \text{J}_\nu(\hat{i}^{-3/2}z)$. Irregular at 0

$$\text{ker}_\nu(z) =$$

$$\text{kei}_\nu(z) =$$

Note $\text{ker}_\nu(z) \pm \hat{i} \text{kei}_\nu(z) = \hat{i}^{\mp\nu} \text{K}_\nu(\hat{i}^{\pm 1/2}z)$. `ber`, `bei`, `ker`, `kei`, `d_ber`, `d_bei`, ..., `Int_ber`, ...

6.9 Struve functions

$$\mathbf{H}_\nu(z) = \sum_{n=0}^{\infty} (-)^n \frac{(z/2)^{2n+\nu+1}}{(n+1/2)!(n+\nu+1/2)!}$$

$$\mathbf{L}_\nu(z) = -\hat{i}e^{-i\nu\pi/2} \mathbf{H}_\nu(\hat{i}z) = \sum_{n=0}^{\infty} \frac{(z/2)^{2n+\nu+1}}{(n+1/2)!(n+\nu+1/2)!}$$

`Struve_Hn`, `d_Struve_Hn`, ..., `Ln`, `Integrate_Struve_Hn`, ...

From A&S [A&S]:

The Struve function $\mathbf{H}_\nu(z)$ is a particular solution to the non-homogeneous Bessel differential equation $z^2 w'' + zw' + (z^2 - \nu^2)w = \frac{4(z/2)^{\nu+1}}{\sqrt{\pi}\Gamma(\nu+1/2)}$ (which thus has a general solution $w = \alpha \text{J}_\nu(z) + \beta \text{Y}_\nu(z) + \mathbf{H}_\nu(z)$) and $z^{-\nu} \mathbf{H}_\nu(z)$ is an entire function in z .

$$\mathbf{H}_\nu(z) = \left(\frac{z}{2}\right)^{\nu+1} \sum_{k=0}^{\infty} (-)^k \frac{(z/2)^{2k}}{\Gamma(k+3/2)\Gamma(k+\nu+3/2)}$$

thus

$$\mathbf{H}_0(z) = \frac{2}{\pi} \left(z - \frac{z^3}{1^2 \cdot 3^2} + \frac{z^5}{1^2 \cdot 3^2 \cdot 5^2} - \dots \right)$$

and

$$\mathbf{H}_1(z) = \frac{2}{\pi} \left(\frac{z^2}{1^2 \cdot 3} - \frac{z^4}{1^2 \cdot 3^2 \cdot 5} + \frac{z^6}{1^2 \cdot 3^2 \cdot 5^2 \cdot 7} \dots \right)$$

We can also write

$$\mathbf{H}_\nu(z) = \frac{2(z/2)^{\nu+1}}{\sqrt{\pi}\Gamma(\nu+3/2)} {}_1\text{F}_2 \left(\begin{matrix} 1 \\ \frac{3}{2} + \nu, \frac{3}{2} \end{matrix} \middle| -\frac{z^2}{4} \right)$$

We have the recurrence

$$\mathbf{H}_{\nu-1} + \mathbf{H}_{\nu+1} = \frac{2\nu}{z} \mathbf{H}_\nu + \frac{(z/2)^\nu}{\sqrt{\pi}\Gamma(\nu+3/2)}$$

((upward recursion is unstable... power-series seems quite robust enough (for $|x| < 20\dots$)))

And we have the asymptotic expansion for large z with $|\arg z| < \pi$

$$\mathbf{H}_\nu(z) - \text{Y}_\nu(z) = \frac{1}{\pi} \sum_{k=0}^{m-1} \frac{\Gamma(k+1/2)}{\Gamma(\nu-k+1/2)(z/2)^{2k-\nu+1}} + R_m$$

with $R_m = O(|z|^{\nu-2m-1})$; if ν is real, z positive, and $m - \nu + \frac{1}{2} \geq 0$, the remainder after m terms is of the same sign and numerically less than the first term neglected.

The modified Struve function is

$$\mathbf{L}_\nu(z) = -\hat{i}e^{-i\nu\pi/2} \mathbf{H}_\nu(\hat{i}z) = \left(\frac{z}{2}\right)^{\nu+1} \sum_{k=0}^{\infty} \frac{(z/2)^{2k}}{\Gamma(k+3/2)\Gamma(k+\nu+3/2)}$$

Other miscellaneous notes:

$$\mathbf{H}'_\nu = \frac{\mathbf{H}_{\nu-1} - \mathbf{H}_{\nu+1}}{2} + \frac{(z/2)^\nu}{2\sqrt{\pi}\Gamma(\nu+3/2)}$$

$$\mathbf{H}'_0 = (2/\pi) - \mathbf{H}_1$$

$$\mathbf{L}_{\nu-1} - \mathbf{L}_{\nu+1} = \frac{2\nu}{z} \mathbf{L}_\nu + \frac{(z/2)^\nu}{\sqrt{\pi}\Gamma(\nu+3/2)}$$

$$\mathbf{L}'_\nu = \frac{\mathbf{L}_{\nu-1} + \mathbf{L}_{\nu+1}}{2} + \frac{(z/2)^\nu}{\sqrt{\pi}\Gamma(\nu+3/2)}$$

[double check for typos here...]

$$\int_0^z \mathbf{H}_\nu(t) dt = \sum_{k=0}^{\infty} \frac{(-)^k (z/2)^{2k+2+\nu}}{(2k+2+\nu)\Gamma(k+3/2)\Gamma(\nu+k+3/2)} = \left(\frac{z}{2}\right)^{\nu+2} \sum_{k=0}^{\infty} \frac{(-)^k (z/2)^{2k}}{(2k+2+\nu)\Gamma(k+3/2)\Gamma(\nu+k+3/2)}$$

$$\int_0^z \mathbf{H}_0(t) dt = \frac{2}{\pi} \left(\frac{z^2}{2} - \frac{z^4}{1^2 \cdot 3^2 \cdot 4} + \frac{z^6}{1^2 \cdot 3^2 \cdot 5^2 \cdot 6} - \dots \right)$$

(just integrate term-by-term the power-series expansion)

From [ACMF]: To compute the Struve function $\mathbf{H}_n(x)$ for $n = 0, \dots, N$ (for real x), we:

- if $x = 0$, then all values are 0
- if $x < N/2$, use a power-series to compute $\mathbf{H}_{N+8}(x)$, $\mathbf{H}_{N+7}(x)$ and then use the recursion downward to $n = 0$
- if $N/2 \leq x \leq 25$, use power-series to compute $\mathbf{H}_0(x)$, $\mathbf{H}_1(x)$ and use the recursion upward to $n = N$
- else ($N/2, 25 < x$) use the asymptotic expansion to get $\mathbf{H}_0(x)$, $\mathbf{H}_1(x)$ and use the recursion upward to $n = N$

To compute the modified Struve function $\mathbf{L}_n(x)$ for $n = 0, \dots, N$ (for real x), we (for $x > 0$) use the power-series or asymptotic expansion to compute $\mathbf{L}_N(x)$, $\mathbf{L}_{N-1}(x)$, then use the backwards recursion.

From [ZJ]:

- for $\mathbf{H}_0(x)$, $\mathbf{H}_1(x)$ — use power-series for $x < 20$, asymptotic (with Y_ν) for $x > 20$
- for $\mathbf{H}_\nu(x)$, (ν arbitrary), use power-series for $x < 20$, asymptotic o.w.
- *OR* use

$$\mathbf{H}_{-n-1/2}(z) = (-)^n J_{n+1/2}(z)$$

$$\mathbf{H}_\nu(z) = \frac{1}{\Gamma(\nu+1/2)} \sum_{k=0}^{\infty} \frac{(z/2)^{k+\nu+1/2}}{k!(\nu+k+1/2)} J_{k+1/2}(z)$$

and note that the $J_{n+1/2}$ are easy to evaluate

7 Spherical harmonics, Legendre functions

spherical Legendre, toroidal Legendre, conical Legendre (Atlas [ACMF] for Computing...) `Pn`, `Pnu`, `dPn`, `dPnu`, `assocPmn`, `assocPmnu`, `Qn`, ..., `Integrate_Pn`, ... polynomials ... “spherical legendre”, “conical legendre”, “toroidal legendre”, ...

$$\begin{aligned} P_\nu(z) &= \\ Q_\nu(z) &= \frac{\pi}{2} \frac{P_\nu(z) \cos \nu z - P_\nu(-z)}{\sin \nu \pi} \\ P_\nu^m(z) &= (-)^m (1-z^2)^{m/2} \frac{d^m}{dz^m} P_\nu(z) \\ Q_\nu^m(z) &= (-)^m (1-z^2)^{m/2} \frac{d^m}{dz^m} Q_\nu(z) \end{aligned}$$

According to Lebedev, “spherical harmonics” are solutions of $(1-z^2)u'' - 2zu' + [\nu(\nu+1) - \mu^2/(1-z^2)]u = 0$ (typically $z \in (-1, 1)$ or sometimes $z \in [-\infty, 1]$; $\mu = 0, 1, 2, \dots$; $\nu \in \mathbb{R}$ or $\nu \in \mathbb{C}$).

[Lebedev] Legendre functions take $\mu = 0$ so we have $(1-z^2)u'' - 2zu' + \nu(\nu+1)u = 0$ and get Legendre functions of the first kind:

$$P_\nu(z) = F[-\nu, \nu+1; 1; \frac{1-z}{2}] \quad |z-1| < 2$$

and Legendre functions of the second kind:

$$Q_\nu(z) =$$

Both of these can be analytically continued to a larger domain. Note that $P_n(z)$ are the Legendre polynomials for $n \in \mathbb{N}_0$.

[Lebedev] The associated Legendre functions (of the first/second kinds) solve the more general equation with $\mu = m = 0, 1, 2, \dots$ where

$$\begin{aligned} P_\nu^m(z) &= (z^2-1)^{m/2} \frac{d^m}{dz^m} P_\nu(z) \\ Q_\nu^m(z) &= (z^2-1)^{m/2} \frac{d^m}{dz^m} Q_\nu(z) \end{aligned}$$

and the associated Legendre functions of the first/second kind *for the interval* $(-1, 1)$ are given by:

$$\begin{aligned} P_\nu^m(x) &= (-)^m (1-x^2)^{m/2} \frac{d^m}{dx^m} P_\nu(x) \\ Q_\nu^m(x) &= (-)^m (1-x^2)^{m/2} \frac{d^m}{dx^m} Q_\nu(x) \end{aligned}$$

8 Exponential integrals

$$\text{Ei}(z) = \int_{-\infty}^z \frac{e^t}{t} dt \quad |\arg -z| < \pi \quad (\text{correct?}) \text{ Lebedev agrees}$$

$$\text{Ei}_1(x) = \gamma + \log z + \sum_{k=1}^{\infty} \frac{x^k}{k!k} \quad x > 0$$

$$\text{E}_n(z) = \int_1^{\infty} \frac{e^{-zt}}{t^n} dt$$

Generalized exponential-integral for $p \in \mathbb{C}$:

$$\text{E}_p(z) = z^{p-1} \Gamma(1-p, z) = z^{p-1} \int_z^{\infty} \frac{e^{-t}}{t^p} dt$$

$$\text{li}(z) = \int_0^z \frac{dt}{\log t} = \text{Ei}(\ln z) \quad |\arg z| < \pi \text{ and } |\arg 1-z| < \pi$$

$\text{li}_1(z) = xxx$ modified log-integral (Lebedev)

Ei, **En**, **Ep**, **Li**, **Ci**, **Chi**, **Si**, **Shi**, **atan_int** — real / complex / adjusted real

Implementation notes for $\text{Ei}(x)$:

- For small x , use $\text{Ei}(x) = \gamma + \ln x + \sum_{m=1}^{\infty} \frac{x^m}{m!m}$ (this has some issues near the root at $0.3725074107813666\dots$ with relative precision, dropping to 14 digits using the expression $\text{Ei}(x) = \ln(x \cdot \exp(\gamma + \sum \dots))$ helps with accuracy in that area)
- For large x , use asymptotics $\text{Ei}(x) \sim \frac{e^x}{x} (1 + \sum_{m=0}^{\infty} \frac{m!}{x^m})$ (this only gives 15 digits accuracy for around $x > 38$, so use, say, for $x >= 40$)

Continued fraction for $\text{E}_1(z)$ for $|\text{ph } z| < \pi$:

$$\text{E}_1(z) = \frac{e^{-z}}{z+} \frac{1}{1+} \frac{1}{z+} \frac{2}{1+} \frac{2}{z+} \frac{3}{1+} \frac{3}{z+} \dots$$

and factorial series for $\Re z > 0$:

$$\text{E}_1(z) = e^{-z} \left(\frac{c_0}{z} + \frac{c_1}{z(z+1)} + 2! \frac{c_2}{z(z+1)(z+2)} + 3! \frac{c_3}{(z)_4} + \dots \right)$$

where $c_0 = 1$, $c_1 = -1$, $c_2 = 1/2$, $c_3 = -1/3$, $c_4 = 1/6$, and $c_k = -\sum_{j=0}^{k-1} \frac{c_j}{k-j}$ for $k \geq 1$.

Implementation notes for $\text{E}_n(x)$:

- For $n = 0$, use $\text{E}_0(x) = e^{-x}/x$
- For $n = 1$, use $\text{E}_1(x) = -\gamma - \ln x - \sum_{k=1}^{\infty} \frac{(-x)^k}{k!k}$
- For $x \leq 1$, use series $\text{E}_n(x) = \frac{(-x)^{n-1}}{(n-1)!} (-\ln x + \psi(n)) - \sum_{m=0, m \neq n-1}^{\infty} \frac{(-x)^m}{(m-(n-1))m!}$
- for $x > 1$, use continued fraction $e^{-x} \left[\frac{1}{x+n-} \frac{1 \cdot n}{x+n+2-} \frac{2 \cdot (n+1)}{x+n+4-} \dots \right]$ (see [ZJ])
- the recurrence $\text{E}_{n+1}(x) = \frac{1}{n}(e^{-x} - x \text{E}_n(x))$ ($n > 1$) is forward and backward unstable but works ok for small values (tested for $x < 1$ and $n < 10$ - should be fine for $x < 1$ and any n ; seems to work decent if $x < n$)
- asymptotic expansion $\text{E}_n(x) \sim \frac{e^{-x}}{x} (1 - \frac{n}{x} + \frac{n(n+1)}{x^2} - \frac{n(n+1)(n+2)}{x^3} + \dots)$ gives quite disappointing results - works only when $x \gg n$

9 Wave equation solutions

(From Temme ...)

Spheroidal Wave Functions - first/second kind, prolate/oblate, characteristic values, expansion coefficients, wave/angular functions, radial/angular functions, spheroidal Bessel functions, ...

10 Theta functions

10.1 Jacobian theta functions

Notation: τ is the “lattice parameter”, $q = e^{i\pi\tau}$ is the “nome”, z is the “argument” Assume that $\Im \tau > 0$ (thus $0 < |q| < 1$ and $q \notin [-\infty, 0]$).

$$\vartheta_1(z|\tau) = \vartheta_1(z, q) = 2 \sum_{n=0}^{\infty} (-)^n q^{(n+1/2)^2} \sin(2n+1)z$$

$$\vartheta_2(z|\tau) = \vartheta_2(z, q) = 2 \sum_{n=0}^{\infty} q^{(n+1/2)^2} \cos(2n+1)z$$

$$\vartheta_3(z|\tau) = \vartheta_3(z, q) = 1 + 2 \sum_{n=1}^{\infty} q^{n^2} \cos 2nz$$

$$\vartheta_4(z|\tau) = \vartheta_4(z, q) = 1 + 2 \sum_{n=1}^{\infty} (-)^n q^{n^2} \cos 2nz$$

theta_1, theta_2, theta_3, theta_4, ..., ϑ'_j/ϑ_j , Neville theta functions: $\vartheta_{s,n,d,c}$; conversions k to/from q , etc.; real/imaginary argument/complex, etc.

- For $q \in \mathbb{R}$, we can simply use the series definitions and we'll get fast convergence, though it slows down near 1
- Near $q = 1$ use the transformations (where $\tau' = -1/\tau$ and square roots assume principal values)

$$\sqrt{-i\tau}\vartheta_1(z|\tau) = -i e^{i\tau' z^2/\pi} \vartheta_1(z\tau'|\tau')$$

$$\sqrt{-i\tau}\vartheta_2(z|\tau) = e^{i\tau' z^2/\pi} \vartheta_4(z\tau'|\tau')$$

$$\sqrt{-i\tau}\vartheta_3(z|\tau) = e^{i\tau' z^2/\pi} \vartheta_3(z\tau'|\tau')$$

$$\sqrt{-i\tau}\vartheta_4(z|\tau) = e^{i\tau' z^2/\pi} \vartheta_2(z\tau'|\tau')$$

- For example, letting $\varphi = -\ln q/\pi$ and $q' = e^{-\pi/\varphi}$, we end up with the very rapidly converging (near $q = 1$):

$$\vartheta_3(z, q) = \frac{e^{-z^2/\pi\varphi}}{\sqrt{\varphi}} \left[1 + 2 \sum_{n=1}^{\infty} (q')^{n^2} \cosh 2nz/\varphi \right]$$

(actually, in practice it seems to work generally better (for $z = 0.3$) than the original sum with just a couple more terms needed near $q = 0...$ oops with $z \sim \pi/2$ we have some issues with the transformed version dying with NaN for $q \sim 1$ (though giving much better results until then than the original series — just need to deal with overflow/underflow issues with exp and cosh... yes, writing $(q')^{n^2} \cosh(2nz/\varphi) = e^{-n^2\pi/\varphi + 2nz/\varphi} (1 + e^{-4nz/\varphi})/2$ gives much better results (though seem to lose a bit of precision...))

11 Elliptic integrals

Weierstrass elliptic function \mathbf{p} , inverse weierstrass, elliptic zeta function, etc. Fock functions, etc.

General notation notes: the “nome” q and the “modulus” k can be related as follows, where $k' = \sqrt{1 - k^2}$:

$$\begin{aligned} q &= e^{-\pi K'(k)/K(k)} \\ k &= \vartheta_2^2(0, q)/\vartheta_3^2(0, q) \\ k' &= \vartheta_4^2(0, 1)/\vartheta_3^2(0, q) \\ K(k) &= \frac{\pi}{2} \vartheta_3^2(0, q) \end{aligned}$$

(conversions)

11.1 Elliptic integrals

First kind: $(x = \sin \varphi)$, $0 \leq k \leq 1$

$$F(\varphi, k) = \int_0^\varphi \frac{d\vartheta}{\sqrt{1 - k^2 \sin^2 \vartheta}} = \int_0^x \frac{dt}{\sqrt{(1 - t^2)(1 - k^2 t^2)}}$$

Second kind: $(x = \sin \varphi)$, $0 \leq k \leq 1$

$$E(\varphi, k) = \int_0^\varphi \sqrt{1 - k^2 \sin^2 \vartheta} d\vartheta = \int_0^x \frac{\sqrt{1 - k^2 t^2}}{\sqrt{1 - t^2}} dt$$

Third kind: $(x = \sin \varphi)$, $0 \leq k \leq 1$

$$\Pi(\varphi, k, c) = \int_0^\varphi \frac{d\vartheta}{(1 - c \sin^2 \vartheta) \sqrt{1 - k^2 \sin^2 \vartheta}} = \int_0^x \frac{dt}{(1 - ct^2) \sqrt{(1 - t^2)(1 - k^2 t^2)}}$$

Special case:

$$\Pi(\varphi, 0, k) = \int_0^\varphi \frac{d\vartheta}{1 - \xi \sin^2 \vartheta} = \frac{\text{atan}(\sqrt{1 - \xi} \tan(\varphi))}{\sqrt{1 - \xi}}$$

To implement $F(\varphi, k)$ and $E(\varphi, k)$, use the ascending Landen transforms.
To implement $\Pi(k, c, \varphi)$, the Gauss transform is effective.

11.2 Complete elliptic integrals

$$K(k) = F(\pi/2, k)$$

$$E(k) = E(\pi/2, k)$$

$$\Pi(k, c) = \Pi(\pi/2, k, c)$$

We use AGM approaches to compute these

To implement $\Pi(k, c)$, we use the AGM approach: for $-\infty < k^2 < 1$, $-\infty < c < 1$, we have

$$\Pi(c, k) = \frac{\pi}{4M(1, k')} \left(2 + \frac{c}{1 - c} \sum_{n=0}^{\infty} Q_n \right)$$

where $a_0 = 1$, $b_0 = k'$, $p_0^2 = 1 - c$, $Q_0 = 1$, and

$$\begin{aligned} p_{n+1} &= \frac{p_n^2 + a_n b_n}{2p_n} \\ Q_{n+1} &= \frac{1}{2} Q_n \varepsilon_n \\ \varepsilon_n &= \frac{p_n^2 - a_n b_n}{p_n^2 + a_n b_n} \end{aligned}$$

Also $Z, FK, CE, \Lambda_0, RC, RD, RF, RJ \dots$

11.3 Complementary complete elliptic integrals

Let $k' = \sqrt{1 - k^2}$. Then

$$K'(k) = K(k')$$

$$E'(k) = E(k')$$

11.4 Carlson's symmetric elliptic integrals

$$R_C(x, y) = \frac{1}{2} \int_0^\infty \frac{dt}{\sqrt{t+x}(t+y)} \quad x \in \mathbb{C} \setminus (-\infty, 0), y \in \mathbb{C} \setminus \{0\}$$

To compute $R_C(x, y)$ for $x, y \in \mathbb{R}$, we simply use the closed-form expressions

- $R_C(x, x) = x^{-1/2}$

- For $0 \leq x < y$:

$$R_C(x, y) = \frac{1}{\sqrt{y-x}} \operatorname{atan} \sqrt{\frac{y-x}{x}} = \frac{1}{\sqrt{y-x}} \operatorname{acos} \sqrt{\frac{x}{y}}$$

- For $0 < y < x$: (note there is a typo in DLMF for this formula)

$$R_C(x, y) = \frac{1}{\sqrt{x-y}} \operatorname{atanh} \sqrt{\frac{x-y}{x}} = \frac{1}{\sqrt{x-y}} \ln \frac{\sqrt{x} + \sqrt{x-y}}{\sqrt{y}}$$

- For $y < 0 \leq x$:

$$R_C(x, y) = \sqrt{\frac{x}{x-y}} R_C(x-y, -y) = \frac{1}{\sqrt{x-y}} \operatorname{atanh} \sqrt{\frac{x}{x-y}} = \frac{1}{\sqrt{x-y}} \ln \frac{\sqrt{x} + \sqrt{x-y}}{\sqrt{-y}}$$

11.5 Jacobian elliptic functions

We have implicit parameter k in the following

$$\operatorname{sn}(u) = x \quad \text{s.t.} \quad u = F(k, x) = \int_0^x \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}}$$

$$\operatorname{cn}(u) = \sqrt{1 - \operatorname{sn}^2 u}$$

$$\operatorname{dn}(u) = \sqrt{1 - k^2 \operatorname{sn}^2 u}$$

I.e. $\operatorname{sn}^2 + \operatorname{cn}^2 = 1$ and $\operatorname{dn}^2 + k^2 \operatorname{sn}^2 = 1$. other inverse?

From [DLMF];

- Three basic techniques to compute: use ϑ functions, use AGM, use Landen transformations and then series in k

- Theta functions: let $\zeta = \frac{\pi z}{2K(k)}$

$$\begin{aligned}\operatorname{sn}(z, k) &= \frac{\vartheta_3(0, q)}{\vartheta_2(0, q)} \frac{\vartheta_1(\zeta, q)}{\vartheta_4(\zeta, q)} = \frac{1}{ns} \\ \operatorname{cn}(z, k) &= \frac{\vartheta_4(0, q)}{\vartheta_2(0, q)} \frac{\vartheta_2(\zeta, q)}{\vartheta_4(\zeta, q)} = \frac{1}{nc} \\ \operatorname{dn}(z, k) &= \frac{\vartheta_4(0, q)}{\vartheta_3(0, q)} \frac{\vartheta_3(\zeta, q)}{\vartheta_4(\zeta, q)} = \frac{1}{nd} \\ \operatorname{sd}(z, k) &= \frac{\vartheta_3(0, q)^2}{\vartheta_2(0, q)\vartheta_4(0, q)} \frac{\vartheta_1(\zeta, q)}{\vartheta_3(\zeta, q)} = \frac{1}{ds} \\ \operatorname{cd}(z, k) &= \frac{\vartheta_3(0, q)}{\vartheta_2(0, q)} \frac{\vartheta_1(\zeta, q)}{\vartheta_2(\zeta, q)} = \frac{1}{dc} \\ \operatorname{sc}(z, k) &= \frac{\vartheta_3(0, q)}{\vartheta_4(0, q)} \frac{\vartheta_1(\zeta, q)}{\vartheta_2(\zeta, q)} = \frac{1}{cs}\end{aligned}$$

- AGM:

1. Let $a_0 = 1$, $b_0 = k' \in (0, 1)$
2. Compute a_N, b_N, c_N via AGM
3. Let $\varphi_N = 2^N a_N x$
4. and $\varphi_{n-1} = \frac{1}{2}(\varphi_n + \operatorname{asin}(\frac{c_n}{a_n} \sin \varphi_n))$
5. Then $\operatorname{sn}(x, k) = \sin \varphi_0$, $\operatorname{cn}(x, k) = \cos \varphi_0$, and $\operatorname{dn}(x, k) = \cos \varphi_0 / \cos(\varphi_1 - \varphi_0)$

12 Parabolic cylinder functions

$$\begin{aligned}D_\nu(z) &= U(a, z) = & \nu &= -a - \frac{1}{2} \\ V_\nu(z) &= V(a, z) = & \nu &= -a - \frac{1}{2} \\ W(a, \pm z) &= \end{aligned}$$

Dnu, Vnu, Wa_p/m, dDnu, ..., Ua, Va, ...

12.1 Lebedev

(According to Lebedev [Lebedev], solutions to $u'' + (2\nu + 1 - z^2)u = 0$ are the *parabolic cylinder functions*.)

According to Lebedev: solutions to $v'' - 2zv' + 2\nu v = 0$ are *Hermite functions*

$$H_\nu(z) = \frac{2^\nu \Gamma(1/2)}{\Gamma((1-\nu)/2)} \Phi[-\nu/2, 1/2; z^2] + \frac{2^\nu \Gamma(-1/2)}{\Gamma(-\nu/2)} \Phi[(1-\nu)/2, 3/2; z^2]$$

And if $\nu = n = 0, 1, 2, \dots$ then we get the Hermite Polynomials

12.2 NIST handbook

According to NIST handbook [NIST]: Parabolic cylinder functions are solutions of the differential equation $w'' + (az^2 + bz + c)w = 0$ with three distinct standard forms:

1. $w'' - (z^2/4 + a)w = 0$ with solutions $U(a, \pm z)$, $V(a, \pm z)$, $\bar{U}(a, \pm z)$ (not complex conjugate), and $U(-a, \pm iz)$; for real values of $z (= x)$, numerically satisfactory pairs of solutions are $U(a, x)$ and $V(a, x)$ for $x > 0$ and $U(a, -x)$, $V(a, -x)$ for $x < 0$.

2. $w'' + (z^2/4 - a)w = 0$ with solutions $W(a, \pm x)$; for all real x , a numerically satisfactory pair is $W(a, x)$, $W(a, -x)$.
3. and $w'' + (\nu + 1/2 - z^2/4)w = 0$ with solutions $D_\nu(\pm z)$ where $D_\nu(z) = U(-1/2 - \nu, z)$ (Whittaker function);

In \mathbb{C} , for $j = 0, 1, 2, 3$, a numerically satisfactory pair of solutions is given by $U((-)^{j-1}a, (-i)^{j-1}z)$ and $U(-)^ja, (-i)^jz$ in the half-plane $\frac{1}{4}(2j-3)\pi \leq \text{ph } z \leq \frac{1}{4}(2j+1)\pi$.

We can express U and V in terms of D via

$$\begin{aligned} U(a, x) &= D_{-a-1/2}(x) \\ V(a, x) &= \frac{\Gamma(a + \frac{1}{2})}{\pi} (\sin(\pi a) D_{-a-1/2}(x) + D_{-a-1/2}(-x)) \end{aligned}$$

All solutions are entire functions of z and a or ν .

Values at $z = 0$ are given by

$$\begin{aligned} U(a, 0) &= \frac{\sqrt{\pi}}{2^{a/2+1/4}\Gamma(\frac{3}{4} + \frac{1}{2}a)} \\ U'(a, 0) &= -\frac{\sqrt{\pi}}{2^{a/2-1/4}\Gamma(\frac{1}{4} + \frac{1}{2}a)} \\ V(a, 0) &= \frac{\pi 2^{a/2+1/4}}{\Gamma^2(\frac{3}{4} - \frac{1}{2}a)\Gamma(\frac{1}{4} + \frac{1}{2}a)} = \frac{2^{1/4+a/2} \sin(\pi(\frac{3}{4} - \frac{1}{2}a))}{\Gamma(\frac{3}{4} - \frac{1}{2}a)} \\ V'(a, 0) &= \frac{\pi 2^{a/3+3/4}}{\Gamma^2(\frac{1}{4} - \frac{1}{2}a)\Gamma(\frac{3}{4} + \frac{1}{2}a)} = \frac{2^{3/4+a/2} \sin(\pi(\frac{1}{4} - \frac{1}{2}a))}{\Gamma(\frac{1}{4} - \frac{1}{2}a)} \\ W(a, 0) &= 2^{-3/4} \left| \frac{\Gamma(\frac{1}{4} + \frac{1}{2}a\hat{i})}{\Gamma(\frac{3}{4} + \frac{1}{2}a\hat{i})} \right|^{1/2} \\ W'(a, 0) &= -2^{-1/4} \left| \frac{\Gamma(\frac{3}{4} + \frac{1}{2}a\hat{i})}{\Gamma(\frac{1}{4} + \frac{1}{2}a\hat{i})} \right|^{1/2} \quad ?\text{sign?} \end{aligned}$$

`initial_value_U, ...`

Series expansions converging for all z are given by

$$\begin{aligned} U(a, z) &= U(a, 0)u_1(a, z) + U'(a, 0)u_2(a, z) \\ V(a, z) &= V(a, 0)u_1(a, z) + V'(a, 0)u_2(a, z) \\ W(a, z) &= W(a, 0)w_1(a, z) + W'(a, 0)w_2(a, z) \end{aligned}$$

where

$$\begin{aligned} u_1(a, z) &= e^{-z^2/4} M(\frac{1}{4} + \frac{1}{2}a, \frac{1}{2}, \frac{z^2}{2}) = e^{-z^2/4} \left(1 + (a + \frac{1}{2}) \frac{z^2}{2!} + (a + \frac{1}{2})(a + \frac{5}{2}) \frac{z^4}{4!} + \dots \right) \\ &= e^{z^2/4} M(\frac{1}{4} - \frac{1}{2}a, \frac{1}{2}, \frac{-z^2}{2}) = e^{z^2/4} \left(1 + (a - \frac{1}{2}) \frac{z^2}{2!} + (a - \frac{1}{2})(a - \frac{5}{2}) \frac{z^4}{4!} + \dots \right) \\ u_2(a, z) &= ze^{-z^2/4} M(\frac{3}{4} + \frac{1}{2}a, \frac{3}{2}, \frac{z^2}{2}) = e^{-z^2/4} \left(z + (a + \frac{3}{2}) \frac{z^3}{3!} + (a + \frac{3}{2})(a + \frac{7}{2}) \frac{z^5}{5!} + \dots \right) \\ &= ze^{z^2/4} M(\frac{3}{4} - \frac{1}{2}a, \frac{3}{2}, \frac{-z^2}{2}) = e^{z^2/4} \left(z + (a - \frac{3}{2}) \frac{z^3}{3!} + (a - \frac{3}{2})(a - \frac{7}{2}) \frac{z^5}{5!} + \dots \right) \end{aligned}$$

or we have the series

$$\begin{aligned} u_1(a, z) &= 1 + a \frac{z^2}{2!} + (a^2 + \frac{1}{2}) \frac{z^4}{4!} + (a^3 + \frac{7}{2}a) \frac{z^6}{6!} + \dots \\ u_2(a, z) &= z + a \frac{z^3}{3!} + (a^2 + \frac{3}{2}) \frac{z^5}{5!} + (a^3 + \frac{13}{2}a) \frac{z^7}{7!} + \dots \end{aligned}$$

where the coefficients of $x^n/n!$ are given by $a_{n+2} = aa_n + \frac{1}{4}n(n-1)a_{n-2}$.

Also,

$$\begin{aligned} w_1(a, z) &= \sum_{n=0}^{\infty} \alpha_n(a) \frac{x^{2n}}{(2n)!} \\ w_2(a, z) &= \sum_{n=0}^{\infty} \beta_n(a) \frac{x^{2n+1}}{(2n+1)!} \end{aligned}$$

with $\alpha_0(a) = \beta_0(a) = 1$, $\alpha_1(a) = \beta_1(a) = a$, and

$$\begin{aligned} \alpha_{n+2} &= a\alpha_{n+1} - \frac{1}{2}(n+1)(2n+1)\alpha_n \\ \beta_{n+2} &= a\beta_{n+1} - \frac{1}{2}(n+1)(2n+3)\beta_n \end{aligned}$$

(Can get recursions for the full series terms directly: $\hat{\alpha}_n = \alpha_n/(2n)!$ then $\hat{\alpha}_{n+2} = \frac{a\hat{\alpha}_{n+1} - \frac{1}{4}\hat{\alpha}_n}{(2n+3)(2n+4)}$, etc.

Recurrence relations are

$$zU(a, z) - U(a-1, z) + (a + \frac{1}{2})U(a+1, z) = 0$$

$$zV(a, z) - V(a+1, z) + (a - \frac{1}{2})V(a-1, z) = 0$$

Asymptotic expansions for $z \rightarrow \infty$ (δ is an arbitrarily small positive constant)

$$\begin{aligned} U(a, z) &\sim e^{-z^2/4} z^{-a-1/2} \sum_{s=0}^{\infty} (-)^s \frac{(\frac{1}{2} + a)_{2s}}{s!(2z^2)^s} \quad |\text{ph } z| \leq \frac{3}{4}\pi - \delta \\ U(a, z) &\sim e^{-z^2/4} z^{-a-1/2} \sum_{s=0}^{\infty} (-)^s \frac{(\frac{1}{2} + a)_{2s}}{s!(2z^2)^s} \pm i \frac{\sqrt{2\pi}}{\Gamma(\frac{1}{2} + a)} e^{\mp i\pi a} e^{z^2/4} z^{a-1/2} \sum_{s=0}^{\infty} \frac{(\frac{1}{2} - a)_{2s}}{s!(2z^2)^s} \\ &\quad \frac{1}{4}\pi + \delta \leq \pm \text{ph } z \leq \frac{5}{4}\pi - \delta \\ V(a, z) &\sim \sqrt{\frac{2}{\pi}} e^{z^2/4} z^{a-1/2} \sum_{s=0}^{\infty} \frac{(\frac{1}{2} - a)_{2s}}{s!(2z^2)^s} \quad |\text{ph } z| \leq \frac{1}{4}\pi - \delta \\ V(a, z) &\sim \sqrt{\frac{2}{\pi}} e^{z^2/4} z^{a-1/2} \sum_{s=0}^{\infty} \frac{(\frac{1}{2} - a)_{2s}}{s!(2z^2)^s} \pm i \frac{1}{\Gamma(\frac{1}{2} - a)} e^{-z^2/4} z^{-a-1/2} \sum_{s=0}^{\infty} (-)^s \frac{(\frac{1}{2} + a)_{2s}}{s!(2z^2)^s} \\ &\quad -\frac{1}{4}\pi + \delta \leq \pm \text{ph } z \leq \frac{3}{4}\pi - \delta \end{aligned}$$

Initial testing seems to indicate that, at least for small a , the series expansions can give decent results, though not necessarily a full 15 digits (with $a = -20$, looks like only 13 digits, even for small z ; with $a = -2$, getting around 15 digits).

13 Lambert functions

$$W(z) = \rho \quad \text{s.t.} \quad \rho e^{\rho} = z$$

$$W_{[k]}(z) = k\text{'th branch}$$

14 Hypergeometric, Meijer G functions

\mathcal{G} -function, \mathcal{H} -function, Macdonald E function, Carr $\varphi[\alpha, \beta\gamma|x, y]$ function,

0f0, 1f0, 0f1, 1f1, 1f2, 2f1, 2f2, 3f2, 3f3, 2f3, pfq, ... Confluent M and U (“first” and “second” kinds) [have alternative ode solutions for all hypergeom funcs?]

Utility functions for hypergeometric-type *series*(?): odd/even parts of series, alternating odd/even parts, mixed odd/even ($x^n y^{2n}$) etc.?

“Gamma series” variations (and modifications as above): (?)

$$\sum_{n=0}^{\infty} \frac{\Gamma(a_1 + n) \cdots \Gamma(a_p + n)}{\Gamma(b_1 + n) \cdots \Gamma(b_q + n)} x^n$$

For example,

$$\sum_{n=0}^{\infty} \frac{x^n}{\Gamma(\alpha + n)} = \frac{1}{\Gamma(\alpha)} \sum_{n=0}^{\infty} \frac{x^n}{(\alpha)_n} = \frac{1}{\Gamma(\alpha)} {}_1F_1 \left(\begin{matrix} 1 \\ \alpha \end{matrix} \middle| x \right)$$

Hybrid series, e.g. $\sum_{n=0}^{\infty} \frac{(a)_n}{\Gamma(b+n)} \frac{x^n}{n!}$ etc. (works for $b = 0, -1, -2, \dots$ also!)

Hypergeometric *functions* vs. hypergeometric *series*

Scaled versions...

Define

$${}_2F_1 \left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} \middle| z \right) = \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n} \frac{z^n}{n!} = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\beta)} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha + n)\Gamma(\beta + n)}{\Gamma(\gamma + n)} \frac{x^n}{n!}$$

This series is defined for $|z| < 1$ and $\gamma \neq 0, -1, -2, \dots$. It can be extended to an analytic function on $\mathbb{C} \setminus [1, \infty]$ via the integral

$${}_2F_1 \left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix} \middle| z \right) = \frac{\Gamma(\gamma)}{\Gamma(\beta)\Gamma(\gamma - \beta)} \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-tz)^\alpha dt$$

for $\Re \gamma > \Re \beta > 0$ and $|\arg 1 - z| < \pi$.

The hypergeometric polynomials are given by $F(-n, \beta; \gamma; z)$.

Note that Lebedev defines the hypergeometric function of the second kind as $G(\alpha, \beta; \gamma; z) = \dots$ which is the other solution of the hypergeometric equation $z(1-z)u'' + [\gamma - (\alpha + \beta + 1)z]u' - \alpha\beta u = 0$ (for $|z| < 1$.)

The generalized hypergeometric function is:

$${}_pF_q \left(\begin{matrix} \vec{\alpha} \\ \vec{\gamma} \end{matrix} \middle| z \right) = \sum_{n=0}^{\infty} \frac{(\alpha_1)_n \cdots (\alpha_p)_n}{(\gamma_1)_n \cdots (\gamma_q)_n} \frac{z^n}{n!}$$

$$M(\alpha, \beta; z) = {}_1F_1 \left(\begin{matrix} \alpha \\ \beta \end{matrix} \middle| z \right) = \sum_{n=0}^{\infty} \frac{(\alpha)_n}{(\beta)_n} \frac{z^n}{n!}$$

For $\beta \neq 0, -1, -2, \dots$, $|z| < \infty$ (defines an entire function of z on \mathbb{C}). (In Lebedev's notation we have $\Phi(\alpha, \beta; z)$.) (Lebedev defines the confluent hypergeometric function of the second kind as $\Psi(\alpha, \gamma; z) = \dots$)

$$U(\alpha, \beta; z) = \frac{\pi}{\sin \pi \beta} \left(\frac{M(\alpha, \beta; z)}{\Gamma(\alpha - \beta + 1)\Gamma(\beta)} - z^{1-\beta} \frac{M(\alpha - \beta + 1, 2 - \beta; z)}{\Gamma(\alpha)\Gamma(2 - \beta)} \right)$$

14.1 Whittaker functions

$$M_{k,\mu}(z) = z^{\mu+1/2} e^{-z/2} M(\mu + \frac{1}{2} - k, 2\mu + 1; z)$$

$$W_{k,\mu}(z) = \frac{\Gamma(-2\mu)}{\Gamma(1/2 - \mu - k)} M_{k,\mu}(z) + \frac{\Gamma(2\mu)}{\Gamma(1/2 + \mu - k)} M_{k,-\mu}(z)$$

15 Zeta functions

Need *zeros* also

Hurwitz Zeta

Lerch Phi

15.1 Riemann zeta function

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} \quad \Re s > 1$$

$$\eta(s) = \sum_{n=1}^{\infty} (-1)^{n-1} n^{-s} = (1 - 2^{1-s})\zeta(s) \quad \Re s > 1$$

$$\lambda(s) = \sum_{n=1}^{\infty} (2n-1)^{-s} = (1 - 2^{-s})\zeta(s) \quad \Re s > 1$$

$$\beta(s) = \sum_{n=1}^{\infty} (-1)^{n-1} (2n-1)^{-s} \quad \Re s > 1$$

We should have implementation of $\zeta(s) - 1$ for large s , (especially for $s \in \mathbb{N}$). (and for other such funcs) [also $\zeta(s) - 2^{-s}$, etc.??] Also, a version of just $\text{zeta}(\frac{1}{2} + it)$ for real t ...

Reflection:

$$\zeta(1-s) = \frac{2}{(2\pi)^s} \cos(s\pi/2) \Gamma(s) \zeta(s)$$

$$\zeta(s) = 2(2\pi)^{s-1} \sin(s\pi/2) \Gamma(1-s) \zeta(1-s)$$

we can write this as $\xi(s) = \xi(1-s)$ where we define

$$\xi(s) = \frac{s(s-1)}{2} \Gamma(s/2) \pi^{-s/2} \zeta(s)$$

$$\zeta(-n) = -\frac{B_{n+1}}{n+1}$$

(A) For all $s \in \mathbb{C} \setminus \{1\}$,

$$\zeta(s) = \frac{1}{2} + \frac{1}{s-1} + 2 \int_0^{\infty} \frac{\sin(s \operatorname{atan} t)}{(1+t^2)^{s/2} (e^{2\pi t} - 1)} dt$$

(B) For $s \in \mathbb{C}$,

$$\pi^{-s/2} \Gamma(s/2) \zeta(s) = -\frac{1}{s} - \frac{1}{1-s} + \frac{1}{2} \int_1^{\infty} (t^{(1-s)/2} + t^{s/2}) (\vartheta_3(e^{-\pi t}) - 1) \frac{dt}{t}$$

(B) Following from **(B)**, we get

$$\zeta(s) \Gamma(s/2) = \frac{\pi^{s/2}}{s(s-1)} + \sum_{n=1}^{\infty} n^{-s} \Gamma(\frac{s}{2}, \pi n^2) + \pi^{s-1/2} \sum_{n=1}^{\infty} n^{s-1} \Gamma(\frac{1-s}{2}, \pi n^2)$$

[DLMF] claims that a principal approach is using (Euler-Maclaurin)

$$\zeta(s) = \sum_{k=1}^N \frac{1}{k^s} + \frac{N^{1-s}}{s-1} - \frac{N^{-s}}{2} + \sum_{k=1}^n \binom{s+2k-2}{2k-1} \frac{B_{2k}}{2k} N^{1-s-2k} - \binom{s+2n}{2n+1} \int_N^\infty \frac{\tilde{B}_{2n+1}(x)}{x^{s+2n+1}} dx$$

for $\Re s > -2n; n, N = 1, 2, 3, \dots$ (A quick test shows that just the first 2 terms after the initial sum improve the accuracy significantly...) [INCLUDE EXPLICIT FIRST FEW TERMS...]

Notes on using the Euler-Maclaurin expansion above:

- stopping criterion is a little bit trickier (but in practice not a bit problem)
- improves convergence speed significantly, especially for small $s > 1$, chart below gives some sample data (running with long doubles, indicating number of extra E-M terms): gives terms needed to convergence and achieved digits of accuracy

s	raw series	2-terms + n=0	2-terms + n=1	2-terms + n=2	2-terms + n=3	2-terms + n=4	2-terms + n=5
2	> 9999/3.8	> 9999/12.6	1089/16.5	201/17.6	77/18.1	40/18.6	26/18.3
5	> 9999/15.2	910/16.7	204/17.5	78/17.7	48/18.6	33/18.6	24/18.7
10	170/18	83/19.1	40/18.7	34/18.5	25/19	22/18.5	23/18.5
40	8/full	8/full	8/full	8/full	8/full	9/full	8/full

Notes on the Dirichlet Beta function $\beta(\nu)$:

- $\beta(-n) = \frac{1}{2} E_n$ for $n = 0, 1, 2, \dots$
- $\beta(n) = \left(\frac{\pi}{2}\right)^n \frac{|E_{n-1}|}{2(n-1)!}$ for $n = 1, 3, 5, \dots$
- $\beta(\nu) = \frac{1}{2\Gamma(\nu)} \int_0^\infty t^{\nu-1} \operatorname{sech}(t) dt \quad (\nu > 0)$
- $\beta(1-\nu) = \frac{\Gamma(\nu) \sin(\nu\pi/2)}{(\pi/2)^\nu} \beta(\nu) \quad \beta \neq 0, -1, -2, \dots$
- $\beta(\nu) = 4^{-\nu} (\zeta(\nu, 1/4) - \zeta(\nu, 3/4))$ (Hurwitz zeta function)
- $\beta(\nu) = 2^{-\nu} \eta(\nu, 1/2)$ (bivariate eta function - from [Atlas])
- It is claimed in [Atlas] that this is the Euler-Maclaurin expansion, but it works very poorly:

$$\beta(\nu) = \lim_{J \rightarrow \infty} \left\{ \sum_{j=1}^J \frac{(-)^{j+1}}{(2j-1)^\nu} + \sum_{k=1}^K \frac{2^{k-1}(2^k-1)(\nu)_{k-1} B_k}{k!(2J-1)^{k+\nu-1}} \right\}$$

- A much more effective Euler-Maclaurin expansion is given by

$$\begin{aligned} \beta(\nu) &= \sum_{k=0}^{2N} \frac{(-)^k}{(2k+1)^\nu} + \frac{(4N+1)^{1-\nu} - (4N+3)^{1-\nu}}{4(\nu-1)} - \frac{(4N+1)^{-\nu} - (4N+3)^{-\nu}}{2} \\ &\quad - \sum_{j=2}^K (-4)^{j-1} (\nu)_{j-1} \frac{B_j}{j!} [(4N+1)^{-\nu-j+1} - (4N+3)^{-\nu-j+1}] \end{aligned}$$

15.2 Hurwitz' zeta function

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s} \quad \Re s > 1, a \neq 0, -1, -2, \dots$$

(with continuation in s to \mathbb{C} with only a simple pole at $s = 1$).

We have the following simple identities

$$\begin{aligned} \zeta(s, a) &= \zeta(s, a+1) + a^{-s} \\ \zeta(s, a) &= \zeta(s, a+m) + \sum_{n=0}^{m-1} (n+a)^{-s} \\ \frac{\partial}{\partial a} \zeta(s, a) &= -s \zeta(s+1, a) \end{aligned}$$

We have a series expansion for $s \neq 1, |a-1| < 1$:

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{\Gamma(n+s)}{\Gamma(s)} \frac{(1-a)^n}{n!} \zeta(n+s)$$

As $a \rightarrow 0$ with $s \neq 1$ fixed

$$\zeta(s, a+1) = \zeta(s) - s \zeta(s+1)a + O(a^2)$$

We have the following Euler-Maclaurin-type expansions

$$\begin{aligned} \zeta(s, a) &= \sum_{n=0}^N (n+a)^{-s} + \frac{(N+a)^{1-s}}{s-1} - s \int_N^{\infty} \dots \\ \zeta(s, a) &= a^{-s} \left(\frac{1}{2} + \frac{a}{s-1} \right) - s(s+1) \int_0^{\infty} \dots \\ \zeta(s, a) &= a^{-s} + (1+a)^{-s} \left(\frac{1}{2} + \frac{1+a}{s-1} \right) + \sum_{k=1}^{\infty} \binom{s+2k-2}{2k-1} \frac{B_{2k}}{2k} \frac{1}{(1+a)^{s+2k-1}} - R \dots \end{aligned}$$

15.3 Lerch's transcendent

$$\Phi(z, s, a) = \sum_{n=1}^{\infty} \frac{z^n}{(a+n)^s} \quad a \neq 0, -1, -2, \dots; |z| < 1; (|z| = 1 \implies \Re(s) > 1)$$

(To implement, try out Euler-Maclaurin...)

We have the following identities

$$\begin{aligned} \Phi(z, s, a) &= z^m \Phi(z, s, a+m) + \sum_{n=0}^{m-1} \frac{z^n}{(a+n)^s} \\ \Phi(z, s, a) &= \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{x^{s-1} e^{-ax}}{1 - ze^{-x}} dx \end{aligned}$$

(for $\Re s > 0, \Re a > 0, z \in \mathbb{C} \setminus [1, \infty)$.)

15.4 Dirichlet L-function

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} \quad \Re(s) > 1, \chi \text{ a Dirichlet character (mod } k)$$

(express as sum of zeta-functions, allow a general k -periodic character χ .)

The following is valid for all s if $\chi \neq \chi_1$; otherwise valid for all $s \neq 1$:

$$L(s, \chi) = k^{-s} \sum_{r=1}^{k-1} \chi(r) \zeta(s, \frac{r}{k})$$

15.5 Polylogarithms and related

15.5.1 Dilogarithm

The dilogarithm is defined via

$$\mathrm{Li}_2(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^2} \quad |z| \leq 1$$

and can be analytically continued to the complex plane split along the real line from +1 to ∞ as

$$\mathrm{Li}_2(z) = - \int_0^z \ln(1-t) \frac{dt}{t} \quad z \in \mathbb{C} \setminus (1, \infty)$$

(Compare also with Spence integral) `Dilog`, `Li_2`, `Polylog`, `Li_n`, ... generalized polylog $\mathrm{Li}_s(z)$...
Noodling around:

$$\begin{aligned} \mathrm{Li}_2(z) &= \sum_{n=1}^{\infty} \frac{z^n}{n^2} \\ \frac{\mathrm{Li}_2(z^2)}{4} &= \sum_{n=1}^{\infty} \frac{z^{2n}}{(2n)^2} \\ \mathrm{Li}_2(z) - \frac{\mathrm{Li}_2(z^2)}{4} &= \sum_{n=1}^{\infty} \frac{z^{2n+1}}{(2n+1)^2} \\ \frac{1}{i} \left(\mathrm{Li}_2(iz) - \frac{\mathrm{Li}_2(-z^2)}{4} \right) &= \sum_{n=1}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)^2} \end{aligned}$$

We can derive the following Euler-Maclaurin expansion for the power-series to get

$$\mathrm{Li}_2(z) = \sum_{k=1}^N \frac{z^k}{k^2} + \frac{\mathrm{E}_2(-N \ln z)}{N} - \frac{z^N}{2N^2} - \sum_{j=2}^K \frac{B_j}{j!} f^{(j-1)}(N) + R_K$$

where $\mathrm{E}_2(z)$ is the exponential integral, and $f(x) = z^x/x^2$. This improves convergence, but the cost of computing the exponential integral outweighs any benefits. Note that we have

$$\begin{aligned} f'(n) &= \frac{z^n}{n^2} \left(\ln z - \frac{2}{n} \right) \\ f^{(2)}(n) &= \frac{z^n}{n^2} \left((\ln z)^2 - 4 \frac{\ln z}{n} + \frac{6}{n^2} \right) \\ f^{(3)}(n) &= \frac{z^n}{n^2} \left((\ln z)^3 - 6 \frac{(\ln z)^2}{n} + 18 \frac{\ln z}{n^2} - \frac{24}{n^3} \right) \\ f^{(4)}(n) &= \frac{z^n}{n^2} \left((\ln z)^4 - 8 \frac{(\ln z)^3}{n} + 36 \frac{(\ln z)^2}{n^2} - 96 \frac{\ln z}{n^3} + \frac{120}{n^4} \right) \\ f^{(5)}(n) &= \frac{z^n}{n^2} \left((\ln z)^5 - 10 \frac{(\ln z)^4}{n} + 60 \frac{(\ln z)^3}{n^2} - 240 \frac{(\ln z)^2}{n^3} + 600 \frac{\ln z}{n^4} - \frac{720}{n^5} \right) \end{aligned}$$

Many identities are useful:

$$\mathrm{Li}_2(z) + \mathrm{Li}_2\left(\frac{z}{z-1}\right) = -\frac{1}{2}(\ln 1-z)^2 \quad z \in \mathbb{C} \setminus [1, \infty)$$

$$\mathrm{Li}_2(z) + \mathrm{Li}_2\left(\frac{1}{z}\right) = -\frac{\pi^2}{6} - \frac{1}{2}(\ln -z)^2 \quad z \in \mathbb{C} \setminus [0, \infty)$$

$$\operatorname{Li}_2(x) + \operatorname{Li}_2(1-x) = \frac{\pi^2}{6} - (\ln x)(\ln 1-x) \quad 0 < x < 1$$

$$\operatorname{Li}_2(z^m) = m \sum_{k=0}^{m-1} \operatorname{Li}_2(z e^{2\pi i k/m}) \quad |z| < 1, m = 1, 2, 3, \dots$$

When $z = e^{i\vartheta}$, the original definition becomes

$$\operatorname{Li}_2(e^{i\vartheta}) = \sum_{n=1}^{\infty} \frac{\cos n\vartheta}{n^2} + i \sum_{n=1}^{\infty} \frac{\sin n\vartheta}{n^2} = \left(\frac{\pi^2}{6} - \frac{\pi\vartheta}{2} + \frac{\vartheta^2}{4} \right) - i \int_0^{\vartheta} \ln(2 \sin(x/2)) dx$$

(and notice the final integral is Clausen's integral).

`dilog(z) = Li2(z)`

```
## -*- texinfo -*-
## @deftypefn {Function File} {@var{res} =} sf_dilog (@var{z})
## Compute the dilogarithm $Di_2(z) = \sum_{n=1}^{\infty} z^n/n^2$
## @end deftypefn
function res = sf_dilog(z)
    if (nargin < 1) print_usage; endif
    res = zeros(size(z));
    for n = 1:prod(size(z))
        res(n) = sf_dilog_1(z(n));
    endfor
endfunction
function res = sf_dilog_1(z)
    # branch cut on positive real axis [1,+infty)
    if (imag(z)==0 && z>=1) res = NaN; return; endif

    if (imag(z)==0 && z>1/2)
        # for efficiency for real values
        # L(z) + L(1-z) = pi^2/6 - (ln z)(ln 1-z)
        res = pi^2/6 - sf_log(z)*sf_log_p1(-z) - power_series(1-z);
    elseif (abs(z) <= 2/3)
        # direct series (radius of convergence for |z|<1)
        res = power_series(z);
    elseif (abs(z) >= 3/2)
        # reflect large values to small
        # L(z) + L(1/z) = -pi^2/6 - ln(-z)^2/2
        res = -pi^2/6 - sf_log(-z)^2/2 - power_series(1/z);
    else
        # clean up the rest
        # L(z) + L(z/(z-1)) = -ln(1-z)^2/2
        res = -sf_log(1-z)^2/2 - power_series(z/(z-1));
    endif
endfunction
function res = power_series(z)
    res = 0.0; n = 1; term = 1.0;
    do
        term *= z;
        old_res = res;
        res += term / (n^2);
        ++n; if (n>999) break; endif
    end
```

```
dilog(z) = Li2(z) (cont)
```

```
until (res == old_res)
endfunction
```

15.5.2 Polylogarithm

For real or complex s and z , the polylogarithm is defined by

$$\text{Li}_s(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^s}$$

For fixed $s \in \mathbb{C}$, the series defines an analytic function of z for $|z| < 1$. The series also converges when $|z| = 1$ if $\Re s > 1$. (For other values of z , define by analytic continuation).

We have the integral representation for $\Re s > 0$ and $|\text{ph } 1 - z| < \pi$, or $\Re s > 1$ and $z = 1$:

$$\text{Li}_s(z) = \frac{z}{\Gamma(s)} \int_0^{\infty} \frac{x^{s-1}}{e^x - z} dx$$

Finally we have the properties

$$\text{Li}_s(z) = \Gamma(1-s)(\ln 1/z)^{s-1} + \sum_{n=0}^{\infty} \zeta(s-n) \frac{(\ln z)^n}{n!} \quad |\ln z| < 2\pi, s \neq 1, 2, 3, \dots$$

and

$$\text{Li}_s(e^{2\pi i a}) + e^{\pi i s} \text{Li}_s(e^{-2\pi i a}) = \frac{(2\pi)^s e^{\pi i s/2}}{\Gamma(s)} \zeta(1-s, a)$$

valid when $\Re s > 0$, $\Im a > 0$; or $\Re s > 1$, $\Im a = 0$.

15.5.3 Fermi-Dirac and Bose-Einstein integrals

$$\text{FD}(k, \eta, \vartheta) = \int_0^{\infty} \frac{x^k (1 + \vartheta x/2)^{1/2}}{e^{x-\eta} + 1} dx \quad k > -1; \vartheta \geq 0$$

The DLMF [DLMF] defines the Fermi-Dirac integrals:

$$F_s(x) = \frac{1}{\Gamma(s+1)} \int_0^{\infty} \frac{t^s}{e^{t-x} + 1} dt \quad (s > -1)$$

and the Bose-Einstein integrals:

$$G_s(x) = \frac{1}{\Gamma(s+1)} \int_0^{\infty} \frac{t^s}{e^{t-x} - 1} dt \quad (s > -1, x < 0; \text{ or } s > 0, x \leq 0)$$

(“Sometimes the Gamma factor is omitted.”)

In terms of polylogarithms we have: $F_s(x) = -\text{Li}_{s+1}(-e^x)$ and $G_s(x) = \text{Li}_{s+1}(e^x)$.

We can integrate the definition directly using numerical quadrature ...

From [AppFD] we have:

- Start with an expansion for $\Re(x) < 0$:

$$F_q(x) = \sum_{n=1}^{\infty} (-)^{n-1} \frac{e^{xn}}{n^{q+1}}$$

- Consider the representation (with $p = q + 1$), where \mathcal{L} is a vertical contour cutting the real s -axis between 0 and 1:

$$F_{p-1}(x) = \frac{1}{2i} \int_{\mathcal{L}} \frac{e^{xs}}{s^p \sin \pi s} ds \quad p > 0, x > 0$$

- (etc.)

16 Coulomb wave functions

These are solutions $w(\rho)$ to the Coulomb (wave) equation, for $\ell = 0, 1, 2, \dots$; $\eta \in \mathbb{R}$; $\rho > 0$:

$$w'' + \left(1 - \frac{2\eta}{\rho} - \frac{\ell(\ell+1)}{\rho^2}\right) w = 0$$

The *normalizing constant* is defined

$$C_\ell(\eta) = \frac{2^\ell e^{-\pi\eta/2} |\Gamma(\ell+1+\eta\hat{i})|}{(2\ell+1)!} = \frac{2^\ell \left(\frac{2\pi\eta}{e^{2\pi\eta}-1} \prod_{k=1}^\ell (\eta^2 + k^2) \right)^{1/2}}{(2\ell+1)!}$$

[use $\exp(\log(\dots))$ in the product form to avoid some blow-up for large ℓ .] The *Coulomb phase shift* is defined (where the branch of phase is chosen as zero when $\eta = 0$)

$$\sigma_\ell(\eta) = \text{ph } \Gamma(\ell+1+\eta\hat{i})$$

and we define

$$\vartheta_\ell(\eta, \rho) = \rho - \eta \ln(2\rho) - \frac{\pi}{2}\eta + \sigma_\ell(\eta)$$

The solution, recessive at $\rho = 0$ is $F_\ell(\eta, \rho)$ (analytic on $\rho \in (0, \infty)$ and for $\eta \in \mathbb{R}$) defined via (where $M_{\kappa, \mu}(z)$ is the Whittaker function and $M(a, b, z)$ is the Kummer function)

$$\begin{aligned} F_\ell(\eta, \rho) &= C_\ell(\eta) \left(\frac{\mp \hat{i}}{2} \right)^{\ell+1} M_{\pm \eta \hat{i}, \ell+1/2}(\pm 2\rho \hat{i}) \\ &= C_\ell(\eta) \rho^{\ell+1} e^{\mp \rho \hat{i}} M(\ell+1 \mp \eta \hat{i}, 2\ell+2, \pm 2\rho \hat{i}) \end{aligned}$$

The irregular solutions (both analytic on $\rho \in (0, \infty)$ and with $e^{\mp \sigma \hat{i}} H$ analytic on $\eta \in \mathbb{R}$) are

$$\begin{aligned} H_\ell^\pm(\eta, \rho) &= G_\ell(\eta, \rho) \pm \hat{i} F_\ell(\eta, \rho) \\ &= (\mp \hat{i})^\ell e^{(\pi\eta/2) \pm \hat{i} \sigma_\ell(\eta)} W_{\mp \eta \hat{i}, \ell+1/2}(\mp 2\rho \hat{i}) \\ &= (\mp 2\rho \hat{i})^{\ell+1 \pm \eta \hat{i}} e^{\pm \hat{i} \vartheta_\ell(\eta, \rho)} U(\ell+1 \pm \eta \hat{i}, 2\ell+2, \mp 2\rho \hat{i}) \end{aligned}$$

Interrelations: $\mathcal{W}\{G_\ell, F_\ell\} = \mathcal{W}\{H_\ell^\pm, F_\ell\} = 1$ and $F_{\ell-1}G_\ell - F_\ell G_{\ell-1} = \frac{\ell}{\sqrt{\ell^2 + \eta^2}}$.

We have various recursion formula. Let $R_\ell = \sqrt{1 + \eta^2/\ell^2}$, $S_\ell = \ell/\rho + \eta/\ell$, and $T_\ell = S_\ell + S_{\ell+1}$, then with X_ℓ being $F_\ell(\eta, \rho)$ or $G_\ell(\eta, \rho)$, or $H_\ell^\pm(\eta, \rho)$, we have

$$R_\ell X_{\ell-1} - T_\ell X_\ell + R_{\ell+1} X_{\ell+1} = 0 \quad (\ell \geq 1)$$

$$X'_\ell = R_\ell X_{\ell-1} - S_\ell X_\ell \quad (\ell \geq 1)$$

$$X'_\ell = S_{\ell+1} X_\ell - R_{\ell+1} X_{\ell+1} \quad (\ell \geq 0)$$

[These recursions should be used in decreasing ℓ for regular solutions and increasing ℓ for irregular solutions.]

We have a series in ρ :

$$F_\ell(\eta, \rho) = C_\ell(\eta) \sum_{k=\ell+1}^{\infty} A_k^\ell(\eta) \rho^k$$

$$F'_\ell(\eta, \rho) = C_\ell(\eta) \sum_{k=\ell+1}^{\infty} k A_k^\ell(\eta) \rho^{k-1}$$

where $A_{\ell+1}^\ell = 1$, $A_{\ell+2}^\ell = \frac{\eta}{\ell+1}$, and $(k+\ell)(k-\ell-1)A_k^\ell = 2\eta A_{k-1}^\ell - A_{k-2}^\ell$ for $k = \ell+3, \ell+4, \dots$ (Alternatively, can write this as $A_k^\ell(\eta) = \frac{(-i)^{k-\ell-1}}{(k-\ell-1)!} {}_2F_1\left(\begin{matrix} \ell+1-k, \ell+1-\eta i \\ 2\ell+2 \end{matrix} \middle| 2\right)$.)

Various limiting forms:

As $\rho \rightarrow \infty$ with η fixed: $F_\ell(\eta, \rho) = \sin(\vartheta_\ell(\eta, \rho)) + o(1)$, $G_\ell(\eta, \rho) = \cos(\vartheta_\ell(\eta, \rho)) + o(1)$, and $H_\ell^\pm(\eta, \rho) \sim \exp(\pm i\vartheta_\ell(\eta, \rho))$.

For $\eta = 0$, $F_\ell(0, \rho) = \rho j_\ell(\rho) = \sqrt{\rho\pi/2} J_{\ell+\frac{1}{2}}(\rho)$, $G_\ell(0, \rho) = -\rho y_\ell(\rho) = -\sqrt{\rho\pi/2} Y_{\ell+\frac{1}{2}}(\rho)$, and $C_\ell(0) = \frac{2^\ell \ell!}{(2\ell+1)!} = \frac{1}{(2\ell+1)!!}$. (Note thus that $F_0(0, \rho) = \sin \rho$, $G_0(0, \rho) = \cos \rho$, and $H_0^\pm(0, \rho) = e^{\pm i\rho}$.)
etc.

17 Orthogonal polynomials

- name, symbol, parameters (and restrictions)
- differential equation (and second solution to DE); integral equation(s)
- integral representation(s)
- recurrence relation(s), “Rodrigues’ formula”
- representation(s) as hypergeometric series
- weight, interval — orthogonality relations, orthonormality weightings, ortho functions (basically including sqrt of weight)
- alternative tricks: chebyshev in terms of trig functions, etc.
- zeros, polynomial coeffs, ... Gauss integration weights, ...
- relationships (Jacobi \rightarrow Gegenbauer \rightarrow Chebyshev1&2/Legendre)

Computation of *general* orthogonal polynomials — compute coefficients / zeros / evaluate / etc. (Give arbitrary weight function or arbitrary zero locations or ...) Evaluate functions given as a sequence of orthogonal polynomials; compute such series representations.

function-evaluation, coefficients, zeros, weights, etc. Other orthogonal polynomials (...) generalized ... etc.

Note that *orthonormal* p_n satisfy

$$p_{n+1} - (a_n x + b_n) p_n + c_n p_{n-1} = 0$$

where $c_0 = 0$, $c_n = a_n/a_{n-1}$; $b_n = -a_n \langle x p_n, p_n \rangle$; $a_n = k_{n+1}/k_n$, with k_n being the coefficient of x^n in p_n .

Let

$$K_n(x, y) = \sum_{k=0}^n p_k(x) p_k(y) = \frac{k_n}{k_{n+1}} \frac{p_n(y) p_{n+1}(x) - p_n(x) p_{n+1}(y)}{x - y}$$

then we have

$$K_n(x, x) = \frac{k_n}{k_{n+1}} (p_n(x) p'_{n+1}(x) - p'_n(x) p_{n+1}(x)) = \sum_{k=0}^n p_k^2(x) > 0$$

For Gaussian quadrature we have

$$\int w(x)f(x) dx \approx \sum_{k=1}^n \lambda_{k,n} f(x_k)$$

where x_1, \dots, x_n being the zeros of p_n and where

$$\lambda_{k,n} = \left(\sum_{j=0}^{n-1} p_j(x_k)^2 \right)^{-1}$$

To find the zeros of orthogonal polynomials p_n (from SF book): suppose that p_n satisfy the recurrence (with $c_0 p_{-1} = 0$):

$$xp_k = a_k p_{k+1} + b_k p_k + c_k p_{k-1}$$

then the zeros of p_n are exactly the eigenvalues of the Jacobian matrix

$$\begin{pmatrix} b_0 & a_0 & 0 & & & \\ c_1 & b_1 & a_1 & 0 & & \\ 0 & c_2 & b_2 & a_2 & 0 & \\ & 0 & \ddots & \ddots & \ddots & 0 \\ & & 0 & a_{n-2} & b_{n-2} & a_{n-2} \\ & & & 0 & c_{n-1} & b_{n-1} \end{pmatrix}$$

For instance, for the generalized Laguerre polynomials we have $a_n = -(n+1)$, $b_n = (2n+\alpha+1)$, $c_n = -(n+\alpha)$. For Hermite H_n we have $a_n = 1/2$, $b_n = 0$, $c_n = n$. Note that due to numerical issues, an eigenvalue solver may find complex eigenvalues for large values of n (for some cases - Hermite especially, whereas Laguerre works much more effectively). Even if all eigenvalues found are real, it is useful to “polish” the accuracy by applying root-finders to improve the accuracy of the roots found (such as Newton — Halley’s method hasn’t been found to be very effective).

Alternative approach might be an iterative method using the roots of p_{n-1} to bracket the roots of p_n for a general root-finder.

An alternative symmetric matrix (whose eigenvalues give roots) is given by

$$\begin{pmatrix} \beta_0 & \alpha_0 & 0 & & & \\ \alpha_1 & \beta_1 & \alpha_1 & 0 & & \\ 0 & \alpha_2 & \beta_2 & \alpha_2 & 0 & \\ & 0 & \ddots & \ddots & \ddots & 0 \\ & & 0 & \alpha_{n-2} & \beta_{n-2} & \alpha_{n-2} \\ & & & 0 & \alpha_{n-1} & \beta_{n-1} \end{pmatrix}$$

(for details see NR4SF book...)

17.1 Chebyshev polynomials

17.1.1 First kind

$T_n(z)$

- $T_0 = 1$, $T_1 = x$
- Weight $w(x) = (1 - x^2)^{-1/2}$ on interval $(-1, 1)$
- Differential equation $(1 - x^2) T_n'' + (-x) T_n' + (n^2) T_n = 0$
- Normalization $(\pi/2)^{-1/2}$ for $n \neq 0$ and $(\pi)^{-1/2}$ for $n = 0$

- Relation $T_n(x) = \frac{n!}{(1/2)_n} P_n^{(-1/2, -1/2)}(x)$
- Recurrence $T_n = 2x T_{n-1} - T_{n-2}$
- Derivative $T'_n = \frac{n}{1-x^2} (T_{n-1} - x T_n)$
- Zeros
- Weights
- Rodrigues' formula $T_n(z) = (-2)^n \frac{n!}{(2n)!} (1-x^2)^{1/2} \frac{d^n}{dx^n} [(1-x^2)^{n-1/2}]$
- $T_n(x) = \cos(n \arccos x)$.

Shifted: $T_n^*(x) = T_n(2x-1)$ on $(0, 1)$ with weight $(x(1-x))^{-1/2}$

17.1.2 Second kind

$U_n(z)$

- $U_0 = 1, U_1 = 2x$
- Weight $w(x) = (1-x^2)^{1/2}$ on interval $(-1, 1)$
- Differential equation $(1-x^2) U''_n + (-3x) U'_n + (n(n+2)) U_n = 0$
- Normalization
- Relation $U_n(x) = \frac{(n+1)!}{(3/2)_n} P_n^{(1/2, 1/2)}(x)$
- Recurrence $U_n = 2x U_{n-1} - U_{n-2}$
- Derivative $U'_n = \frac{1}{1-x^2} ((n+1) U_{n-1} - nx U_n)$
- Zeros
- Weights
- Rodrigues' formula $U_n(z) = (-2)^n \frac{n+1}{2n+1} \frac{n!}{(2n)!} (1-x^2)^{-1/2} \frac{d^n}{dx^n} [(1-x^2)^{n+1/2}]$
-

Shifted: $U_n^*(x) = U_n(2x-1)$ on $(0, 1)$ with weight $(x(1-x))^{1/2}$

17.2 Legendre polynomials

$P_n(z)$

- $P_0 = 1, P_1 = x$
- Weight $w(x) = 1$ on interval $(-1, 1)$
- Differential equation $(1-x^2) P''_n + (-2x) P'_n + (n(n+1)) P_n = 0$
- Normalization $\sqrt{\frac{2}{2n+1}}$
- Relation $P_n(x) = P_n^{(0,0)}(x)$
- Recurrence $P_n = \frac{(2n-1)x}{n} P_{n-1} - \frac{n-1}{n} P_{n-2}$
- Derivative $P'_n =$
- Zeros
- Weights
- Rodrigues' formula $P_n(z) = \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2-1)^n]$
-

17.3 Laguerre (generalized) polynomials

$L_n^{(\alpha)}(z)$ with $\alpha > -1$

- $L_0^{(\alpha)} = 1, L_1^{(\alpha)} = 1 + \alpha - x$
- Weight $w(x) = e^{-x}x^\alpha$ on interval $(0, \infty)$
- Differential equation $(x) L_n'' + (\alpha + 1 - x) L_n' + (n) L_n = 0$
- Normalization $\sqrt{\frac{n!}{\Gamma(n+\alpha+1)}}$
- Relation
- Recurrence $L_n^{(\alpha)}(z) = \frac{2n+\alpha-1-z}{n} L_{n-1}^{(\alpha)}(z) - \frac{n+\alpha-1}{n} L_{n-2}^{(\alpha)}(z)$
- Derivative $L' = \frac{n}{x} (L_n - L_{n-1})$ [TODO: generalized form]
- Zeros
- Weights
- Rodrigues' formula $L_n^{(\alpha)}(z) = e^x \frac{x^{-\alpha}}{n!} \frac{d^n}{dx^n} [e^{-x} x^{n+\alpha}]$
-

17.4 Gegenbauer (ultraspherical) polynomials

$C_n^{(\gamma)}(z)$ with $\gamma \dots$

- Weight $w(x) = (1 - x^2)^{\gamma-1/2}$ on interval $(-1, 1)$
- Differential equation $(1 - x^2) C_n'' + (-(2\gamma + 1)x) C_n' + (n(n + 2\gamma)) C_n = 0$
- Normalization $\left(\frac{(2\gamma)_n}{(\gamma+1/2)_n} \frac{2^\gamma \Gamma(n+\gamma+1/2)}{\sqrt{2(n+\gamma)n! \Gamma(n+2\gamma)}} \right)^{-1}$
- Relation $C_n^{(\gamma)}(x) = \frac{(2+\gamma)_n}{(\gamma+1/2)_n} P_n^{(\gamma-1/2, \gamma-1/2)}(x)$
- Recurrence
- Zeros
- Weights
- Rodrigues' formula $C_n^{(\gamma)}(z) = \frac{d^n}{dx^n} []$
- Eigenvalue matrix terms $\beta_j = 0,$

$$\alpha_j = \frac{2}{2j + 2\gamma - 1} \sqrt{\frac{j(j + \gamma - 1/2)^2(j + 2\gamma - 1)}{(2j + 2\gamma)(2j + 2\gamma - 2)}} \quad j \geq 1$$

17.5 Jacobi polynomials

$P^{(\alpha,\beta)}(z)$ with $\alpha, \beta > -1$

- Weight $w(x) = (1-x)^\alpha(1+x)^\beta$ on interval $(-1, 1)$
- Differential equation $(1-x^2)P_n'' + (\beta - \alpha - (\alpha + \beta + 2)x)P_n' + (n(n + \alpha + \beta + 1))P_n = 0$
- Normalization $\sqrt{\frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)} \frac{\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)}}$
- Relation
- Recurrence
- Zeros
- Weights
- Rodrigues' formula $P_n^{(\alpha,\beta)}(z) = \frac{(-)^n}{2^n n!} (1-x)^{-\alpha}(1+x)^{-\beta} \frac{d^n}{dx^n} [(1-x)^{n+\alpha}(1+x)^{n+\beta}]$
- Eigenvalue matrix terms $\beta_0 = \frac{\beta-\alpha}{\alpha+\beta+2}$,

$$\beta_j = \frac{\beta^2 - \alpha^2}{(2j + \alpha + \beta)(2j + \alpha + \beta + 2)} \quad j \geq 1$$

$$\alpha_j = \frac{2}{2j + \alpha + \beta} \sqrt{\frac{j(j + \alpha)(j + \beta)(j + \alpha + \beta)}{(2j + \alpha + \beta + 1)(2j + \alpha + \beta - 1)}} \quad j \geq 1$$

Shifted: $R^{(\alpha,\beta)}(x) = P_n^{(\alpha,\beta)}(2x - 1)$ on $(0, 1)$ with weight $(1-x)^\alpha x^\beta$

17.6 Hermite polynomials

17.6.1 Physicists'

$H_n(z)$

- $H_0 = 1, H_1 = 2x$
- Weight $w(x) = e^{-x^2}$ on interval $(-\infty, \infty)$
- Differential equation $(1)H_n'' + (-2x)H_n' + (2n)H_n = 0$
- Normalization $(\sqrt{\pi}2^n n!)^{-1/2}$
- Relation
- Recurrence $H_n = 2xH_{n-1} - 2(n-1)H_{n-2}$
- Derivative $H_n' = 2nH_{n-1}$
- Zeros
- Weights
- Rodrigues' formula $H_n(z) = (-)^n e^{x^2} \frac{d^n}{dx^n} [e^{-x^2}]$
- Eigenvalue matrix terms $\beta_j = 0, \alpha_j = \sqrt{j/2}, \mu_0 = \sqrt{\pi}$

17.6.2 Probabilists'

$\text{He}_n(z)$

- Weight $w(x) = e^{-x^2/2}$ on interval $(-\infty, \infty)$
- Differential equation $(2) \text{He}_n'' + (-4x) \text{He}_n' + (2n) \text{He}_n = 0$ [double-check this]
- Normalization $(\sqrt{2\pi}n!)^{-1/2}$
- Relation $\text{He}_n(z) = 2^{-n/2} \text{H}_n(x/\sqrt{2})$
- Recurrence
- Zeros: zeros of $\text{He}_n(x)$ are $\sqrt{2} \times$ zeros of $\text{H}_n(x)$
- Weights
- Rodrigues' formula $\text{He}_n(z) = (-)^n e^{x^2/2} \frac{d^n}{dx^n} [e^{-x^2/2}]$
-

18 Misc functions

synchrotron, transport, Einstein, Lobachevsky, Erdelyi μ/ν ,

Heun

Lamé

Painlevé

18.1 Anger, Weber functions

[Move this section??]

18.1.1 Lommel functions

For $\mu \pm \nu \neq \pm 1, \pm 3, \pm 5, \dots$

$$S_{\mu,\nu}(z) = \frac{z^{\mu+1}}{(\mu+1)^2 - \nu^2} \sum_{k=0}^{\infty} t_k$$

where $t_0 = 1$ and $t_k = t_{k-1} \frac{-z^2}{(\mu+2k+1)^2 - \nu^2}$.

And the second Lommel function given via asymptotic expansion:

$$s_{\mu,\nu}(z) \sim \sum_{k=0}^{\infty} u_k$$

where $u_0 = 1$, $u_k = u_{k-1} \frac{-(\mu-2k+1)^2 - \nu^2}{z^2}$.

18.1.2 Anger function

$$\mathbf{J}_{\nu}(z) = \int_0^{\pi} \frac{\cos(\nu\vartheta - z \sin \vartheta)}{\pi} d\vartheta$$

We have

$$\mathbf{J}_{\nu}(z) = \frac{\sin \nu\pi}{\pi} S_{0,\nu}(z) - \nu \frac{\sin \nu\pi}{\pi} S_{-1,\nu}(z)$$

and

$$\mathbf{J}_{\nu}(z) = \mathbf{J}_{\nu}(z) + \frac{\sin \nu\pi}{\pi z} s_{0,\nu}(z) - \nu \frac{\sin \nu\pi}{\pi z^2} s_{-1,\nu}(z)$$

Power-series if $|z| < 10$ or $|z/\nu| < q??$

From [ACMF]: to get an array of values for index $\nu, \nu + 1, \dots, \nu + m$ with $0 < \nu < 1$:

- Compute $\mathbf{J}_\nu(x)$, $\mathbf{J}_{\nu+1}(x)$ by power-series for $x < 30$ and by asymptotic expansion for $x \geq 30$
- Use upwards recurrence in ν — for small x this can be bad, so we can use a downwards recurrence (with modifications to the previous step)
- For the special case of $x = 0$, use the fact that $\mathbf{J}_\nu(0) = \sin(\nu\pi)/(\nu\pi)$ exactly

18.1.3 Weber function

$$\mathbf{E}_\nu(z) = \int_0^\pi \frac{\sin(\nu\vartheta - z \sin \vartheta)}{\pi} d\vartheta$$

This is computed the same basic way as $\mathbf{J}_\nu(x)$ above... [ACMF].

We have

$$\mathbf{E}_\nu(z) = -\frac{1 + \cos \nu\pi}{\pi} S_{0,\nu}(z) - \nu \frac{1 - \cos \nu\pi}{\pi} S_{-1,\nu}(z)$$

and

$$\mathbf{E}_\nu(z) = -Y_\nu(z) + \frac{1 + \cos \nu\pi}{\pi} s_{0,\nu}(z) - \nu \frac{1 - \cos \nu\pi}{\pi z^2} s_{-1,\nu}(z)$$

[CHECK THE DENOMINATOR ON THE FIRST FRACTION IN THE PREVIOUS EQUATION! (compare with Anger function...)]

18.2 Howland integrals

$$I_k = \frac{1}{2(k!)} \int_0^\infty \frac{w^k}{\sinh w + w} dw \quad (k \geq 1)$$

$$I_k^* = \frac{1}{2(k!)} \int_0^\infty \frac{w^k}{\sinh w - w} dw \quad (k \geq 3)$$

18.3 Mathieu functions

(elliptic cylinder functions) [This needs to be cleaned up!!] [MOVE TO ITS OWN SECTION!]

characteristic values, expansion coefficients, sem, cem dsem, dsem, McM, dMcM, MsM, dMsM, etc. ...

From Zhang-Jin [ZJ] Mathieu's differential equation is

$$y'' + (\lambda - 2q \cos 2z)y = 0$$

And note that due to periodicity requirements, there are conditions on $\lambda = \lambda(q)$ as a function of q . Let $\lambda(0) = m^2$, then we get the following Mathieu functions of order m :

$$\text{ce}_m(z, q) = \sum_{k=0}^{\infty} A_k^m(q) \cos kz$$

$$\text{se}_m(z, q) = \sum_{k=0}^{\infty} B_k^m(q) \cos kz$$

for some coeffs $A, B...$

18.3.1 Modified Mathieu functions

$$y'' - (\lambda - 2q \cosh 2z)y = 0$$

$$\text{Ce}_m(z, q) = \text{ce}_m(z, q)(\hat{iz}, q)$$

$$\text{Se}_m(z, q) = -\hat{i} \text{se}_m(z, q)(\hat{iz}, q)$$

$$\text{Mc}_m^{(j)}(z, q) = (j = 1..4)$$

$$\text{Ms}_m^{(j)}(z, q) = (j = 1..4)$$

Matheiu-Hankel for $j = 3, 4$
 ? others: ?

$$\text{Cc}_m^{(1)}(z, q) =$$

$$\text{Cs}_m^{(1)}(z, q) =$$

$$\text{Cc}_m^{(2)}(z, q) =$$

$$\text{Cs}_m^{(2)}(z, q) =$$

$$\text{Cc}_m^{(3)}(z, q) =$$

$$\text{Cs}_m^{(3)}(z, q) =$$

$$\text{Cc}_m^{(4)}(z, q) =$$

$$\text{Cs}_m^{(4)}(z, q) =$$

$$\lambda(q) =$$

18.4 Debye functions

$$\text{D}_n(x) = \int_0^x \frac{t^n}{e^t - 1} dt$$

$$\tilde{\text{D}}_n(x) = \frac{n}{x^n} \text{D}_n(x)$$

A series representation (compare with generating function for Bernoulli numbers) (for $|x| < 2\pi$ and $n \geq 1$):

$$\int_0^x \frac{t^n}{e^t - 1} dt = x^n \left(\frac{1}{n} - \frac{x}{2(n+1)} + \sum_{k=1}^{\infty} \frac{B_{2k}}{(2k+n)(2k)!} x^{2k} \right)$$

This series works ok, but requires increasing number of terms as n or x increase (for $n = 1$, $x = 1$ we already need 13 terms, for $n = 6$, $x = 3$ we need 31 terms).

Reflection:

$$\text{D}_n(-x) = (-)^n \text{D}_n(x) + (-)^n \frac{x^{n+1}}{n+1}$$

$$\tilde{\text{D}}_n(-x) = \tilde{\text{D}}_n(x) + \frac{n}{n+1} x$$

Also for the complementary integral (for $x > 0$ and $n \geq 1$):

$$\int_x^{\infty} \frac{t^n}{e^t - 1} dt = \sum_{k=1}^{\infty} e^{-kx} \left(\frac{x^n}{k} + \frac{nx^{n-1}}{k^2} + \frac{n(n-1)x^{n-2}}{k^3} + \dots + \frac{n!}{k^{n+1}} \right)$$

Note that

$$\int_0^\infty \frac{t^n}{e^t - 1} dt = n! \zeta(n+1)$$

The complementary integral is useful for $|x| \geq 2$ but for large n (say $n > 20$) we get bad cancellation.
Note

$$D_n(x) = n! \zeta(n+1) - \int_x^\infty \frac{t^n}{e^t - 1} dt = n! \left\{ [\zeta(n+1) - 1] + e^{-x} \left[\sum_{j=n+1}^\infty \frac{z^j}{j!} \right] - \sum_{k=2}^\infty \frac{e^{-kx}}{k^{n+1}} \left[\sum_{j=0}^n \frac{z^j}{j!} \right] \right\}$$

which might offer improved numerical stability (with bracketed terms computed individually with high precision)

Asymptotic expansion:

$$D_n(x) \sim n! \zeta(n+1) - e^{-x} \cdot n \cdot n! \sum_{k=0}^n \frac{x^k}{k!} + \log(1 - \cosh(x) + \sinh(x)) x^n$$

18.5 Sievert integral

$$S(\vartheta, x) = \int_0^\vartheta e^{-x \sec \varphi} d\varphi = \int_0^\vartheta e^{-x/\cos \varphi} d\varphi$$

$$\tilde{S}(\vartheta, x) = \frac{S(\vartheta, x)}{\vartheta}$$

From [A&S]: As $x \rightarrow \infty$,

$$S(x, \vartheta) \sim \sqrt{\frac{\pi}{2x}} e^{-x} \operatorname{erf}(\vartheta \sqrt{x/2})$$

This can be derived by expanding $\sec \varphi$ as a power-series at zero, taking the first 2 terms: $\sec \varphi = 1 + \frac{\varphi^2}{2} + O(\varphi^4)$ and integrating the resulting integral: $\int_0^\vartheta e^{-z(1+t^2/2)} dt$.

Also, if $a_0 = 1$, $a_n = \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots (2n)}$ then

$$S(x, \vartheta) = S(x, \pi/2) - \sum_{k=0}^\infty \alpha_k \cos(\vartheta)^{2k+1} E_{2k+2}\left(\frac{x}{\cos \vartheta}\right)$$

for $x \geq 0$, $0 < \vartheta < \pi/2$. ($E_n(z) = \int_1^\infty e^{-zt} t^{-n} dt$ are exponential integrals.)

$$S(x, \pi/2) = \operatorname{Ki}_1(x) = \int_x^\infty K_0(t) dt$$

We can derive the following expansion by expanding the exponential and integrating, but it doesn't seem particularly helpful for calculations:

$$S(x, \vartheta) = \sum_{n=0}^\infty (-x)^n {}_2F_1\left(\frac{1}{2}, (n+1)/2 \middle| \sin^2 \vartheta\right)$$

18.6 Abramowitz functions

$$f_m(x) = \int_0^\infty t^m e^{-t^2 - x/t} dt$$

for $m = 0, 1, 2, \dots$

A few potentially useful facts (from [A&S]):

- A recurrence: $2f_m = (m-1)f_{m-2} + xf_{m-3}$
- $f'_m = -f_{m-1}$ ($m = 1, 2, \dots$)
- A series expansion for f_1 :

$$f_1(x) = \frac{1}{2} \sum_{k=0}^{\infty} (a_k \ln x + b_k) x^k$$

where

$$\begin{aligned} - a_0 &= a_1 = 0; a_2 = -b_0 \\ - b_0 &= 1; b_1 = -\sqrt{\pi}; b_2 = \frac{3}{2}(1-\gamma) \\ - a_k &= \frac{-2a_{k-2}}{k(k-1)(k-2)} \\ - b_k &= \frac{-2b_{k-2} - (3k^2 - 6k + 2)a_k}{k(k-1)(k-2)} \end{aligned}$$

- An asymptotic expansion, let $v = 3(x/2)^{2/3}$, then

$$f_m(x) \sim \sqrt{\frac{\pi}{3}} 3^{-m/2} v^{m/2} e^{-v} \left(a_0 + \frac{a_1}{v} + \frac{a_2}{v^2} + \dots \right)$$

where $a_0 = 1$; $a_1 = \frac{1}{12}(3m^2 + 3m - 1)$; and $12(k+2)a_{k+2} = -(12k^2 + 36k - 3m^2 - 3m + 25)a_{k+1} + \frac{1}{2}(m-2k)(2k+3-m)(2k+3+2m)a_k$

18.7 Abramowitz 2 functions

$$f(x) = \int_0^{\infty} \frac{e^{-t^2}}{t+x} dt$$

Representation in terms of exponential integral and Dawson's integral:

$$f(x) = -\frac{1}{2}e^{-x^2} \text{Ei}(x^2) + \sqrt{\pi}e^{-x^2} \int_0^x e^{t^2} dt$$

Series expansion:

$$f(x) = -e^{-x^2} \ln x + e^{-x^2} \left[\sqrt{\pi} \sum_{k=0}^{\infty} \frac{x^{2k+1}}{k!(2k+1)} - \sum_{k=1}^{\infty} \frac{x^{2k}}{k!(2k)} - \frac{\gamma}{2} \right]$$

or a series with the digamma function:

$$f(x) = -e^{-x^2} \ln x + \frac{1}{2} \sum_{k=0}^{\infty} \frac{\psi(k+1)x^{2k}}{k!} + \sqrt{\pi} \sum_{k=0}^{\infty} \frac{(-2)^k x^{2k+1}}{1 \cdot 3 \cdot 5 \cdots (2k+1)}$$

18.8 Clausen integral

[and generalizations...] For $0 \leq \vartheta \leq \pi$:

$$C(\vartheta) = -\int_0^{\vartheta} \ln(2 \sin t/2) dt = \sum_{n=1}^{\infty} \frac{\sin n\vartheta}{n^2} = \Im(\text{Li}_2(e^{i\vartheta}))$$

Another series:

$$C(\vartheta) = -\vartheta \ln \vartheta + \vartheta + \sum_{k=1}^{\infty} (-)^{k-1} \frac{B_{2k}}{(2k)!} \frac{\vartheta^{2k+1}}{2k(2k+1)}$$

or equivalently:

$$\frac{C(\vartheta)}{\vartheta} = 1 - \ln \vartheta + \sum_{k=1}^{\infty} \frac{\zeta(2k)}{n(2n+1)} \left(\frac{\vartheta}{2\pi} \right)^{2k}$$

Another series (from wikipedia)

$$C(\vartheta) = 3\vartheta - \vartheta \ln \left(\vartheta \left(1 - \frac{\vartheta^2}{4\pi^2} \right) \right) - 2\pi \ln \left(\frac{2\pi + \vartheta}{2\pi - \vartheta} \right) + \vartheta \sum_{n=1}^{\infty} \frac{\zeta(2n) - 1}{n(2n+1)} \left(\frac{\vartheta}{2\pi} \right)^n$$

[check and test this...]

Relation:

$$C(\pi - \vartheta) = C(\vartheta) - \frac{1}{2} C(2\vartheta) \quad (0 \leq \vartheta \leq \pi/2)$$

18.9 Voigt (plasma dispersion) function

$$V(u, \alpha) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{e^{-(\alpha y - u)^2}}{y^2 + 1} dy$$

$$V(u, \alpha) = \Re(e^{z^2} \operatorname{erfc}(z)) \quad z = \alpha + u\hat{i}$$

The versions from DLMF:

$$U(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} \frac{e^{-(x-y)^2/4t}}{1+y^2} dy$$

$$V(x, t) = \frac{1}{\sqrt{4\pi t}} \int_{-\infty}^{\infty} \frac{ye^{-(x-y)^2/4t}}{1+y^2} dy$$

then with $z = \frac{1-i\hat{x}}{2\sqrt{t}}$ we have

$$U(x, t) + \hat{i} V(x, t) = \sqrt{\frac{\pi}{4t}} e^{z^2} \operatorname{erfc}(z)$$

The “line-broadening function” (from DLMF) is defined by (and related to Voigt functions via):

$$H(a, u) = \frac{a}{\pi} \int_{-\infty}^{\infty} \frac{e^{-t^2}}{(u-t)^2 + a^2} dt = \frac{1}{a\sqrt{\pi}} U\left(\frac{u}{a}, \frac{1}{4a^2}\right)$$

18.10 Spence integral

$$S(x) = - \int_1^x \frac{\ln t}{t-1} dt = \operatorname{Li}_2(1-x) \quad \Re x \geq 0 \vee x \notin \mathbb{R}$$

Implementation notes for $S(z)$, $z \geq 0$:

- Recall $\operatorname{Li}_2(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^2}$ (with $z < 1$). Recall the following identities (and their applicability when $z \in \mathbb{R}$)

(A) $\operatorname{Li}_2(z) + \operatorname{Li}_2\left(\frac{1}{z}\right) = -\frac{\pi^2}{6} - \frac{\ln(-z)^2}{2}$ for $z < 0$

(B) $\operatorname{Li}_2(z) + \operatorname{Li}_2(1-z) = \frac{\pi^2}{6} - \ln(z) \ln(1-z)$ for $0 < z < 1$

(C) $\operatorname{Li}_2(z) + \operatorname{Li}_2\left(\frac{z}{z-1}\right) = -\frac{\ln(1-z)^2}{2}$ for $z < 1$

- Using these identities, we have the following computations:

– For $x = 0$ use $S(0) = \frac{\pi^2}{6}$

- For $0 < x < 0.5$, use **(B)** and compute $S(x) = \frac{\pi^2}{6} - \ln(x) \ln(1-x) - \sum_{k=1}^{\infty} \frac{x^k}{k^2}$
- For $0.5 \leq x < 1$, use $S(x) = \text{Li}_2(1-x)$ and compute $S(x) = \sum_{k=1}^{\infty} \frac{(1-x)^k}{k^2}$
- For $1 \leq x < 2.5$, use **(C)** and compute $S(x) = -\frac{\ln(x)^2}{2} - \sum_{k=1}^{\infty} \frac{1}{k^2} \left(\frac{x-1}{x}\right)^k$
- For $2.5 \leq x$, use **(A)** and compute $S(x) = -\frac{\pi^2}{6} - \frac{\ln(x-1)^2}{2} - \sum_{k=1}^{\infty} \frac{(1-x)^{-k}}{k^2}$

- This gives 15 digits precision everywhere but up to 7 ulps near 1
- Generally have less than 50 terms needed everywhere in the domain, as $x \rightarrow \infty$, need fewer terms
- Atlas recommends series near 0 and 1 and in other cases to integrate $S(x) = -\int_0^{x-1} \frac{\ln 1+u}{u} dt$ using Simpson's rule (140 points for $x \leq 2$ and more points for larger x)

18.11 Angular momentum coupling coefficients

Define $\Delta(abc) = \sqrt{\frac{(a+b-c)!(a-b+c)!(-a+b+c)!}{(a+b+c+1)!}}$

18.11.1 3-j coefficients

The sum below is over all nonnegative integers s such that factorial arguments are nonnegative:

$$\begin{aligned} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} &= (-)^{a-b-\gamma} \Delta(abc) \sqrt{(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!(c-\gamma)!} \\ &\cdot \sum_s \frac{(-)^s}{s!(a+b-c-s)!(a-\alpha-s)!(b+\beta-s)!(c-b+\alpha+s)!(c-a-\beta+s)!} \end{aligned}$$

Alternatively:

$$\begin{aligned} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} &= \delta_{\alpha+\beta+\gamma,0} (-)^{a-b-\gamma} \\ &\cdot \sqrt{\frac{(c+a-b)!(c-a+b)!(a+b-c)!(c-\gamma)!(c+\gamma)!}{(a+b+c+1)!(a-\alpha)!(a+\alpha)!(b-\beta)!(b+\beta)!}} \\ &\cdot \sum_k' \frac{(-)^{k+b+\beta} (b+c+\alpha-k)!(a-\alpha+k)!}{k!(c-a+b-k)!(c-\gamma-k)!(k+a-b+\gamma)!} \end{aligned}$$

18.11.2 6-j coefficients

$$\begin{aligned} \begin{Bmatrix} a & b & c \\ d & e & f \end{Bmatrix} &= (-)^{a+b+c+d} \Delta(abe) \Delta(acf) \Delta(bdf) \Delta(cde) \\ &\cdot \sum_k' (a+b+c+d+1-k)! \\ &\quad (k!(e+f-a-d+k)!(e+f-b-c+k)!)^{-1} \\ &\quad ((a+b-e-k)!(c+d-e-k)!(a+c-f-k)!(b+d-f-k-k)!)^{-1} \end{aligned}$$

18.11.3 9-j coefficients

$$\begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{Bmatrix} = \sum_k (-)^{2k} (2k+1) \begin{Bmatrix} a & i & k \\ h & d & g \end{Bmatrix} \begin{Bmatrix} b & f & k \\ d & h & e \end{Bmatrix} \begin{Bmatrix} a & i & k \\ f & b & c \end{Bmatrix}$$

19 Combinatorial functions and numbers

Fibonacci numbers/polynomials, Lucas numbers/polynomials, Stirling numbers (1/2), factorial, binomial, multinomial, Pochhammer symbol, rising/falling factorial, generalized Lucas/Fibonacci numbers, ((brackets and numbers from “concrete mathematics”))

integer & generalized versions

Bell numbers = # of partitions of set. Touchard's congruence: $B_{p+k} \cong B_k + B_{k+1} \pmod{p}$ where p is prime and B_n is a Bell number. $e^{e^x-1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}$. Bell polynomials — Dobinski's formula: $B_n(x) = e^{-x} \sum_{k=0}^{\infty} k^n \frac{x^k}{k!}$.

19.1 Bernoulli numbers, polynomials

$$\frac{t}{e^t - 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} B_n$$

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} B_n(x)$$

Fact: $\zeta(2n) = \frac{(2\pi)^{2n}}{2} \frac{|B_{2n}|}{(2n)!}$ which implies that $|B_{2n}| \sim \frac{2(2n)!}{(2\pi)^{2n}}$ for large n ... ** Use ζ to generate B_n for large/scaled n (very easy to compute)...

Compute: B_n , $B_n/n!$, $\ln|B_n|$, etc.

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_k x^{n-k}$$

$$B_n(1-x) = (-)^n B_n(x)$$

$$B_n(x+1) - B_n(x) = nx^{n-1}$$

19.2 Euler numbers, polynomials

$$\frac{1}{\cosh t} = \frac{2e^t}{e^{2t} + 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} E_n \quad |z| < \frac{\pi}{2}$$

$$\frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} E_n(x) \quad |z| < \pi$$

$$E_n = 2^n E_n\left(\frac{1}{2}\right) \in \mathbb{Z}$$

$$E_n(x) = \sum_{k=0}^n \binom{n}{k} \frac{E_k}{2^k} \left(x - \frac{1}{2}\right)^{n-k}$$

$$E_n(1-x) = (-)^n E_n(x)$$

$$E_n(x+1) + E_n(x) = 2x^n$$

We also have the *secant numbers*, with $\sec x = \sum_{k=0}^{\infty} S_k \frac{x^{2k}}{(2k)!}$:

$$S_n = |E_{2n}| = E_n^*$$

19.3 Tangent numbers, polynomials

$$T_{2n-1} = \frac{2^{2n}(2^{2n}-1)|B_{2n}|}{2n} = (-)^{n-1} \frac{2^{2n}(2^{2n}-1)B_{2n}}{2n} \in \mathbb{Z}$$

and $T_{2n} = 0$ with $\tan x = \sum_{k=1}^{\infty} T_k \frac{x^k}{k!}$ Note that

19.4 Genocchi numbers, polynomials

$$\frac{2t}{e^t + 1} = \sum_{n=1}^{\infty} G_n \frac{t^n}{n!}$$

whence $G_n = 2(1 - 2^n)B_n$. We can define the scaled numbers also $G_n^* = G_n/n!$.

19.5 Number theoretic functions

for unsigned long long (64-bit): isprime, factor (note ≤ 16 distinct factors of $n \leq 2^{64}$), nth_prime, Chinese Remainder Theorem, Möbius function, Jacobi function, σ , τ , etc.

20 Mathematical finance

digital, power, log, OT, NT, DOT, DNT, KI, KO, DKI, DKO, sequentials, windowed, aisan, lookback, ratchet, cliquet, forward-start, chooser, compound, reset, extendible, fader, accumulator, accrual, digital-accrual, ..., fadelets, ..., quanto, basket,

implied-volatility function

various associated functionals of Brownian motion: pdf, hitting-time pdfs, BF with drift, lognormal, local-times, etc.

20.1 Brownian motion

pdf/cdf of BM at time T (with starting point); with drift/volatility; pdf/cdf of hitting time BH (w/drift) for level L , H , L and H , L before H , etc.; conditional density of B_T on not hitting barrier(s); pdf/cdf of minimum/maximum before T ; joint pdf of B_T , M_T , m_T (& combinations); occupation times; average densities (Asians...); joint densities of B_{T_1} , B_{T_2} , etc.;

Brownian bridge; lognormal;

copulas?

sampling?; generating Weiner paths?; generic SDE simulator? (Euler/Milstein/etc.)

20.2 Bachelier model

20.3 Black-Scholes-Merton (BS) model

$$d_{\pm}(k, t, s, r, g, \sigma) = \frac{\ln(s/k) + (r - g \pm \sigma^2/2)t}{\sigma\sqrt{t}}$$

20.3.1 Vanilla

$$\text{Prem}_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \omega e^{-gt} s N(\omega d_+) - \omega e^{-rt} k N(\omega d_-)$$

$$\Delta_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \frac{\partial P}{\partial s} = \omega e^{-gt} N(\omega d_+)$$

$$\Gamma_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \frac{\partial^2 P}{\partial s^2} = n(d_+) \frac{e^{-gt}}{\sigma\sqrt{t}}$$

$$\text{Ve}_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \frac{\partial P}{\partial \sigma} = n(d_+) s e^{-gt} \sqrt{t}$$

$$\text{Va}_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \frac{\partial^2 P}{\partial s \partial \sigma} = n(d_+) \frac{-e^{-gt} d_-}{\sigma}$$

$$\text{Vo}_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \frac{\partial^2 P}{\partial \sigma^2} = \text{Ve} \frac{d_+ d_-}{\sigma}$$

$$\rho_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \frac{\partial P}{\partial r} = \omega t k e^{-rt} N(\omega d_-)$$

$$\varphi_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = \frac{\partial P}{\partial g} = -\omega t s e^{-gt} N(\omega d_+)$$

$$\vartheta_{\text{BS}}^{\text{V}}(k, t, \omega; s, r, g; \sigma) = -\frac{\partial P}{\partial t} = -\frac{s\sigma e^{-gt} n(d_+)}{2\sqrt{t}} + \omega g s e^{-gt} N(d_+) - \omega r k e^{-rt} N(d_-)$$

20.4 Heston stochastic-volatility model

$$\text{Prem}_{\text{H}}^{\text{Van}}(k, t; s, r, g; \dots) =$$

20.5 Variance-gamma (VG) model

$$\Psi(a, b, \gamma) = \frac{1}{\Gamma(\gamma)} \int_0^\infty N[au^{-1/2} + bu^{1/2}] u^{\gamma-1} e^{-u} du$$

$$\Phi(\alpha, \beta, \gamma; x, y) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int_0^1 u^{\alpha-1} (1-u)^{\gamma-\alpha-1} (1-ux)^{-\beta} e^{uy} du$$

$$\text{Prem}_{\text{VG}}^{\text{Van}}(k, t; s, r, g; \dots) =$$

20.6 Constant-elasticity-of-variance (CEV) model

$$\text{Prem}_{\text{CEV}}^{\text{Van}}(k, t; s, r, g; \dots) =$$

21 Probability functions

pdf, cdf, co-cdf, inverse cdf, inverse co-cdf, shifted/scaled versions (give mean/variance *or* location/scale parameters), moments (mean, variance, general),

sampling, generation, integration against density, fitting, etc.

normal, lognormal, t, gamma, exponential, logistic, F, χ^2 , beta, cauchy, Weibull, Bessel (I), α -stable, Pareto, Laplace, inverse-gamma, inverse-normal, Bernoulli, Cauchy-Lorentz, “extreme-value distribution”, erlang, inverse χ^2 , non-centered beta/F/t, Rayleigh, triangular, uniform, “logarithmic series distribution”
bivariate normal, trivariate normal, n-variate normal,
binomial, poisson, hypergeometric, Kolmogorov-Smirnov, negative binomial, geometric
copulas

22 Useful techniques

22.1 Misc

22.1.1 Euler-Maclaurin summation

A few results for possible use:

$$\begin{aligned} \int \frac{1}{x^s} dx &= \frac{x^{1-s}}{1-s} \\ \int \frac{1}{(x+a)^s} dx &= \frac{(x+a)^{1-s}}{1-s} \\ \int \frac{z^x}{x^s} dx &= -x^{1-s} \Gamma(1-s, -x \ln z) (-x \ln z)^{s-1} \\ \int \frac{z^x}{(x+a)^s} dx &= -(x+a)^{1-s} \Gamma(1-s, -(x+a) \ln z) (-(x+a) \ln z)^{s-1} \end{aligned}$$

The Euler-Maclaurin summation formula

$$\sum_{n=a}^b f(n) = \int_a^b f(t) dt - \frac{f(b) + f(a)}{2} + \sum_{j=2}^k \frac{B_j}{j!} \left(f^{(j-1)}(b) - f^{(j-1)}(a) \right) - \int_a^b \frac{B_k(\{1-t\})}{k!} f^{(k)}(t) dt$$

where $\{x\} \in [0, 1)$ is the fractional part of x and B_j , $B_j(x)$ are Bernoulli numbers, polynomials (resp.).

If f and its derivatives go to zero as $x \rightarrow \infty$ then we have

$$\sum_{n=a}^{\infty} f(n) = \int_a^{\infty} f(t) dt - \frac{f(a)}{2} - \sum_{j=2}^k \frac{B_j}{j!} f^{(j-1)}(a) - \int_a^{\infty} \frac{B_k(\{1-t\})}{k!} f^{(k)}(t) dt$$

This is very useful for getting accurate approximations to the tail of truncated summations...

22.1.2 Boole summation

Let $h \in [0, 1]$, $f : [m, \infty) \rightarrow \mathbb{R}$ with k continuous derivatives, $f^{(i)}(x) \rightarrow 0$ as $x \rightarrow \infty$ for $i = 0, \dots, k$, then

$$\sum_{i=m}^{\infty} (-)^{i-m} f(h+i) = \frac{1}{2} \sum_{i=0}^{k-1} \frac{E_i(h)}{i!} f^{(i)}(m) + R_k$$

(recall that the Euler numbers E_n are related to the Euler polynomials $E_n = E_n(1/2)$ and are $E_{2n+1} = 0$, $E_0 = 1$, $E_2 = -1$, $E_4 = 5$, $E_6 = -61$, $E_8 = 1385$, ...).

22.1.3 Borel summation

(From Wikipedia)

Let $y(z) = \sum_{k=0}^{\infty} y_k z^k$ be a formal power series in z . Define the Borel transform of y

$$\mathcal{B}_y(t) = \sum_{k=0}^{\infty} \frac{y_k}{k!} t^k$$

Suppose that \mathcal{B}_y converges to an analytic function near 0 that can be analytically continued along the positive real axis to a function growing sufficiently slowly that the following integral is well-defined (as an improper integral). Then the *Borel sum* of y is given by

$$\int_0^{\infty} e^{-t} \mathcal{B}_y(tz) dt$$

A slightly weaker form of Borel's summation method gives the Borel sum of y as

$$\lim_{t \rightarrow \infty} e^{-t} \sum_n \frac{t^n}{n!} \cdot \sum_{k \leq n} y_k z^k$$

If the sum exists in this sense then it also exists in the previous sense and is the same, but there are some series that can be summed with the previous method but not with this method.

Example: $y(z) = \sum_{k=0}^{\infty} z^k = \frac{1}{1-z}$ for $|z| < 1$. Then $\mathcal{B}_y(t) = \sum_{k=0}^{\infty} \frac{1}{k!} t^k = e^t$ and the Borel sum is $\int_0^{\infty} e^{-t} e^{tz} dt = \frac{1}{1-z}$ for $\Re z < 1$, giving analytic continuation of the original series to a larger region.

Example: $y(z) = \sum_{k=0}^{\infty} k!(-z)^k$ diverges for all $z \neq 0$. Then $\mathcal{B}_y(t) = \sum_{k=0}^{\infty} (-t)^k = \frac{1}{1+t}$ and the Borel sum is $\int_0^{\infty} \frac{e^{-t}}{1+t} dt = \frac{1}{z} e^{1/z} \Gamma(0, 1/z)$.

22.1.4 Kahan's compensated summation trick

Note that optimization will destroy these algorithms!!

To sum up the numbers x_1, \dots, x_n :

```

s = 0; e = 0;
for i = 1:n
    temp = s
    y = x[i] + e
    s = temp + y
    e = (temp - s) + y
end
s = s + e // optional correction

```

Then s is a more accurate summation.

22.2 Continued fractions

Add Wallis algorithm also?

22.2.1 Pincherle's theorem

(From Temme: "Numerical Aspects of Special Function") Pincherle's theorem: given a three-term recurrence relation

$$y_{n+1} + b(n)y_n + a(n)y_{n-1} = 0$$

then the continued fraction

$$\frac{-a_k}{b_k +} \frac{-a_{k+1}}{b_{k+1} +} \dots$$

converges iff the recurrence relation possesses a minimal solution. Furthermore, if f_n is a minimal solution, then the continued fraction converges to f_k/f_{k-1} .

22.2.2 Modified Lentz algorithm

(cribbed from Numerical Recipes) Suppose $f(x) = b_0 + \frac{a_1}{b_1 +} \frac{a_2}{b_2 +} \frac{a_3}{b_3 +} \dots$. Then compute the following (where $\varepsilon \sim 10^{-15}$ and $\zeta \sim 10^{-30}$):

- Set $f_0 = b_0$ (if $b_0 = 0$ then set $f_0 = \zeta$)
- Set $C_0 = f_0$
- Set $D_0 = 0$
- For $j = 1, 2, \dots$
 - Set $D_j = b_j + a_j D_{j-1}$ (if $D_j = 0$ then set $D_j = \zeta$)
 - Set $C_j = b_j + a_j / C_{j-1}$ (if $C_j = 0$ then set $C_j = \zeta$)
 - Set $D_j = 1/D_j$
 - Set $\Delta_j = C_j D_j$
 - Set $f_j = f_{j-1} \Delta_j$
 - if $|\Delta_j - 1| < \varepsilon$ then exit.

22.2.3 Quotient-difference algorithm

Given a series $f(z) = c_0 + c_1 + c_2 z^2$, we can find an equivalent continued fraction via the table

$$\begin{array}{ccccccc}
 & & q_1^0 & & & & \\
 e_0^1 & & & e_1^0 & & & \\
 & q_1^1 & & & q_2^0 & & \\
 e_0^2 & & e_1^1 & & e_2^0 & & \\
 \vdots & \vdots & \vdots & \vdots & & \ddots &
 \end{array}$$

where $e_0^n = 0$, $q_1^n = \frac{c_{n+1}}{c_n}$ and we continue via the rules

$$\begin{aligned} e_j^k &= e_{j-1}^{k+1} + (q_j^{k+1} - q_j^k) \\ q_{j+1}^k &= q_j^{k+1} \frac{e_j^{k+1}}{e_j^k} \end{aligned}$$

Then we get the continued fraction

$$c = \frac{a_0}{1-} \frac{a_1 z}{1-} \frac{a_2 z}{1-} \dots$$

where $a_0 = c_0$, $a_1 = q_1^0$, $a_2 = e_1^0$, $a_3 = q_2^0$, $a_4 = e_2^0$, \dots

22.3 Sequence acceleration

22.3.1 Aitken acceleration

Given $\{x_n\}_n$, produce

$$\alpha_n = \frac{x_{n+2}x_n - x_{n+1}^2}{x_{n+2} - 2x_{n+1} + x_n} = x_n - \frac{(\Delta x_n)^2}{\Delta^2 x_n}$$

where $\Delta x_n = x_{n+1} - x_n$ and $\Delta^2 x_n = x_{n+2} - 2x_{n+1} + x_n$.

22.3.2 Cohen-Villegas-Zagier convergence acceleration of alternating series

Suppose we have $S = \sum_{k=0}^{\infty} (-)^k a_k$. Then fixing N , we compute

```
Let d = (3 + sqrt(8)) ^ N;
d = (d + 1/d) / 2;
b = -1; c = -d; s = 0;
for k = 0 : (N-1)
    c = b - c;
    s = s + c * a_k;
    b = b * ((k+N) * (k-N)) / ((k+1/2) * (k+1));
output s/d;
```

Note that this uses universal rational coefficients $c_k^{(N)}/d^{(N)}$ independent of the particular sequence (so they could be pre-computed...)

This algorithm gives relative accuracy $\sim 5.828^{-N}$, so to get D decimal digits, we need $N \sim 1.31D$ (so for 16 digits, need $N \sim 21$). (Assuming a_k are the moments of a positive measure on $[0, 1]$, equivalent to ...)

Can also use the partial sums $s_m = \sum_{k=0}^{m-1} (-)^k a_k$ (rather than the individual terms) and compute

$$\sum_{k=0}^N c_k^{(N)} a_k = \sum_{m=1}^N \frac{N}{n+m} \binom{N+m}{2m} 2^{2m} s_m$$

22.4 Root finding

22.4.1 Newton's method

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

22.4.2 Halley's method

$$x_{n+1} = x_n - \frac{2f(x_n)f'(x_n)}{2f'(x_n)^2 - f(x_n)f''(x_n)}$$

23 Notes on implementations

tolerance, accuracy, error estimate (max, expected), iterations, time, cost (mul,div,plus,sub,exp,... ops)

- Taylor series
 - at fixed points
 - at arbitrary points
 - analytic continuation
 - (modified series)
- (For inverse functions)
 - root-finders
 - direct approximation / inverse interpolation
 - power series (Lagrange inversion, Dominici techniques)
- quadrature
 - real
 - complex (contour)
 - ODE (system)
- local approximation
 - polynomial (power-series, Chebyshev, other orthogonal polynomial)
 - Padé (rational)
 - other series
- continued fraction
 - fixed depth, direct
 - iteration (P,Q)
- asymptotic expansion
- recurrence (backward, forward, scaling trick)
- expansion in other special functions

24 Notes on testing

1. tabulated values (Mathematica, Maple, etc.) [$\operatorname{erf} 0 = 0$]
2. comparison for special cases (relations between functions) [$\operatorname{erf} z = \gamma(z^2, 1/2)$]
3. identities [$\cos^2 z + \sin^2 z = 1$]
4. comparison among alternative implementations [power series vs. Padé approximation vs. continued fraction vs. integration]
5. properties of function (e.g. check monotonicity — strictest test is 1ulp bump...; inequalities, etc.)
6. for functions with many parameters, use a lattice (say Sobol' qrng) to generate points that uniformly sample the space for testing specific values

25 General mathematical library

1. Data-types
 - (a) double
 - (b) double-double, quad-double
 - (c) arbitrary precision
 - (d) complex_i
 - (e) vector_i, matrix_i, matrix_ndim_i
 - (f) Common Lisp has: Number=Complex/Real; Real=Float/Rational; Float=Short/Single/Double/Long; Rational=Ratio/Integer; Integer=Fixnum/Bignum
 - (g) Scheme (R6RS) has: Complex - Real - Rational - Integer and distinguishes Exact/Inexact and Flonum/Fixnum
2. Monte-carlo methods
 - (a) Random-number generation (Mersenne Twister prng, Sobol' qrng)
 - (b) Distribution sampling (all major types as well as generic; discrete and continuous)
3. Root-finding
 - (a) 1-dimension: bracketer, bisection, secant, Newton, inverse-quadratic, Halley, Brent-Dekker, (monotonic)
 - (b) 2-,*n*-dimension: conjugate-gradient, homotopy
 - (c) polynomial methods — specialized methods to find *all* real/complex roots, etc.
4. Sequence acceleration (Aitken, Wynn, Δ , etc.)
5. Differentiation (basic, *n*-th order, Savitzky-Golay, adaptive; weight generators)
6. Optimization
 - (a) 1-dim minimization / maximization
 - (b) *n*-dim minimization / maximization
 - (c) least-squares optimization (linear / non-linear)
7. Integration
 - (a) Quadrature (1-dim)
 - i. Open/closed/infinite intervals, singularities
 - ii. Trapezoidal, Simpson, Gauss-type rules (& weight generators)
 - iii. Contour integration in \mathbb{C} (branch tracking?)
 - (b) Cubature (2-dim)
 - (c) High-dimensional integration (lattice, MC, quasi-MC; dimension reduction)
 - (d) ODE
 - i. 1-dim second-order equation
 - ii. *n*-dim first-order (RK (4 & others), adaptive, Boer-Stoerlisch, etc.)
 - (e) PDE (generic IC, BC, coeffs)
 - i. 1-dim parabolic
 - ii. 2-dim parabolic
 - iii. 2-,3-dim elliptic

8. Solving integral equations ...

9. Interpolation

- (a) 1-dim (splines (cubic, B-, Akima, general), Chebyshev, RBF, etc.)
- (b) 2-dim (bicubic, scattered data, etc.)
- (c) n -dim (scattered, etc.)
- (d) probability distribution

10. Linear algebra

(a) ...

- function grouping / classification
- name, symbol
- parameters, arguments: largest domain of definition, branch cuts, etc.
- different representations (and domains of validity)
- relations
- implementation notes — for different parameter/argument restrictions (e.g. \mathbb{N} vs \mathbb{R} vs \mathbb{C}) — for scaled versions, etc.
- NR/ISML/Boost/GSL/C library/NAG/Pari/cephes/&c. availability

Generic functions templated with input-type, working-type, output-type (which can be allowed to be real or complex variants). Can be useful for balancing needed accuracy vs. speed (especially inside other routines...)

Precision: [also 20/25/30 digits for double-double, etc.; arbitrary precision? (template implementation...)]

- Green: ≥ 15 digits
- Yellow: ≥ 10 digits
- Red: ≥ 5 digits
- Grey: < 5 digits and/or restricted domain
- Black: not supported

Timing:

- $\leq 10\text{ms}$
- $\leq 100\text{ms}$
- $\leq 1\text{s}$
- $> 1\text{s}$

Domain:

- integer
- real interval
- reals
- complex domain

- complex
- principal value + branches

Platform:

- Octave
- C
- C++

Modify with $+/-$ for some discrepancies (lower in extreme ranges, etc.)

Also, intrinsic limitations (over/under-flow) [solve with scaled versions also] vs. extrinsic (unstable algorithm, etc.)

Idea: number representation as Integer \pm Fractional-part, where both of the two components have an independent exponent/mantissa representation. (Similar to the quad/double-double representations in two parts...) This allows to represent $1 + \varepsilon$ with high precision in ε ...

Estimation of cancellation errors in summations — compare magnitude of result to largest term / partial-sum.

Test data-files: # comment or nnn—nnn—nnn (to many digits of accuracy, minimum 72 (for quad-double+guards)).

Plot test points — color-coded for error/etc. (for 2 parameter cases).

References

- [A&S] Abramowitz and Stegun, *Handbook of*
- [Concrete] R.L. Graham, D.E. Knuth, and O. Patashnik, *Concrete mathematics*, Addison-Wesley, Reading, MA, 1989.
- [Luke1] Yudell Luke, *The Special Functions and Their Approximations, Volume 1*.
- [Atlas] K. Oldham, J. Myland, and J. Spanier, *An Atlas of Special Functions: With Equator, the Atlas Function Calculator*, second edition Springer, 2009.
- [NIST] eds F.W.J. Olver, D.W. Lozier, R.F. Boisvert, and C.W. Clark, *NIST Handbook of Mathematical Functions*, NIST and Cambridge University Press, 2010.
- [NR] W.H. Press, S.A. Teukolsky, W.T. Vetterling, and B.P. Flannery, *Numerical Recipes: The art of scientific computing*, third edition, Cambridge University Press, 2007.
- [ACMF] W.J. Thompson, *Atlas for Computing Mathematical Functions: An Illustrated Guide for Practitioners; With Programs in Fortran 90 and Mathematica*, John Wiley & Sons, Inc., 1997.
- [ZJ] S. Zhang and J. Jin, *Computation of Special Functions*, John Wiley & Sons, Inc., 1996.
- [DLMF] *DLMF*
- [Lebedev] Lebedev *Special Functions...*
- [Marsaglia] G. Marsaglia, “Evaluating the Normal Distribution” *Journal of Statistical Software*, July 2004, Volume 11, Issue 4.
- [Precise] M. Sofroniou and G. Spaletta, “Precise Numerical Computation”, in
- [AppFD] N.M. Temme and A.B. Olde Daalhuis, “Approximation of Fermi-Dirac integrals” in *Journal of Computational and Applied Mathematics*, **31** (1990) 383–387, North-Holland.
- [ZetaE] Edwards ... *Riemann Zeta Function*.