Seth Ricks ECEN 470 PS03: 3.45(a)(b)(c), 3.46(a)(b)(Hint: see c) 3.53(a), 3.55

1/6

(3.45) Find TF below:

a.) Which meet an overshoot of $M_p \leq 5\%$?
b.) " " a rise time of $t_r \leq 0.5$sec?
c.) " " a setting time of $t_s \leq 2.5$sec?

$$G_1(s) = \frac{40}{s^2+4s+40}$$
$$G_2(s) = \frac{40}{(s+1)(s^2+4s+40)}$$
$$G_3(s) = \frac{120}{(s+3)(s^2+4s+40)}$$
$$G_4(s) = \frac{20(s+2)}{(s+1)(s^2+4s+40)}$$
$$G_5(s) = \frac{36040/401\,(s^2+s+401)}{(s^2+4s+40)(s^2+s+401)}$$

$$H(s) = \frac{\omega_n^2}{s^2+2\zeta\omega_n s+\omega_n^2}$$

$G_1(s)$: $4s = 2\zeta\omega_n s \rightarrow 2 = \zeta\omega_n$

$40 = \omega_n^2$

$\omega_n = \sqrt{40} = 2\sqrt{10}$

$2 = \zeta(2\sqrt{10})$

$\zeta = \frac{1}{\sqrt{10}}$

$t_r = \frac{1.8}{\omega_n} = \frac{1.8}{2\sqrt{10}} = 0.285$ ✓

$t_s = \frac{4.6}{\zeta\omega_n} = \frac{4.6}{(\frac{1}{\sqrt{10}})(2\sqrt{10})} = 2.3$ ✓

$M_p = e^{-\pi\zeta/\sqrt{1-\zeta^2}}$

$= e^{-\pi/\sqrt{10}/\sqrt{1-(1/\sqrt{10})^2}}$

$= 0.35 \rightarrow 35\%$ X

$M_p \leq 5\%$? NO
$t_r \leq 0.5$ sec? Yes
$t_s \leq 2.5$ sec? Yes

2/6

$G_2(s) = \frac{40}{(s+1)(s^2+4s+40)}$ Poles: $s = -2 \pm j6$, $s = -1$

* within factor of 4 → significant effect *

For $G_2(s)$, since pole is within factor of 4 of the real part of the original poles, overshoot will decrease, rise time will increase, and settling time will increase.

$M_p \downarrow \leq 5\%$? → probably ✓
$t_r \uparrow \leq 0.5$ sec? → probably not ×
$t_s \uparrow \leq 2.5$ sec? → probably not ×

$G_3(s) = \frac{120}{(s+3)(s^2+4s+40)}$

Original: $6 = -2$
new: $s = -3$
$Re\{p\} < 46$: yes
-3 further left than -2 → faster
$M_p \downarrow \leq 5\%$? → probably ✓
$t_r \uparrow \leq 0.5$ sec? → probably not ×
$t_s \uparrow \leq 2.5$ sec? → probably not ×

$G_4(s) = \frac{20(s+2)}{(s+1)(s^2+4s+40)}$

Original: $6 = -2$ closer to origin
new p: $s = -1$
new z: $s = -2$
} → within factor of 4 = yes
$M_p \downarrow$ → ✓
$t_r \uparrow$ → ×
$t_s \uparrow$ → ×

$G_5(s) = \frac{36040/401\,(s^2+s+4011)}{(s^2+4s+40)(s^2+s+901)}$ Original: $6 = -2$
New poles: $s = -0.5 \pm j30$
New zeros: $z = -0.5 \pm j20$

both -0.5, balance each other out.
M_p ×
t_r ✓ t_s ×

After MATLAB:

3/6

	$M_p \leq 5\%$	$t_r \leq 0.5$ sec	$t_s \leq 2.5$ sec
G_1	×	✓	✓
G_2	✓	×	×
G_3	✓	✓	✓
G_4	✓	×	×
G_5	×	✓	✓
	↑ a.)	↑ b.)	↑ c.)

3.46 Consider following two nonminimum phase systems.

$$G_1(s) = -\frac{2(s-1)}{(s+1)(s+2)} \qquad G_2(s) = \frac{3(s-1)(s-2)}{(s+1)(s+2)(s+3)}$$

a.) Sketch unit step response for $G_1(s)$ & $G_2(s)$, paying close attention to the transient part of response
b.) Explain difference in behavior of two responses as it relates to the zeros

---

* step: $\mathcal{L}\{u(t)\} = \frac{1}{s}$


$$G_1(s)\cdot\frac{1}{s} = \frac{-2(s-1)}{s(s+1)(s+2)} = \frac{1}{s} - \frac{4}{s+1} + \frac{3}{s+2}$$

$$g_1(t) = [1 - 4e^{-t} + 3e^{-2t}]u(t)$$

$$G_2(s)\cdot\frac{1}{s} = \frac{3(s-1)(s-2)}{s(s+1)(s+2)(s+3)} = \frac{1}{s} - \frac{9}{s+1} + \frac{18}{s+2} - \frac{10}{s+3}$$

$$g_2(t) = [1 - 9e^{-t} + 18e^{-2t} - 10e^{-3t}]u(t)$$

a.)



b.) G_1 = odd # of zeros = undershoot
G_2 = even # of zeros = overshoot

3.53 Use Routh's stability criterion to determine how many roots w/ + real parts.

a.) $s^4 + 8s^3 + 32s^2 + 80s + 100 = 0$

Routh array:

s^4: 1 32 100

s^3: 8 80

s^2: $\frac{(8)(32)-(1)(80)}{8} = 22$ $\frac{8(100)-(1)(0)}{8} = 100$

s^1: $\frac{(22)(80)-8(100)}{22} = 43.63$

s^0: 100

No poles in RHP

0 roots with pos real parts

3.55 TF: $KG(s) = \frac{K(s+4)}{s[(s+0.5)(s+1)(s^2+0.4s+4)]}$ 6/6

Using Routh's stability criterion, determine range of K for which system is stable when $1+KG=0$.

Char eq: $1+KG(s)=0$

$$1+\frac{K(s+4)}{s[(s+0.5)(s+1)(s^2+0.4s+4)]}=0$$

$$s[(s+0.5)(s+1)(s^2+0.4s+4)]+K(s+4)=0$$

$$s^5+1.9s^4+5.1s^3+6.2s^2+2s+Ks+4K=0$$

$$s^5+1.9s^4+5.1s^3+6.2s^2+(2+K)s+4K=0$$

Routh array:

s^5:	1	5.1	$(2+K)$	← (odd powers)
s^4:	1.9	6.2	$4K$	← (even powers)
s^3:	1.837	$2-1.105K$		
s^2:	$4.13+1.14K$	$4K$		
s^1:	$\frac{-1.26K^2-9.63K+8.26}{4.13+1.14K}$			
s^0:	$4K$			

$\frac{1.9(2+K)-4K}{1.9} = 2-1.105K$

$\frac{(1.837)(6.2)-1.9(2-1.105K)}{1.837}$

$\frac{11.389-3.8+2.099K}{1.837}$

$4.131+1.143K$

$\frac{1.837(4K)-1.9(0)}{1.837}=4K$

$\frac{(4.13+1.14K)(2-1.105K)+(1.837)(4K)}{4.13+1.14K}$

$\frac{-1.26K^2-9.63K+8.26}{4.13+1.14K}$

every element in 1st column must be >0

s^0: $K>0$
s^1: $K<0.78$
s^2: $K>-3.62$
s^3: $K<1.81$

$\boxed{0<K<0.78}$