

1) Lesson 1

1.1) Inequalities

In general:

1. Find domain D
2. Rewrite so that it is function bigger/smaller than 0
3. Divide in different products and analyze them individually
 1. find zeroes
 2. study sign
4. Final sign analysis

From bolzano weierstrass theorem (you will see this later) we have that every intersection at 0, that is, at every zero of the function. there is a change of sign.

Ex c)

Solve the following inequality:

$$\frac{x-1}{x-2} > \frac{2x-3}{x-3}$$

First find the domain:

$$D = x \in \mathbb{R} \setminus \{2, 3\}$$

Now compute

$$\frac{x-1}{x-2} - \frac{2x-3}{x-3} > 0 \implies \frac{(x-1)(x-3) - (2x-3)(x-2)}{(x-2)(x-3)} > 0$$

This is > 0 if numerator and denominator share the same sign, that is, where the function is positive.

Numerator:

First we find the zeroes of the numerator function:

$$\begin{aligned}(x-1)(x-3) - (2x-3)(x-2) &= (x^2 - 4x + 3) - (2x^2 - 7x + 6) \\ &= -x^2 + 3x - 3\end{aligned}$$

Recall the quadratic formula:

Let $a, b, c \in \mathbb{R}$, then $ax^2 + bx + c = 0$ has the following solution:

$$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$


Where the discriminant $\Delta = b^2 - 4ac$ determines the types of solution:

- $\Delta > 0$: two real (distinct) solutions
- $\Delta = 0$: two real (identical) solutions
- $\Delta < 0$: no real solution

In this case since $\Delta = 9 - 12 = -3 < 0$ we don't have any solutions, hence since $a < 0$ the whole numerator is negative.

Denominator:

Here the function already shows the zeroes, so we can plot the graph to see the regions:



Curve

So we have

$$D > 0 \iff x < 2 \cup x > 3$$

By noticing the interval where they have the same sign:

x	$(-\infty, 2)$	$(2, 3)$	$(3, +\infty)$
Numerator	-	-	-
Denominator	+	-	+
Fraction	-	+	-

Finally we have:

$$\boxed{2 < x < 3}$$

Consider the following:

1.2) Subsets of $\mathbb{R}$

Ex a)

Describe the following set:

$$A = S_1 \cap S_2 = \{x \in \mathbb{R} : x^2 + 4x + 13 < 0\} \cap \{x \in \mathbb{R} : 3x^2 + 5 > 0\}$$

Subset 1:

$$x^2 + 4x + 13 < 0 \rightarrow \Delta = -36 \implies S_1 = \emptyset$$

any intersection with an empty set is an empty set:

$$\boxed{A = \emptyset \cap S_2 = \emptyset}$$

In general recall:

$$\emptyset \cap S = \emptyset \quad \emptyset \cup S = S$$

Ex c)

Describe the following set:

$$C = \left\{x \in \mathbb{R} : \frac{x^2 - 5x + 4}{x^2 - 9} < 0\right\} \cup \left\{x \in \mathbb{R} : x + \sqrt{7x + 1} = 17\right\}$$

Subset 1:

Numerator:

$$x^2 - 5x + 4 = 0 \rightarrow x_{1,2} = 1, 4 \rightarrow N > 0 \iff x < 1 \cup x > 4$$

Denominator:

$$x^2 - 9 = (x + 3)(x - 3) = 0 \rightarrow D > 0 \iff x < -3 \cup x > 3$$

x	$(-\infty, -3)$	$(-3, 1)$	$(1, 3)$	$(3, 4)$	$(4, +\infty)$
Numerator	+	+	-	-	+
Denominator	+	-	-	+	+
Fraction	+	-	+	-	+

so $S_1 = (-3, 1) \cup (3, 4)$

Subset 2:

This is just an equality:

$$\sqrt{7x+1} = 17-x$$

The domain is $7x+1 \geq 0 \rightarrow D = \{x \leq 17\}$

Square both sides:

$$7x+1 = (17-x)^2 \rightarrow 7x+1 = x^2 - 34x + 17^2 \rightarrow x^2 - 41x + 288 = 0$$

$$x_{1,2} = 9, 32$$

But only 9 is acceptable as $32 \notin D$

Finally:

$$C = (-3, 1) \cup (3, 4) \cup \{9\}$$

1.3) Max/Min, Sup/Inf

Recall the theory

Definition (Upper/Lower Bound).

$M \in \mathbb{Q}$ is called **upper bound** for A if $M \geq a, \forall a \in A$.

Respectively it is called **lower bound** if $M \leq a, \forall a \in A$

Notice M does not need to be in A

Example: consider set $A = [0, 1)$ any $m \leq 0$ is a lower bound and any $M \geq 1$ is upper bound

Definition (Maximum/Minimum of A).

If $\overline{M}$ is an upper bound for A and $\overline{M} \in A$, if $\overline{M}$ is the *minimum* possible *upper* bound, then it is the **maximum** of A .

If $\overline{m}$ is a lower bound for A and $\overline{m} \in A$, if $\overline{m}$ is the *maximum* possible *lower* bound, then it is the **minimum** of A .

In this case $\overline{m} \in A$ by definition.

From the previous example: $\overline{m} = \max \{m \leq 0\} = 0 \in A$ is the correct minimum of A , however

$\overline{M} = \min \{M \geq 1\} = 1 \notin A$ means that we don't have a maximum

Definition (Supremum/Infimum).

The **supremum** of A $\sup A$ is the minimum of the upper bounds of A . If a set allows for a maximum, then it is also the supremum

The **infimum** of A $\inf A$ is the maximum of the lower bounds of A . If a set allows for a minimum, then it is also the infimum.

Finish the example: we know $\overline{m} = 0 \in A$ so this is both minimum and infimum of A . We know also $\overline{M} = 1 \notin A$, therefore $\sup A = 1$ but it is not the maximum.

Ex a)

Tell whether this subset is bounded and the sup/inf and if it allows for maximum/minimum.

$$A = \left\{ x \in \mathbb{R} : x = n \text{ or } x = \frac{1}{n^2}, n \in \mathbb{N} \setminus \{0\} \right\}$$

It is immediately clear that A is the union of two sets, mainly

$$A = S_1 \cup S_2 \text{ where } \begin{aligned} S_1 &= \{n \in \mathbb{N} \setminus \{0\}\} = \{1, 2, 3, 4, \dots\} \\ S_2 &= \left\{ \frac{1}{n^2} \mid n \in \mathbb{N} \setminus \{0\} \right\} = \left\{ 1, \frac{1}{4}, \frac{1}{9}, \frac{1}{16}, \dots \right\} \end{aligned}$$

Find the bounds on both sets individually (not required but the result is clearer).

Study the first subset S_1 : notice that it is just the natural numbers with zero excluded, so its properties are already known.

- No upper bound, so also no max or sup
- Lower bound: $m_1 \leq 1$
 - $\overline{m}_1 = \max m_1 \leq 1 = 1 \in S$ which is also the infimum!

Study the second subset S_2 : Intuitively we know that the function describing the set is monotonically decreasing: we can express it by saying:

Call x_n the value of x for any chosen n , then $x_{n+1} < x_n$.

This is easily shown as the following inequality is true for $n \in \mathbb{N} \setminus \{0\}$ (check at home):

$$\frac{1}{(n+1)^2} < \frac{1}{n^2}$$

Therefore for every number $x_n > 0$ we can find a smaller number such that $0 < x_{n+1} < x_n$

With this digression aside, we also know that $0 \notin S_2$.

As before analyze:

- Lower bound: $m_2 \leq 0$
 - No minimum: $\overline{m}_2 = \max m_2 \leq 0 = 0 \notin S_2$
 - Infimum is $\inf S_2 = 0$
- Upper bound: $M_2 \geq 1$
 - $\overline{M}_2 = \min M_2 \geq 1 = 1 \in S_2$ which is also supremum!

Putting all together:

Now consider the union, so:

- No upper bound since S_1 is unbounded
 - therefore no max and sup
- Lower bound: the minimum of the two lower bounds is $m \leq 0$
 - no minimum since $0 \notin A$
 - Infimum is $\inf A = 0$

2) Symbols

- $\implies$: implies, necessary condition
- $\impliedby$: implied by, sufficient condition
- $\iff$: necessary + sufficient, if and only if "iff"
- $\emptyset$: empty set
- $\mathbb{N}$: natural numbers; $\{0, 1, 2, \dots\}$
- $\mathbb{Z}$: integer numbers; $\{\dots, -2, -1, 0, 1, 2, \dots\}$
- $\mathbb{Q}$: rational numbers; $\{\frac{m}{n} \mid m, n \in \mathbb{Z}, n \neq 0\}$
- $\mathbb{R}$: real numbers
- $\mathbb{C}$: complex numbers
- $\cup$: Union; or
- $\cap$: Intersection; and