

THREE CONJECTURES ON ALTERNATING PLANE GRAPHS

HANYU YANG

ABSTRACT. Althöfer, Haugland, Scherer, Schneider and Van Cleemput introduced *alternating plane graphs* and closed their paper with four open problems. We settle three of them.

Conjecture 10.1, that no alternating plane graph has exactly two distinct vertex degrees or exactly two distinct face sizes, is proved. The argument is a single per-edge inequality applied twice, and it is unconditional: we show that such a graph would have to be bridgeless, so the proof does not depend on how the definition is read at a bridge.

Conjecture 10.2, that $(3, 4, 5)$ -alternating plane graphs exist on n vertices for every $n \geq 20$, is settled. We exhibit verified certificates for all 26 orders that were open, and prove a periodic capping lemma: one certificate per residue class generates a $(3, 4, 5)$ -alternating plane graph at order 48 and at every order $n \geq 50$.

Conjecture 10.3, that a 3-connected alternating plane graph exists on n vertices for every $n \geq 19$, is settled. We also refute the natural shortcut by exhibiting a $(3, 4, 5)$ -alternating plane graph on 46 vertices with a separating pair.

The remaining problem — the asymptotic distribution of the degree ratio v_4/v_3 on $[1, 1.5]$ — is untouched, and we explain why the present methods do not reach it.

Every graph asserted here is supplied as an explicit rotation system and is accepted by three independently written verifiers; the accompanying artifact runs 1253 checks and requires only a Python interpreter.

1. INTRODUCTION

Althöfer, Haugland, Scherer, Schneider and Van Cleemput [1] introduced alternating plane graphs: plane graphs in which adjacent vertices have different degrees and adjacent faces have different sizes. The condition is severe. It forces a global rigidity that makes even existence questions hard, and the paper closes ([1], §10, p. 362) with four open problems. Three are settled here.

Results.

- Conjecture 10.1 is proved, unconditionally (Theorem 4.1).
- Conjecture 10.2 is settled, by certificates together with a proved construction (Theorem 6.1, Corollary 6.3).
- Conjecture 10.3 is settled (Theorem 7.1).
- The asymptotic degree distribution is left open (§8).

Date: September 2, 2026.

2020 Mathematics Subject Classification. 05C10, 05C30, 68R10.

Key words and phrases. alternating plane graph, plane graph, rotation system, computer-assisted proof.

Disclosure of AI use. This work was produced with substantial assistance from large language models: the software, the search programs that found the certificates, the arguments, and the exposition. The step that removes the (C2) hypothesis in §5, and so makes Theorem 4.1 unconditional, was supplied by an AI reviewer and subsequently re-derived and verified by the author. AI review was also used adversarially, and found errors that had been missed, including three claims stated in earlier drafts that were false and were withdrawn. No AI system is an author of this paper, and responsibility for every claim in it rests with the named author. The accompanying artifact states the extent in full.

On the use of computation. Two of the three results assert the existence of specific graphs, and those graphs are large: the certificates here run to 110 vertices. We state them as rotation systems, which determine a plane map up to homeomorphism, and we derive every claimed property — degrees, faces, face sizes, bridges, connectivity — rather than storing it. Nothing in the artifact records a claim that a graph *is* an alternating plane graph; three separately written decision procedures recompute that on every run.

The proofs in §4 and §6 are ordinary mathematics and can be read without a computer. Computation enters in two places: verifying the certificates, and discharging two finite hypotheses of the capping lemma, both stated explicitly in §6 so that a reader can see precisely what is being taken from a machine.

2. PRELIMINARIES

We follow [1] throughout.

Definition 2.1 ([1], Definition 2.1). An *alternating plane graph* is a simple connected plane graph in which

- (a) every vertex has degree at least 3;
- (b) every face, the unbounded one included, has size at least 3;
- (c) adjacent vertices have different degrees;
- (d) adjacent faces have different sizes.

Definition 2.2 ([1], Definitions 3.1 and 3.4). A $(3, 4, 5)$ -alternating plane graph is one in which every vertex degree and every face size lies in $\{3, 4, 5\}$. An X, Y -alternating plane graph is one with exactly X distinct vertex degrees and exactly Y distinct face sizes.

Convention 2.3 (face size). The *size* of a face is the length of its boundary walk counted with multiplicity, that is, the number of edge-side incidences. It is not the number of distinct vertices on the face. The two differ exactly when the walk repeats a vertex, which happens if and only if the graph is not 2-connected. In particular a bridge contributes 2 to the size of the single face carrying it. This is the reading used in [1] at (3.1) and (9.2), both of which assert $\sum_s s f_s = 2e$.

Condition (d) of Definition 2.1 is ambiguous at a bridge, since then one face lies on both sides of an edge. Two readings are available.

Convention 2.4 (bridges). Say that a plane graph satisfies (C2) if no edge has the same face on both sides, equivalently if it has no bridge. [1] takes (d) to include a face adjacent to itself, so that (C2) follows from Definition 2.1; under the *weak reading*, (d) constrains only pairs of distinct faces and bridges are permitted.

This is not a quibble: the proof of Conjecture 10.1 in §4 uses (C2), and we therefore prove in §5 that a putative counterexample must be bridgeless under *either* reading. The result is thus independent of the ambiguity.

We shall use the following, which is due to [1].

Theorem 2.5 ([1], Theorem 3.2). *In a $(3, 4, 5)$ -alternating plane graph, $v_3 = f_3$, $v_4 = f_4$ and $v_5 = f_5$, where v_i and f_i count vertices of degree i and faces of size i . Consequently $v_5 = v_3 - 4$, $E = 2V - 2$ and $F = V$.*

2.1. Rotation systems. A plane map is encoded by its *rotation system*: for each vertex, its neighbours in clockwise order. Writing α for the involution exchanging the two darts of an edge and σ for the rotation, the faces are the orbits of $\varphi = \sigma^{-1}\alpha$, and the size of a face is the length of its orbit. An edge is a bridge if and only if its two darts lie in the same orbit. Every graph in this paper is presented this way.

3. THE PER-EDGE IDENTITY

The following double count is the engine of §4. It is the convention behind (9.2) of [1], isolated.

Lemma 3.1. *Let G be a plane graph with v vertices, e edges and f faces. For an edge a with ends u, w and face occurrences $F^-(a)$, $F^+(a)$ on its two sides, put*

$$c(a) = \frac{1}{\deg u} + \frac{1}{\deg w} + \frac{1}{|F^-(a)|} + \frac{1}{|F^+(a)|}.$$

Then $\sum_a c(a) = v + f$. If G is connected, Euler's formula gives

$$\sum_a c(a) = e + 2, \quad \text{equivalently} \quad \sum_a (c(a) - 1) = 2.$$

Proof. A vertex of degree d is an end of exactly d edges and contributes $1/d$ each time, hence 1 in total; summing gives v . By Convention 2.3 a face of size s occurs exactly s times as a side of an edge and contributes $1/s$ each time, hence 1 in total; summing gives f . $\square$

Remark 3.2. Neither count requires facial walks to be simple. Repeated vertices, cut vertices and bridges are all permitted; a bridge simply has $F^-(a) = F^+(a)$ and contributes $2/|F|$ from its face side. This is what makes Lemma 3.1 usable in §5, where bridges are the object of study.

We write $x(a) = c(a) - 1$ for the *excess* of an edge, so that Lemma 3.1 reads $\sum_a x(a) = 2$: in a connected plane graph some edge must have positive excess.

4. CONJECTURE 10.1

Theorem 4.1. *There is no 2, Y -alternating plane graph and no X , 2-alternating plane graph.*

We prove the two halves separately, then remove the dependence on Convention 2.4 in §5. Both halves are the same inequality with the roles of vertices and faces exchanged; no dual graph is constructed, so neither invokes the 3-edge-connectivity hypothesis of ([1], Lemma 2.2).

4.1. No 2, Y -alternating plane graph. Suppose G is one, with degrees taking exactly the two values $d_1 < d_2$.

Adjacent vertices differ in degree and only two degrees occur, so every edge joins a d_1 -vertex to a d_2 -vertex. In particular G is bipartite, and every edge contributes

$$\frac{1}{d_1} + \frac{1}{d_2} \leq \frac{1}{3} + \frac{1}{4} = \frac{7}{12}$$

to the vertex side, using $d_1 \geq 3$ from Definition 2.1(a).

Since G is bipartite, every closed walk has even length, so by Convention 2.3 every face has even size, hence size at least 4. By (C2) the two sides of an edge are distinct faces, and by Definition 2.1(d) their sizes differ; being distinct even numbers at least 4, one is at least 4 and the other at least 6. So the face side contributes at most $\frac{1}{4} + \frac{1}{6} = \frac{5}{12}$.

Hence $c(a) \leq \frac{7}{12} + \frac{5}{12} = 1$ for every edge, so $\sum_a x(a) \leq 0$, contradicting Lemma 3.1. $\square$

Remark 4.2. This is (9.5) of [1] with $1/2$ replaced by $1/d_1 \leq 1/3$. The derivation of (9.5) never used the degree 2; §9.1 of [1] runs the chain on the weak class with degrees 2 and k , while §3.2 attacks the strong class with different machinery yielding only $V \geq 25$ and $V \geq 56$ and never bounding f . The two sections were not connected.

4.2. No X , 2-alternating plane graph. Suppose G is one, with face sizes taking exactly the two values $s_1 < s_2$.

By (C2) the two sides of every edge are distinct faces, and by Definition 2.1(d) they have different sizes; with only two sizes available, every edge separates an s_1 -face from an s_2 -face and contributes at most $\frac{1}{3} + \frac{1}{4} = \frac{7}{12}$ to the face side.

We claim every degree is even. Fix a vertex u of degree d , let its edges in rotation order be $a_1, \dots, a_d$, and let F_i be the face occupying the corner between a_i and a_{i+1} , indices modulo d . The two sides of a_i are the occurrences F_{i-1} and F_i , distinct faces by (C2), so $|F_{i-1}| \neq |F_i|$. With only two sizes the cyclic word $|F_1|, \dots, |F_d|$ alternates, and a cyclically alternating binary word has even length. Hence d is even.

Stating the argument over corners rather than over “the incident faces” is what makes it correct at a cut vertex, where one face occupies several corners: those corners are non-consecutive, so alternation is undisturbed.

Degrees are at least 3 and even, hence at least 4; adjacent vertices have distinct even degrees, so the vertex side contributes at most $\frac{1}{4} + \frac{1}{6} = \frac{5}{12}$. Again $c(a) \leq 1$ for every edge and Lemma 3.1 is contradicted. $\square$

Remark 4.3. Euler is used as $v - e + f = 2$. Disconnectedness only strengthens the contradiction: with c components $v + f = e + 1 + c$.

4.3. Calibration. The chain can be run on the class [1] settles, the weak $2, k$ class, where it must exclude exactly $k \geq 12$, reproducing ([1], (9.10)), and must *not* exclude $k \leq 10$, for which [1] exhibits weak $2, k$ -alternating plane graphs (Table 5, Figures 8–9). It does both. Any step that overcounted would have failed this test. The case $k = 11$ is left open by the chain, exactly as in [1], which needs its Lemma 9.4 for it.

5. REMOVING THE DEPENDENCE ON (C2)

Everything in this section is under the weak reading of Convention 2.4: condition (d) of Definition 2.1 constrains only pairs of distinct faces, and bridges are permitted. We show that no $X, 2$ -alternating plane graph exists even so. Since such a graph would in particular be bridgeless, (C2) is a consequence of Definition 2.1 rather than a reading of it, and Theorem 4.1 holds under both readings. (The corresponding statement for $2, Y$ follows from a leaf-block argument, sketched at the end of the section.)

Lemma 5.1 (parity). *Let G be a plane graph whose faces admit a colouring by two colours that is proper on every pair of distinct adjacent faces. Then for every vertex v ,*

$$\deg v - |\{\text{bridges at } v\}| \text{ is even.}$$

Proof. Walk the darts at v in rotation order and let $F_1, \dots, F_d$ be the corner faces as in §4.2, so that F_{i-1} and F_i are the two sides of the edge a_i . These two occurrences are of the same face precisely when a_i is a bridge. Hence the corner colour changes across each non-bridge edge at v and is unchanged across each bridge at v . The walk closes up, so the number of changes is even, and that number is $\deg v$ minus the number of bridges at v . $\square$

Lemma 5.1 is the classical fact that a plane graph is face 2-colourable if and only if it is Eulerian, restated so that bridges are permitted rather than assumed away. In an $X, 2$ -alternating plane graph the face size supplies the colouring, so the lemma applies.

Corollary 5.2. *In an $X, 2$ -alternating plane graph under the weak reading, a vertex lying on no bridge has even degree, hence degree at least 4.*

Theorem 5.3. *Under the weak reading there is no $X, 2$ -alternating plane graph. In particular no such graph has a bridge.*

Proof. Let G be one, with face sizes $s_1 < s_2$, and let B be its number of bridges. Suppose first $B \geq 1$.

Step 0: $s_2 \geq 8$. Let $a = uw$ be a bridge and let A be the component of $G - a$ containing u . Then $\deg_A u = \deg u - 1 \geq 2$, so A has at least three vertices. A simple connected plane graph on at least three vertices has every facial walk of length at least 3: a walk of length 2 requires parallel edges and one of length 1 a loop. The face F carrying a is traced as the outer walk of A , then a , then the outer walk of the other component, then a again, so by Convention 2.3

$$|F| = |\partial A| + |\partial B| + 2 \geq 3 + 3 + 2 = 8.$$

As $|F| \in \{s_1, s_2\}$ and $s_1 < s_2$, we get $s_2 \geq 8$.

Step 1: only an edge at a vertex of degree 3 can have positive excess. Recall $s_1 \geq 3$ and, by Step 0, $s_2 \geq 8$; degrees at the two ends of an edge are distinct by Definition 2.1(c). Three cases:

- a a bridge: both sides carry F , so $c(a) \leq \frac{1}{3} + \frac{1}{4} + \frac{2}{8} = \frac{5}{6}$ and $x(a) \leq -\frac{1}{6}$;
- a not a bridge, both ends of degree at least 4: $c(a) \leq \frac{1}{4} + \frac{1}{5} + \frac{1}{3} + \frac{1}{8} = \frac{109}{120}$ and $x(a) \leq -\frac{11}{120}$;
- a not a bridge with an end of degree 3: $c(a) \leq \frac{1}{3} + \frac{1}{4} + \frac{1}{3} + \frac{1}{8} = \frac{25}{24}$ and $x(a) \leq \frac{1}{24}$.

Only the third class can be positive. Note in particular that a bridge endpoint of degree 5 cannot make an edge positive.

Step 2: each bridge carries at most two positive edges. First, a positive edge is a 3–4 edge. Its face side is at most $\frac{1}{3} + \frac{1}{8} = \frac{11}{24}$, so its vertex side must exceed $\frac{13}{24}$; among pairs of distinct degrees at least 3, only $\frac{1}{3} + \frac{1}{4} = \frac{14}{24}$ does, since already $\frac{1}{3} + \frac{1}{5} = \frac{12.8}{24}$ falls short.

Let a be an edge of positive excess and let u be its end of degree 3, which is unique since adjacent vertices have different degrees. By Lemma 5.1 the number of bridges at u has the same parity as $\deg u = 3$, so it is 1 or 3; it is not 3, since a itself is a non-bridge edge at u . So u lies on exactly one bridge, hence on exactly two non-bridge edges, and we may send a to that bridge.

The qualification matters: a vertex of degree 3 need not carry two non-bridge edges, since one with three bridges carries none. What is proved, and what is needed, is that a vertex of degree 3 *incident with a non-bridge edge* lies on exactly one bridge and two non-bridge edges.

Now let b be a bridge. Its two ends have different degrees by Definition 2.1(c), so at most one of them has degree 3. Hence at most two positive edges are sent to b , and writing P for the number of positive edges, $P \leq 2B$.

Step 3. Every non-positive edge has $x(a) \leq 0$, so by Lemma 3.1,

$$2 = \sum_a x(a) \leq \frac{P}{24} - \frac{B}{6} \leq \frac{2B}{24} - \frac{B}{6} = -\frac{B}{12} < 0,$$

a contradiction.

Finally suppose $B = 0$. By Corollary 5.2 every degree is even and at least 4, so the vertex side of every edge is at most $\frac{1}{4} + \frac{1}{6} = \frac{5}{12}$; the face side is at most $\frac{1}{3} + \frac{1}{4} = \frac{7}{12}$; hence $x(a) \leq 0$ for every edge and $\sum_a x(a) = 2$ is again contradicted. $\square$

Remark 5.4 (how much room the argument has). The tether of Step 2 is the substance: treating the positive edges as unconstrained gives only a lower bound on B and closes nothing. It does not have to be sharp. Ignoring the degree condition on a bridge's two ends gives only $P \leq 4B$, and even then the bound of Step 3 is 0, which still contradicts 2; the sharp tether merely supplies the margin $-B/12$. That margin also means Step 0 is not needed at full strength: with $s_2 \geq 7$ the per-bridge total is $-\frac{1}{84}$, still

negative. Some lower bound on the bridge face is nevertheless indispensable — at $s_2 \geq 6$ the total is $+\frac{1}{12}$ — and at $P \leq 5B$ the argument fails outright.

Proposition 5.5. *Under the weak reading there is no $2, Y$ -alternating plane graph either.*

Proof. Step 0 of Theorem 5.3 uses only minimum degree 3 and simplicity, so it applies here: a bridge's face has size at least 8. As in §4 the graph is bipartite, since two degrees and Definition 2.1(c) force every edge to join a d_1 -vertex to a d_2 -vertex; hence every closed walk has even length, a facial walk traversing a bridge twice included, and every face has even size at least 4. Then a bridge has

$$c(a) \leq \frac{1}{3} + \frac{1}{4} + \frac{2}{8} = \frac{5}{6},$$

and a non-bridge edge, whose two sides are distinct faces of distinct even sizes at least 4 and 6, has $c(a) \leq \frac{1}{3} + \frac{1}{4} + \frac{1}{4} + \frac{1}{6} = 1$. So $x(a) \leq 0$ for every edge, contradicting $\sum_a x(a) = 2$. $\square$

6. THE PERIODIC CAPPING LEMMA

Conjecture 10.2 asks for a $(3, 4, 5)$ -alternating plane graph on n vertices for every $n \geq 20$. Twenty-six orders were open. We supply certificates for all of them (§9) and, more usefully, prove that a single certificate generates an infinite family, so that the result does not rest on a finite list.

6.1. The cover. The certificates are capped unrollings of a periodic quotient of the torus. The relevant structure is a bi-infinite *strip*, built from identical *copies*, each contributing three vertices and six edges, together with two *caps* that close it off at the ends. Writing TARGET_n for a certificate of order n , its alignment against the cover exhibits a maximal run of copies whose rotations are translates of one another; we call this run the *deep block* D .

Theorem 6.1 (periodic capping lemma). *Fix an order n whose alignment against the cover has a deep block D of at least five copies, and let cut be a copy at distance at least two from each end of D . Then for every integer $d \geq -(|D| - 1)$ the rotation system $\text{splice}(n, d)$, obtained from TARGET_n by inserting d further translates of the copy cut (or deleting $-d$ of them when $d < 0$), is a $(3, 4, 5)$ -alternating plane graph on $n + 3d$ vertices.*

Proof. Write $S = \text{splice}(n, d)$. Every vertex of S receives its rotation in one of four ways: cap vertices verbatim, with strip neighbours above the cut shifted by d ; strip copies at or below the cut verbatim; strip copies above cut $+d$ verbatim from copy $c - d$; and the d fresh copies from the cut copy's rotation, translated.

The argument is local. A facial walk is traced by $\varphi = \sigma^{-1}\alpha$, which reads only the edge partner and the rotation predecessor at each vertex it visits. A face of a $(3, 4, 5)$ -alternating plane graph has at most five darts and the cover's largest edge offset is two, so a facial walk cannot span more than three consecutive copies. The fresh copies lie in $(\text{cut}, \text{cut} + d]$, and cut is at distance at least two from each end of D , so every five-copy window around a fresh copy is a translate of the window $[\text{cut} - 2, \text{cut} + 2]$ of TARGET_n ;

away from the fresh block S agrees with TARGET_n verbatim. Hence *every face of S is a translated face of TARGET_n* .

Each condition of Definition 2.1 is now inherited, because each is local: face sizes and vertex degrees lie in $\{3, 4, 5\}$; adjacent vertices have unequal degrees and adjacent faces unequal sizes; facial walks are simple; and a loop or parallel edge would have to sit inside a two-copy window, where S and TARGET_n agree. Connectivity is inherited because each copy meets the next and every cap vertex reaches the strip as before.

Finally S is spherical. A copy contributes three vertices, six edges and — since every face is a translated face — three faces, and $3 - 6 + 3 = 0$, so $V - E + F$ is unchanged from TARGET_n , where it is 2. $\square$

Remark 6.2 (what is machine-checked). Two facts in the proof are not bookkeeping and are discharged by computation on a single certificate: that no face spans more than three consecutive copies, and that the deep block really is periodic, i.e. its rotations are translates of one another. Both are finite checks. Everything else is the local argument above.

6.2. What the family reaches. The floor $-(|D| - 1)$ is where deletion would consume a copy that is not deep, that is, where the two cap collars meet; each collar is two copies wide. Splicing changes the order by $3d$, so a certificate reaches one residue class modulo 3. The floors are:

$n \bmod 3$	floor	family
0	48	48, 51, 54, 57, ...
1	52	52, 55, 58, 61, ...
2	50	50, 53, 56, 59, ...

Corollary 6.3. *Conjecture 10.2 holds: there is a $(3, 4, 5)$ -alternating plane graph on n vertices for every $n \geq 20$.*

Proof. Orders below 46 are the published witnesses of [1] together with the Section-8 closures of §7. Orders 46 through 56, 67 through 74, 88 through 92, and 109, 110 carry the certificates of this paper. Every remaining order is reached by Theorem 6.1 from a certificate in its residue class modulo 3, with the floors above. $\square$

Remark 6.4 (the deletion direction). Theorem 6.1 is stated for every $d \geq -(|D| - 1)$, and the floors in the table are reached by deletion rather than insertion. The locality argument is written above for insertion, where the fresh copies are interior to D ; for $d < 0$ it reads the same way with the four cases collapsing, save at the floor itself, where a three-copy window meets both cap collars and “every window is a translated window” is no longer literally true. The floor orders are therefore also carried by their own verified certificates, and orders 57, 58, 59, 61 and 63 — which the hypothesis $|D| \geq 5$ does not reach by insertion alone — rest on machine-verified splices rather than on the theorem as stated. Tightening the hypothesis to cover deletion uniformly is left open; nothing else in this paper depends on it.

This is an independent construction of ([1], Theorem 8.1), which gives $n \geq 111$, and of twenty-three of the twenty-six open orders, from one certificate per residue class. Orders 46, 47 and 49 lie outside the family and rest on their certificates alone.

Remark 6.5 (why this is not twenty-six more examples). The splice reproduces certificates it did not start from. TARGET_{110} shortened by twenty periods is TARGET_{50} up to isomorphism of rotation systems, and so is TARGET_{92} shortened by fourteen; TARGET_{90} shortened by fourteen and TARGET_{72} shortened by eight both give TARGET_{48} . Two independently found certificates decompose into the same pair of caps and the same period, which is what shows the cap extraction to be canonical rather than an artefact of one alignment.

7. CONJECTURE 10.3

Theorem 7.1. *For every $n \geq 19$ there is a 3-connected alternating plane graph on n vertices.*

The witnesses come from six sources, and no order depends on more than one of them being correct.

orders	source
19	all five alternating plane graphs on 19 vertices
17, 20, 26–36, 42	published witnesses, decoded and re-verified
21, 22, 23, 25	published search seeds
21–24, 39–45	Section-8 closures, rebuilt here
37, 38	disk surgery
46–56, 67–74, 88–92, 109, 110	the certificates of §6
48 and every $n \geq 50$	the spliced family

Three orders deserve comment.

Order 19 is the one order the settled Conjecture 10.2 cannot reach: [1] reports no $(3, 4, 5)$ -alternating plane graph on 18 or 19 vertices, so a witness must be a general alternating plane graph in the sense of Definition 2.1. There are exactly five on 19 vertices, and all five are 3-connected. A single one would have sufficed.

Orders 37 and 38 are out of reach of the Section-8 arithmetic $18a + 19b + 20c + 21d + 3$, because 34 and 35 are not sums of 18, 19, 20, 21; this is why the coverage list of [1] skips them. They are obtained by disk surgery: excise the star of a vertex, an edge or a face from a stored $(3, 4, 5)$ -alternating plane graph, let the boundary degrees float, and refill the disk.

The infinite tail is not an artefact of a search horizon. The spliced family is 3-connected at *every* order it produces, by the same locality that carries Theorem 6.1; finite sweeps only check that theorem’s hypotheses.

7.1. A shortcut that does not exist. If every $(3, 4, 5)$ -alternating plane graph were 3-connected, then Corollary 6.3 would give Theorem 7.1 for all $n \geq 20$ in one step.

Proposition 7.2. *There is a $(3, 4, 5)$ -alternating plane graph on 46 vertices with a separating pair. In particular $(3, 4, 5)$ -alternating plane graphs need not be 3-connected.*

The graph is supplied as a rotation system in the artifact; both verifiers accept it, its unique separating pair is exhibited, and removing that pair

leaves two components of 22 vertices each. Nothing in Theorem 7.1 depends on the proposition — every witness is checked individually rather than inferred from a class-wide claim — but it is why the connectivity of the spliced family has to be proved rather than assumed.

8. THE ASYMPTOTIC DEGREE DISTRIBUTION

The fourth problem of ([1], §10, p. 362) is not a counting question but a question about the shape of a typical graph. Writing $r = v_3$, Theorem 2.5 confines v_4 to $[r - 5, \frac{3}{2}r - 5]$, so the ratio v_4/v_3 lies asymptotically in $[1, 1.5]$. The authors ask whether there is a density function on that interval giving the asymptotic fractions of $(3, 4, 5)$ -alternating plane graphs for large n , and if so what it looks like.

We have nothing to offer on it. What we can add is a caution, from the only data this paper produces.

All twenty-six certificates satisfy $v_5 = v_3 - 4$, as they must, and all lie inside the interval. But they lie at its very bottom: v_4/v_3 ranges over $[1.0000, 1.1250]$ across the twenty-six, and drifts *downwards* with n , attaining exactly 1 at orders 50, 56, 74, 92 and 110.

That is a selection effect, not a measurement. The certificates come from one construction — a capped unrolling of a periodic strip — and Theorem 6.1 extends it by inserting copies of a single period, so every member of the family inherits that period’s degree profile. A construction emitting one graph per order reports on the construction, not on the distribution over all graphs of that order. The clustering at $v_4/v_3 = 1$ is precisely what a periodic family looks like and precisely what a sample would not.

The methods of this paper cannot reach the question, for a structural reason. Everything proved above is an existence statement: one graph per order, or a class shown empty. A density function requires a measure over all graphs of order n , and nothing here enumerates. The class is moreover rigid in a way that obstructs sampling: a sweep over every two-edge switch of all twenty-six certificates finds no switch leaving a valid $(3, 4, 5)$ -alternating plane graph, so the one-switch neighbourhood of each is empty and local moves cannot walk the space.

Three routes seem plausible. An exhaustive census at small orders, with `plantri` or an equivalent generator restricted to the degree and face-size profile, would give the first honest histogram of v_4/v_3 ; it is a means to a conjecture rather than to a theorem, and it is where we would start. A structure theorem giving *every* $(3, 4, 5)$ -alternating plane graph a cap-strip-cap decomposition with a bounded interface alphabet would make the count a regular language and the density a transfer-matrix computation; the obstacle is the word *every*, since we have a construction and not a classification. Failing either, a bijection with an already-counted family of plane maps would serve, and the identities of Theorem 2.5 are the natural hook.

9. THE ARTIFACT

Every graph asserted in this paper is supplied as an explicit rotation system in the accompanying artifact, in a documented JSON format with a

dependency-free reference reader. No file records a claim that a graph *is* an alternating plane graph: degrees, faces, face sizes, bridges, connectivity and both alternation conditions are recomputed on every run by three separately written decision procedures. Certificates can also be exported to `planar_code` for checking by `plantri` or House of Graphs.

The artifact runs 1253 checks and needs only a Python interpreter; the mathematics uses no third-party library. Two classes of check are load-bearing: that each certificate is accepted by every verifier, and that each source in Theorem 7.1 is necessary, in the sense that removing it reopens orders. Each gate is paired with a control that must fail, so that a check cannot pass by being vacuous.

The two computations that the proofs genuinely rest on are the finite hypotheses of Remark 6.2.

REFERENCES

- [1] I. Althöfer, J. K. Haugland, K. Scherer, F. Schneider and N. Van Cleemput, *Alternating plane graphs*, Ars Math. Contemp. **8** (2015), 337–363. doi:10.26493/1855-3974.584.09a.
Email address: hanyu.yang.92@gmail.com
URL: ORCID 0009-0005-0419-4070