

# Seminar on Algebra, Geometry and Discrete Mathematics

Severino Da Dalt

March 22, 2025

## 1 Proof of asymptotic behaviour of eigenvalues in families of expanders

The main topic of the seminar is the study of families of expanders in the context of connected,  $k$ -regular and finite graphs. A family of expanders consists in a sequence of graphs of increasing size such that the isoperimetric number  $h(X)$  satisfies  $h(X) \geq \epsilon$  for some fixed positive  $\epsilon$  and for all graphs  $X$  in the sequence. The isoperimetric number, in the context of finite graphs, can be defined as:

$$h(x) = \min \left\{ \frac{|\partial F|}{|F|} \mid F \subseteq V, |F| \leq \frac{n}{2} \right\}$$

Earlier in the seminar, we established that the isoperimetric number is bounded by the spectral gap, in particular,

$$\frac{k - \mu_1}{2} \leq h(x) \leq \sqrt{2k(k - \mu_1)}$$

where  $\mu_1$  denotes the first nonzero eigenvalue. Moreover, we showed that the spectral gap cannot grow too much, i.e.

$$\liminf_{m \rightarrow +\infty} \mu_1 \geq 2\sqrt{k-1}$$

In this session, we aim to strengthen this result by proving that a non-trivial fraction of the eigenvalues of the graph lies within the interval  $[(2 - \epsilon)\sqrt{k-1}, k]$  for any given  $\epsilon > 0$ .

To prove this, we introduce a set of matrices  $A_r$ , which are polynomials in the incidence matrix  $A$ . The entries of  $A_r$  describe the number of

backtracking-free paths of length  $r$  between two vertices.

After establishing the necessary properties of these polynomials, we compute their generating function which is closely related to the generating function of Chebyshev polynomials  $(U_m)_{m \in \mathbb{N}}$ . This fact allows us to prove the following link between the two:

$$\sum_{x \in V} \sum_{0 \leq r \leq \frac{m}{2}} (A_{m-2r})_{xx} = (k-1)^{\frac{m}{2}} \sum_{j=0}^{n-1} U_m \left( \frac{\mu_j}{2\sqrt{k-1}} \right)$$

for all  $m \in \mathbb{N}$ . Chebyshev polynomials will be instrumental in introducing measures in the proof.

The key measure we use is:

$$\nu = \frac{1}{n} \sum_{j=0}^{n-1} \delta_{\frac{\mu_j}{\sqrt{k-1}}}$$

where  $\delta_a$  is the Dirac measure at  $a$ , and  $\mu_0, \dots, \mu_{n-1}$  are the eigenvalues of the graph. The Dirac measure  $\delta_a$  acts as an indicator function of whether  $a$  lies in a given interval. Hence, the measure  $\nu$  counts the number of eigenvalues within a specific range.

We then prove that any measure  $\nu'$  for which

$$\int_{-L}^L U_m \left( \frac{x}{2} \right) d\nu' \geq 0$$

for all  $m$ , satisfies that

$$\nu'[2 - \epsilon, L] \geq C$$

where  $C$  is a constant that only depends on  $\epsilon$  and  $L$ , and is strictly positive.

Finally, we demonstrate that  $\nu$  satisfies this condition and, with a suitable choice of  $L$ , we conclude the desired result.

As an additional result, we show a similar conclusion for the negative eigenvalues of the graph, generalizing an earlier result.