

Minicourse on *information, complexity and organisation in
multidimensional symbolic dynamics*

Exact computations of entropy for multidimensional SFT


Silvere Gangloff

April 30, 2021

sgangloff@agh.edu.pl ; silvere.gangloff@gmx.com


Multidimensional SFT: a computational 'transition':

Reminder (third lecture):




Multidimensional SFT: a computational 'transition':

Reminder (third lecture):



Multidimensional SFT: a computational 'transition':


Reminder (third lecture):



In practice, formula for entropy: development of tools for analysis of many variables systems.

Multidimensional SFT: a computational 'transition':

Reminder (third lecture):



In practice, formula for entropy: development of tools for analysis of many variables systems.


Question: what makes the entropy of subshifts in the liminal area computable ?

Dimers model:

Subshift X_0 :


Dimers model:

Subshift X_0 :



Dimers model:


Subshift X_0 :



Theorem[[Kasteleyn\(1961\)](#)]: $h(X_0) = \frac{G}{\pi}$,

Dimers model:

Subshift X_0 :




Theorem[[Kasteleyn\(1961\)](#)]: $h(X_0) = \frac{G}{\pi}$, where:

$$G = \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n+1)^2}.$$

Dimers model:

Subshift X_0 :




Theorem[[Kasteleyn\(1961\)](#)]: $h(X_0) = \frac{G}{\pi}$, where:


$$G = \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n+1)^2}.$$

(Called Catalan constant)

Ideas of the proof: Symmetries:




Ideas of the proof: Symmetries:



W


Ideas of the proof: Symmetries:



w

Number of such patterns: $N_n^w(X_0)$.

Ideas of the proof: Symmetries:




w

Number of such patterns: $N_n^{\mathbf{w}}(X_0)$.

Denote $\overline{\mathbf{w}}$ symmetric of \mathbf{w} .

Ideas of the proof: Symmetries:




w

Number of such patterns: $N_n^{\mathbf{w}}(X_0)$.

Denote $\overline{\mathbf{w}}$ symmetric of \mathbf{w} .

Lemma: $N_n^{\mathbf{w}}(X_0) = N_n^{\overline{\mathbf{w}}}(X_0)$.

Ideas of the proof: Symmetries:



w


Number of such patterns: $N_n^w(X_0)$.

Denote \bar{w} symmetric of w .

Lemma: $N_n^w(X_0) = N_n^{\bar{w}}(X_0)$.

Proof: the symmetry map is an involution.

Ideas of the proof: Symmetries:



Number of such patterns: $N_n^w(X_0)$.

Denote \bar{w} symmetric of w .

Lemma: $N_n^w(X_0) = N_n^{\bar{w}}(X_0)$.

Proof: the symmetry map is an involution.

Ideas of the proof: Toric and cylindric entropies:

Ideas of the proof: Toric and cylindric entropies:

Definition: $h_t(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_{n,n}^t(X))}{n^2},$

Ideas of the proof: Toric and cylindric entropies:

Definition: $h_t(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_{n,n}^t(X))}{n^2},$

where $N_{m,n}^t(X)$ is the number of (m, n) rectangle patterns in which rules apply as on torus.

Ideas of the proof: Toric and cylindric entropies:

Definition: $h_t(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_{n,n}^t(X))}{n^2},$

where $N_{m,n}^t(X)$ is the number of (m, n) rectangle patterns in which rules apply as on torus.

Similar definitions $h_c(X)$, $N_{m,n}^c(X)$ for cylinders.


Ideas of the proof: Toric and cylindric entropies:

Definition: $h_t(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_{n,n}^t(X))}{n^2},$

where $N_{m,n}^t(X)$ is the number of (m, n) rectangle patterns in which rules apply as on torus.

Similar definitions $h_c(X)$, $N_{m,n}^c(X)$ for cylinders.

Examples:



Ideas of the proof: Toric entropy equals entropy:

Ideas of the proof: Toric entropy equals entropy:

Lemma: $h_t(X_0) = h(X_0)$.

Ideas of the proof: Toric entropy equals entropy:

Lemma: $h_t(X_0) = h(X_0)$.

Ideas of the proof: Toric entropy equals entropy:

Lemma: $h_t(X_0) = h(X_0)$.

Proof: we have that:


$$\sum_{\mathbf{w}} (N_n^{\mathbf{w}}(X_0))^2 \leq N_{2n,n}^c(X_0) \leq \left(\sum_{\mathbf{w}} N_n^{\mathbf{w}}(X_0) \right)^2.$$

Ideas of the proof: Toric entropy equals entropy:

Lemma: $h_t(X_0) = h(X_0)$.

Proof: we have that:

$$\sum_{\mathbf{w}} (N_n^{\mathbf{w}}(X_0))^2 \leq N_{2n,n}^c(X_0) \leq \left(\sum_{\mathbf{w}} N_n^{\mathbf{w}}(X_0) \right)^2.$$




Ideas of the proof: Toric entropy equals entropy:

Lemma: $h_t(X_0) = h(X_0)$.

Proof: we have that:

$$\sum_{\mathbf{w}} (N_n^{\mathbf{w}}(X_0))^2 \leq N_{2n,n}^c(X_0) \leq \left(\sum_{\mathbf{w}} N_n^{\mathbf{w}}(X_0) \right)^2.$$




Ideas of the proof: Toric entropy equals entropy:

Lemma: $h_t(X_0) = h(X_0)$.

Proof: we have that:

$$\sum_{\mathbf{w}} (N_n^{\mathbf{w}}(X_0))^2 \leq N_{2n,n}^c(X_0) \leq \left(\sum_{\mathbf{w}} N_n^{\mathbf{w}}(X_0) \right)^2.$$




Thus $h_c(X_0) = h(X_0)$.

Ideas of the proof: Toric entropy equals entropy:

Lemma: $h_t(X_0) = h(X_0)$.

Proof: we have that:

$$\sum_{\mathbf{w}} (N_n^{\mathbf{w}}(X_0))^2 \leq N_{2n,n}^c(X_0) \leq \left(\sum_{\mathbf{w}} N_n^{\mathbf{w}}(X_0) \right)^2.$$



Thus $h_c(X_0) = h(X_0)$.

In a similar way $h_t(X_0) = h(X_0)$.

Ideas of the proof:


Consider $n \geq 1$;

Ideas of the proof:

Consider $n \geq 1$; arrow pattern


Ideas of the proof:

Consider $n \geq 1$; arrow pattern (ex $n = 4$):



Ideas of the proof:


Consider $n \geq 1$; arrow pattern (ex $n = 4$):



Denote $K^{(n)}$, $n^2 \times n^2$ matrix

Ideas of the proof:


Consider $n \geq 1$; arrow pattern (ex $n = 4$):



Denote $K^{(n)}$, $n^2 \times n^2$ matrix s.t. $K^{(n)}(\mathbf{i}, \mathbf{j}) = \pm 1$ when \mathbf{i}, \mathbf{j} adjacent, otherwise 0.

Ideas of the proof:


Consider $n \geq 1$; arrow pattern (ex $n = 4$):



Denote $K^{(n)}$, $n^2 \times n^2$ matrix s.t. $K^{(n)}(\mathbf{i}, \mathbf{j}) = \pm 1$ when \mathbf{i}, \mathbf{j} adjacent, otherwise 0. The coefficient is 1 when $\mathbf{i} \rightarrow \mathbf{j}$ and -1 when $\mathbf{i} \leftarrow \mathbf{j}$.

Ideas of the proof:

Consider $n \geq 1$; arrow pattern (ex $n = 4$):




Denote $K^{(n)}$, $n^2 \times n^2$ matrix s.t. $K^{(n)}(\mathbf{i}, \mathbf{j}) = \pm 1$ when \mathbf{i}, \mathbf{j} adjacent, otherwise 0. The coefficient is 1 when $\mathbf{i} \rightarrow \mathbf{j}$ and -1 when $\mathbf{i} \leftarrow \mathbf{j}$.

$$N_n^t(X_0) = \det(K^{(n)}) = \sum_{\sigma \in S_{n^2}} s(\sigma) \prod_{\mathbf{i}} K_{\mathbf{i}, \sigma(\mathbf{i})}^{(n)}$$

Ideas of the proof:

Consider $n \geq 1$; arrow pattern (ex $n = 4$):




Denote $K^{(n)}$, $n^2 \times n^2$ matrix s.t. $K^{(n)}(\mathbf{i}, \mathbf{j}) = \pm 1$ when \mathbf{i}, \mathbf{j} adjacent, otherwise 0. The coefficient is 1 when $\mathbf{i} \rightarrow \mathbf{j}$ and -1 when $\mathbf{i} \leftarrow \mathbf{j}$.


$$N_n^t(X_0) = \det(K^{(n)}) = \sum_{\sigma \in S_{n^2}} s(\sigma) \prod_{\mathbf{i}} K_{\mathbf{i}, \sigma(\mathbf{i})}^{(n)}$$

Diagonalisation of $K^{(n)} \rightarrow$ formula for $N_n^t(X_0)$ as sum of trigonometric functions.


Square ice: Wang tiles representation:




Square ice: Wang tiles representation:




Square ice: Wang tiles representation:




Square ice: Wang tiles representation:





Square ice: Wang tiles representation:




Square ice: Wang tiles representation:



Square ice: Subshift X^S :




Square ice: Subshift X^S :



E.H. Lieb, *Residual entropy of square ice*, Physical Review, 1967.

Square ice: Subshift X^S :



E.H. Lieb, *Residual entropy of square ice*, Physical Review, 1967.

S. Gangloff, *A proof that square ice entropy is $\frac{3}{2} \log_2(4/3)$* , 2019
(based on the work of R.Baxter, K.Kozlowski).

Square ice: Computing entropy using stripes subshifts:

$$h(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_n(X))}{n^2}.$$

Square ice: Computing entropy using stripes subshifts:


$$h(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_n(X))}{n^2}.$$

$$h(X) = \lim_{m,n} \frac{\log_2(N_{m,n}(X))}{mn}.$$

Square ice: Computing entropy using stripes subshifts:

$$h(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_n(X))}{n^2}.$$

$$h(X) = \lim_{m,n} \frac{\log_2(N_{m,n}(X))}{mn}.$$




Square ice: Computing entropy using stripes subshifts:


$$h(X) \stackrel{\text{def}}{=} \lim_n \frac{\log_2(N_n(X))}{n^2}.$$

$$h(X) = \lim_{m,n} \frac{\log_2(N_{m,n}(X))}{mn}.$$


$$h(X) = \lim_N \frac{h(X_N)}{N}$$




Square ice: Cylindric stripes subshifts:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric stripes subshifts:




Symmetry properties:




Square ice: Cylindric stripes subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:




Square ice: Cylindric subshifts:




Symmetry properties:



Square ice: Cylindric stripes subshifts:




Symmetry properties:




$$h(X^s) = \lim_N \frac{h(\overline{X}_N^s)}{N}$$


Square ice: Lieb transfer matrices:




Square ice: Lieb transfer matrices:




Square ice: Lieb transfer matrices:




Square ice: Lieb transfer matrices:




Square ice: Lieb transfer matrices:



Square ice: Lieb transfer matrices:




Square ice: Lieb transfer matrices:



$$V_N(t)[u, v] = \sum_{u \mathcal{R}[w] v} t^{|w|}.$$

where $|w| = \#$ of  and 

Square ice: Lieb transfer matrices:



$$V_N(t)[u, v] = \sum_{u \mathcal{R}[w] v} t^{|w|}.$$


where $|w| = \#$ of  and 

$$h(X^s) = \lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N}$$


Square ice: Computing maximal eigenvalue of $V_N(1)$,
strategy:

$$V_N(1) \quad \bullet$$


Square ice: Computing maximal eigenvalue of $V_N(1)$,
strategy:



Square ice: Computing maximal eigenvalue of $V_N(1)$,
strategy:




Square ice: Computing maximal eigenvalue of $V_N(1)$, strategy:




$$\forall x, y, T_N^y(t) T_N^x(t) = T_N^y(t) T_N^x(t).$$

Square ice: Computing maximal eigenvalue of $V_N(1)$, strategy:




$$\forall x, y, \quad T_N^y(t) T_N^x(t) = T_N^y(t) T_N^x(t).$$

Square ice: Computing maximal eigenvalue of $V_N(1)$, strategy:



$$\forall x, y, \quad T_N^y(t) T_N^x(t) = T_N^x(t) T_N^y(t).$$

Square ice: Computing maximal eigenvalue of $V_N(1)$, strategy:



$$\forall x, y, \quad T_N^y(t) T_N^x(t) = T_N^y(t) T_N^x(t).$$


Square ice: R-matrices and monodromy matrices:

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \left(\begin{array}{cc} & \\ & \end{array} \right)$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & \\ & \end{pmatrix}$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ & \end{pmatrix}$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & \end{pmatrix}$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$


Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$




Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$




Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$




Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$




Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:



$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$




Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$


Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$



Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:




$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$




$$M_N(1,0)[\epsilon, \eta]$$

Square ice: R-matrices and monodromy matrices: \rightarrow :
input/output:



$$R(0,1) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$$




$$M_N(1,0)[\epsilon, \eta]$$

Yang-Baxter transfer matrices:

$$T_N[\epsilon, \eta] = \sum_{u \in \{0,1\}} M_N(u, u)[\epsilon, \eta].$$


Square ice: Commutation of Yang-Baxter matrices:

Composition of R -matrices and monodromy matrices:




Square ice: Commutation of Yang-Baxter matrices:

Composition of R -matrices and monodromy matrices:




Yang-Baxter equation:




Square ice: Commutation of Yang-Baxter matrices:

Composition of R -matrices and monodromy matrices:



Yang-Baxter equation:



Yang-Baxter equation \Rightarrow transfer matrices commute.

Square ice: Trigonometric R -matrices: $t \in (0, \sqrt{2}]$.

Square ice: Trigonometric R -matrices: $t \in (0, \sqrt{2}]$.

Denote $\mu_t \in (0, \frac{\pi}{2})$ s.t. $\cos(\mu_t) = \frac{2-t^2}{2}$.

Square ice: Trigonometric R -matrices: $t \in (0, \sqrt{2}]$.

Denote $\mu_t \in (0, \frac{\pi}{2})$ s.t. $\cos(\mu_t) = \frac{2-t^2}{2}$.

$$R_{\mu_t}^x = \frac{1}{\sin(\mu_t/2)} \begin{pmatrix} \sin(\mu_t - x) & 0 & 0 & 0 \\ 0 & \sin(x) & \sin(\mu_t) & 0 \\ 0 & \sin(\mu_t) & \sin(x) & 0 \\ 0 & 0 & 0 & \sin(\mu_t - x) \end{pmatrix}.$$

Square ice: Trigonometric R -matrices: $t \in (0, \sqrt{2}]$.

Denote $\mu_t \in (0, \frac{\pi}{2})$ s.t. $\cos(\mu_t) = \frac{2-t^2}{2}$.

$$R_{\mu_t}^x = \frac{1}{\sin(\mu_t/2)} \begin{pmatrix} \sin(\mu_t - x) & 0 & 0 & 0 \\ 0 & \sin(x) & \sin(\mu_t) & 0 \\ 0 & \sin(\mu_t) & \sin(x) & 0 \\ 0 & 0 & 0 & \sin(\mu_t - x) \end{pmatrix}.$$

where $R_{\mu_t}^x(0, 0)$ is the up-left 2×2 part of this matrix, etc.

Square ice: Bethe ansatz:

Square ice: Bethe ansatz:

If there exists $(p_j(t))_{j=1}^n$ solution of:

$$Np_j = 2\pi j - (n+1)\pi - \sum_{k=1}^n \Theta_t(p_j, p_k) \quad (E_t)$$

Square ice: Bethe ansatz:

If there exists $(p_j(t))_{j=1}^n$ solution of:

$$Np_j = 2\pi j - (n+1)\pi - \sum_{k=1}^n \Theta_t(p_j, p_k) \quad (E_t)$$

construction of a candidate eigenvector for the transfer matrix for value:

$$\prod_{k=1}^n L_t(e^{ip_k}) + \prod_{k=1}^n M_t(e^{ip_k}).$$

where L_t and M_t are simple rational functions.

Square ice: Bethe ansatz:

If there exists $(p_j(t))_{j=1}^n$ solution of:

$$Np_j = 2\pi j - (n+1)\pi - \sum_{k=1}^n \Theta_t(p_j, p_k) \quad (E_t)$$

construction of a candidate eigenvector for the transfer matrix for value:

$$\prod_{k=1}^n L_t(e^{ip_k}) + \prod_{k=1}^n M_t(e^{ip_k}).$$

where L_t and M_t are simple rational functions.

Lemma: for all t there exists $(p_j(t))_{j=1}^n$ solution of (E_t) and it is an analytic function in t .

Square ice: Identification to $\lambda_{\max}(V_N(t))$: ingredients:

Square ice: Identification to $\lambda_{\max}(V_N(t))$: ingredients:

1. When $t = \sqrt{2}$, simplified expression of candidate eigenvector and eigenvalue.

Square ice: Identification to $\lambda_{\max}(V_N(t))$: ingredients:

1. When $t = \sqrt{2}$, simplified expression of candidate eigenvector and eigenvalue.
2. There is H_N (given by derivative of $V_N(t)$) diagonalisable s.t. $V_N(\sqrt{2})H_N = H_N V_N(\sqrt{2})$: candidate value is the maximal eigenvalue.

Square ice: Identification to $\lambda_{\max}(V_N(t))$: ingredients:

1. When $t = \sqrt{2}$, simplified expression of candidate eigenvector and eigenvalue.
2. There is H_N (given by derivative of $V_N(t)$) diagonalisable s.t. $V_N(\sqrt{2})H_N = H_N V_N(\sqrt{2})$: candidate value is the maximal eigenvalue.
3. Perron-Frobenius: candidate eigenvector has positive coordinates \rightarrow eigenvector for the maximal eigenvalue.

Square ice: Identification to $\lambda_{\max}(V_N(t))$: ingredients:

1. When $t = \sqrt{2}$, simplified expression of candidate eigenvector and eigenvalue.
2. There is H_N (given by derivative of $V_N(t)$) diagonalisable s.t. $V_N(\sqrt{2})H_N = H_N V_N(\sqrt{2})$: candidate value is the maximal eigenvalue.
3. Perron-Frobenius: candidate eigenvector has positive coordinates \rightarrow eigenvector for the maximal eigenvalue.
4. Identification of maximal eigenvalue and eigenvector around $\sqrt{2}$ (positive coordinates), then on $(0, \sqrt{2})$ by analyticity.

Square ice: Asymptotics:

Square ice: Asymptotics: consider $n_k/N_k \rightarrow 1/2$.

Square ice: Asymptotics: consider $n_k/N_k \rightarrow 1/2$.

Denote $(\mathbf{p}_j^{(k)})_j$ solutions of equations (E_t) for N_k, n_k .

Square ice: Asymptotics: consider $n_k/N_k \rightarrow 1/2$.

Denote $(\mathbf{p}_j^{(k)})_j$ solutions of equations (E_t) for N_k, n_k .

Change of variable $(\mathbf{p}_j^{(k)})_j = (\kappa_t(\boldsymbol{\alpha}_j^{(k)}))_j$.

Square ice: Asymptotics: consider $n_k/N_k \rightarrow 1/2$.

Denote $(\mathbf{p}_j^{(k)})_j$ solutions of equations (E_t) for N_k, n_k .

Change of variable $(\mathbf{p}_j^{(k)})_j = (\kappa_t(\boldsymbol{\alpha}_j^{(k)}))_j$.

Lemma:

$$\lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N} = \lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\boldsymbol{\alpha}_j^{(k)}),$$

with $f(x) = \log_2(2|\sin(\kappa_t(x))/2|)$.

Square ice: Asymptotics: consider $n_k/N_k \rightarrow 1/2$.

Denote $(\mathbf{p}_j^{(k)})_j$ solutions of equations (E_t) for N_k, n_k .

Change of variable $(\mathbf{p}_j^{(k)})_j = (\kappa_t(\alpha_j^{(k)}))_j$.

Lemma:

$$\lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N} = \lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\alpha_j^{(k)}),$$


with $f(x) = \log_2(2|\sin(\kappa_t(x))/2|)$.

Theorem: there exists ρ_t s.t. for all $f \in L^1$:

$$\lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\alpha_j^{(k)}) = \int_{\mathbb{R}} f(\alpha) \rho_t(\alpha) d\alpha.$$


Square ice: Counting functions:

$$\xi_t^{(k)} : \alpha \mapsto \frac{1}{2\pi} \kappa_t(\alpha) + \frac{n_k + 1}{2N_k} + \frac{1}{2\pi N_k} \sum_{j=1}^{n_k} \theta_t(\alpha, \alpha_j^{(k)})$$



Square ice: Counting functions:

$$\xi_t^{(k)} : \alpha \mapsto \frac{1}{2\pi} \kappa_t(\alpha) + \frac{n_k + 1}{2N_k} + \frac{1}{2\pi N_k} \sum_{j=1}^{n_k} \theta_t(\alpha, \alpha_j^{(k)})$$



Equality with a Riemann sum:


$$\lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\alpha_j^{(k)}) = \lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} \left(\alpha_{j+1}^{(k)} - \alpha_j^{(k)} \right) \frac{\left(\xi_t^{(k)}(\alpha_{j+1}^{(k)}) - \xi_t^{(k)}(\alpha_j^{(k)}) \right)}{\left(\alpha_{j+1}^{(k)} - \alpha_j^{(k)} \right)} f(\alpha_j^{(k)})$$

Square ice: Asymptotics of counting functions:

Square ice: Asymptotics of counting functions:


Square ice: Asymptotics of counting functions:

1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :




Square ice: Asymptotics of counting functions:

1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :




Square ice: Asymptotics of counting functions:

1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :



Square ice: Asymptotics of counting functions:


1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :



2. Assume $((\xi_t)^{\nu(k)})' \rightarrow F_t$ on any compact K .

Square ice: Asymptotics of counting functions:


1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :



2. Assume $((\xi_t)^{\nu(k)})' \rightarrow F_t$ on any compact K .

Square ice: Asymptotics of counting functions:


1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :



2. Assume $((\xi_t)^{\nu(k)})' \rightarrow F_t$ on any compact K .

Square ice: Asymptotics of counting functions:

1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :




2. Assume $((\xi_t)^{\nu(k)})' \rightarrow F_t$ on any compact K .

Square ice: Asymptotics of counting functions:

1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :


$$(\xi_t^{(+\infty)})'_{|K}$$



2. Assume $((\xi_t)^{\nu(k)})' \rightarrow F_t$ on any compact K .

Square ice: Asymptotics of counting functions:

1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :




2. Assume $((\xi_t)^{\nu(k)})' \rightarrow F_t$ on any compact K .
3. F_t satisfies an integral equation with unique solution.

Square ice: Asymptotics of counting functions:


1. Extend $(\xi_t^{(k)})'$ on a stripe including \mathbb{R} :

$$(\xi_t^{(+\infty)})'_{|K}$$




2. Assume $((\xi_t)^{\nu(k)})' \rightarrow F_t$ on any compact K .
3. F_t satisfies an integral equation with unique solution.
4. Thus, $(\xi_t^{(k)})'$ converges to this solution.


Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
biholomorphisms: $\epsilon > 0$:




Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
biholomorphisms: $\epsilon > 0$:




Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
 biholomorphisms: $\epsilon > 0$:




Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
 biholomorphisms: $\epsilon > 0$:




Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
 biholomorphisms: $\epsilon > 0$:




Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
 biholomorphisms: $\epsilon > 0$:



Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
 biholomorphisms: $\epsilon > 0$:




Square ice: Rarefaction of the roots and $\xi_t^{(k)}$
 biholomorphisms: $\epsilon > 0$:




The functions $\xi_t^{(k)}$ have distinct values on \mathcal{V} and Γ . Thus they are biholomorphisms onto \mathcal{V} (Cauchy formula).

Square ice: Lax integral expression of $\xi_t^{(k)}$:



Square ice: Llace integral expression of $\xi_t^{(k)}$:



By residues theorem:

$$\xi_t^{(k)}(\alpha) = \frac{1}{2\pi} \kappa_t(\alpha) + \frac{n_k + 1}{2N_k} + \oint_{\Gamma} \theta_t \left(\left(\xi_t^{(k)} \right)^{-1}(\alpha) \right) \frac{e^{2i\pi s N_k}}{e^{2i\pi s N_k} - 1} ds + O(\epsilon).$$

Square ice: Fredholm integral equation: Limit and change of variable:

$$\xi_t^{(\infty)}(\alpha) = \frac{1}{2\pi} \kappa_t(\alpha) + \frac{1}{4} + \int_0^{+\infty} \theta_t(\alpha) \left(\xi_t^{(\infty)} \right)'(\alpha) d\alpha.$$

Square ice: Fredholm integral equation: Limit and change of variable:

$$\xi_t^{(\infty)}(\alpha) = \frac{1}{2\pi} \kappa_t(\alpha) + \frac{1}{4} + \int_0^{+\infty} \theta_t(\alpha) \left(\xi_t^{(\infty)} \right)'(\alpha) d\alpha.$$

Unique solution by Fourier transforms.

Square ice: Final computation:

Square ice: Final computation:

$$\lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N} = \lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\alpha_j^{(k)}),$$

with $f(x) = \log_2(2|\sin(\kappa_t(x))/2|)$.

Square ice: Final computation:

$$\lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N} = \lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\alpha_j^{(k)}),$$

with $f(x) = \log_2(2|\sin(\kappa_t(x))/2|)$.

$$h(X^s) = \int_{\mathbb{R}} \log_2(2|\sin(\kappa_t(\alpha))/2|) \cdot \rho_t(\alpha) d\alpha.$$

Square ice: Final computation:

$$\lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N} = \lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\alpha_j^{(k)}),$$

with $f(x) = \log_2(2|\sin(\kappa_t(x))/2|)$.

$$h(X^s) = \int_{\mathbb{R}} \log_2(2|\sin(\kappa_t(\alpha))/2|) \cdot \rho_t(\alpha) d\alpha.$$

Expression of $\rho_t = \left(\xi_t^{(\infty)}\right)'$ and lace integrals computations:

$$h(X^s) = \frac{3}{2} \log_2(4/3).$$

Friedland's theorem:

Theorem[Friedland(1967)]: if the set of forbidden patterns \mathcal{F} is stable by symmetry, $h(X_{\mathcal{F}})$ is a computable number.

Friedland's theorem:

Theorem[Friedland(1967)]: if the set of forbidden patterns \mathcal{F} is stable by symmetry, $h(X_{\mathcal{F}})$ is a computable number.

Examples: dimers, square ice, hard squares.


Friedland's theorem:

Theorem[Friedland(1967)]: if the set of forbidden patterns \mathcal{F} is stable by symmetry, $h(X_{\mathcal{F}})$ is a computable number.



Examples: dimers, square ice, hard squares.

Question: what are the possible values of entropy for symmetric bidimensional subshifts of finite type ?


Baxter's hard hexagons: [Exactly solvable models in statistical physics]



Baxter's hard hexagons: [Exactly solvable models in statistical physics]




Baxter's hard hexagons: [Exactly solvable models in statistical physics]



Formula for entropy as sum of a series:


Baxter's hard hexagons: [Exactly solvable models in statistical physics]



Formula for entropy as sum of a series:

1. Transfer matrices \rightarrow **diagonal** transfer matrices;


Baxter's hard hexagons: [Exactly solvable models in statistical physics]



Formula for entropy as sum of a series:

1. Transfer matrices \rightarrow **diagonal** transfer matrices;
2. Obtain a functional equation;


Baxter's hard hexagons: [Exactly solvable models in statistical physics]



Formula for entropy as sum of a series:

1. Transfer matrices \rightarrow **diagonal** transfer matrices;
2. Obtain a functional equation;
3. Maximal eigenvalue satisfies the same functional equation;


Baxter's hard hexagons: [Exactly solvable models in statistical physics]



Formula for entropy as sum of a series:

1. Transfer matrices \rightarrow **diagonal** transfer matrices;
2. Obtain a functional equation;
3. Maximal eigenvalue satisfies the same functional equation;
4. Maximal eigenvalue equal for the two diagonal transfer matrices \rightarrow other function equation.


Baxter's hard hexagons: [Exactly solvable models in statistical physics]



Formula for entropy as sum of a series:

1. Transfer matrices \rightarrow **diagonal** transfer matrices;
2. Obtain a functional equation;
3. Maximal eigenvalue satisfies the same functional equation;
4. Maximal eigenvalue equal for the two diagonal transfer matrices \rightarrow other function equation.
5. System of equations with unique solution: evaluate it in specific parameter.

Baxter's hard hexagons: [Exactly solvable models in statistical physics]




Formula for entropy as sum of a series:


1. Transfer matrices \rightarrow **diagonal** transfer matrices;
2. Obtain a functional equation;
3. Maximal eigenvalue satisfies the same functional equation;
4. Maximal eigenvalue equal for the two diagonal transfer matrices \rightarrow other function equation.
5. System of equations with unique solution: evaluate it in specific parameter.

Main problems: points **3, 4**.


Eight-vertex model: [Exactly solvable models in statistical physics]




Eight-vertex model: [Exactly solvable models in statistical physics]



Eight-vertex model: [Exactly solvable models in statistical physics]



Eight-vertex model: [Exactly solvable models in statistical physics]



Entropy computation: similar to square ice; analytical part non verified.

Subsidiary questions:

Question: can we use similar methods to talk about invariant measures (for instance $\times 2, \times 3$ conjecture) ?


Subsidiary questions:

Question: can we use similar methods to talk about invariant measures (for instance $\times 2$, $\times 3$ conjecture) ?

Question: can we find solutions of Yang-Baxter equations for other subshifts of finite type ? *Example:* Kari-Culik tilings (know: positive entropy [[Durand, Gamard, Grandjean\(2017\)](#)])).


Combinatorial methods

Definition subshift Δ_r ; ex for $r = 3$:



Combinatorial methods


Definition subshift Δ_r ; ex for $r = 3$:




| | | | | | | | | | |
|---|---|---|---|---|---|---|---|--|--|
| 3 | 1 | 2 | 3 | | | | | | |
| | | | | 1 | 2 | 3 | 1 | | |
| 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | | |
| 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 | | |
| 3 | 1 | 2 | 3 | | | | | | |
| 2 | 3 | | | 1 | 2 | 3 | 1 | | |
| | | 1 | 2 | 3 | | | | | |
| 1 | 2 | 3 | | | 1 | 2 | 3 | | |

Combinatorial methods

Definition subshift Δ_r ; ex for $r = 3$:




| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 3 | 1 | 2 | 3 | | | | |
| | | | | 1 | 2 | 3 | 1 |
| 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 |
| 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 |
| 3 | 1 | 2 | 3 | | | | |
| 2 | 3 | | | 1 | 2 | 3 | 1 |
| | | 1 | 2 | 3 | | | |
| 1 | 2 | 3 | | | 1 | 2 | 3 |




Combinatorial methods

Definition subshift Δ_r ; ex for $r = 3$:




| | | | | | | | | | |
|---|---|---|---|---|---|---|---|--|--|
| 3 | 1 | 2 | 3 | | | | | | |
| | | | | 1 | 2 | 3 | 1 | | |
| 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 | | |
| 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 | | |
| 3 | 1 | 2 | 3 | | | | | | |
| 2 | 3 | | | 1 | 2 | 3 | 1 | | |
| | | 1 | 2 | 3 | | | | | |
| 1 | 2 | 3 | | | 1 | 2 | 3 | | |




Theorem:[G., Sablik] $h(\Delta_r) = \frac{\log_2(r+1)}{r}$.

Combinatorial methods

Definition subshift Δ_r ; ex for $r = 3$:



| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 3 | 1 | 2 | 3 | | | | |
| | | | | 1 | 2 | 3 | 1 |
| 1 | 2 | 3 | 1 | 2 | 3 | 1 | 2 |
| 2 | 3 | 1 | 2 | 3 | 1 | 2 | 3 |
| 3 | 1 | 2 | 3 | | | | |
| 2 | 3 | | | 1 | 2 | 3 | 1 |
| | | 1 | 2 | 3 | | | |
| 1 | 2 | 3 | | | 1 | 2 | 3 |



Theorem:[G.,Sablik] $h(\Delta_r) = \frac{\log_2(r+1)}{r}$.

Question: for what kind of subshifts can we compute entropy with similar methods ?

Proof for $r = 1$

Proof for $r = 1$

Second layer is trivial: we consider only first and third.

Proof for $r = 1$

Second layer is trivial: we consider only first and third.

Lower bound:

Proof for $r = 1$

Second layer is trivial: we consider only first and third.

Lower bound: $\mathcal{L}_n(\Delta_1)$ contains the following patterns:

Proof for $r = 1$

Second layer is trivial: we consider only first and third.


Lower bound: $\mathcal{L}_n(\Delta_1)$ contains the following patterns:



Proof for $r = 1$

Second layer is trivial: we consider only first and third.

Lower bound: $\mathcal{L}_n(\Delta_1)$ contains the following patterns:




$$N_n(\Delta_1) \geq 2^{n^2}.$$

Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :


Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :



Proof for $r = 1$


Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :



2^3 choices in 2nd layer

Proof for $r = 1$


Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :



2^3 choices in 2nd layer
 2^4 choices in total


Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :



Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :




Number of possible extensions into a pattern of $\mathcal{L}_{n+1}(\Delta_1)$:
 $\leq 2^{4n} 2^{3 \cdot 3 \cdot 4} = 2^{4n+4} 2^C.$

$$2^{n^2} \leq N_n(\Delta_1) \leq 2^{Cn} \cdot 2^{n^2}.$$

Proof for $r = 1$

Upper bound: Consider a pattern in $\mathcal{L}_n(\Delta_1)$ for some n :



Number of possible extensions into a pattern of $\mathcal{L}_{n+1}(\Delta_1)$:
 $\leq 2^{4n} 2^{3 \cdot 3 \cdot 4} = 2^{4n+4} 2^C.$

$$2^{n^2} \leq N_n(\Delta_1) \leq 2^{Cn} \cdot 2^{n^2}.$$

$$h(\Delta_1) = 1 = \frac{\log_2(2)}{1}.$$

Combinatorial method for square ice entropy ?

Some ideas:

Combinatorial method for square ice entropy ?

Some ideas:

1. For instance: use Hochman-Meyerovitch's theorem to realize square ice entropy; then simplify the construction without changing the entropy; find a subshift which is isomorphic to square ice.

Combinatorial method for square ice entropy ?

Some ideas:

1. For instance: use Hochman-Meyerovitch's theorem to realize square ice entropy; then simplify the construction without changing the entropy; find a subshift which is isomorphic to square ice.
2. Find smaller and smaller subshifts of square ice with same entropy.

Combinatorial method for square ice entropy ?

Some ideas:

1. For instance: use Hochman-Meyerovitch's theorem to realize square ice entropy; then simplify the construction without changing the entropy; find a subshift which is isomorphic to square ice.
2. Find smaller and smaller subshifts of square ice with same entropy.
3. Try Baxter's method for square ice; more precisely: do both transfer matrices of square ice have same maximal eigenvalues or not ?