

A Cantor dynamical system is slow if and only if it has only attracting finite orbits.

**S.Gangloff**, joint work with **P.Oprocha**

*AGH, Faculty of Applied Mathematics, Kraków.*

sgangloff@agh.edu.pl ; silvere.gangloff@gmx.com

Problem

## Definitions

## Definitions

**Cantor space** : topological space  $X$  with a base of clopen sets and without isolated point.



## Definitions

**Cantor space** : topological space  $X$  with a base of clopen sets and without isolated point.



**Cantor system** :  $f : X \rightarrow X$  continuous on  $X$  Cantor space.

## Definitions

**Cantor space** : topological space  $X$  with a base of clopen sets and without isolated point.



**Cantor system** :  $f : X \rightarrow X$  continuous on  $X$  Cantor space.

**Embedding in  $\mathbb{R}$  with vanishing derivative** : exists  $\phi : X \rightarrow \mathbb{R}$  injective and  $g : \mathbb{R} \rightarrow \mathbb{R}$  s.t.  $\phi \circ f = g \circ \phi$ , and  $g'_{|\phi(X)|} \equiv 0$ .

**Problem :**

*What are the Cantor systems which can be embedded in  $\mathbb{R}$  with vanishing derivative ?*

Partial result



## Result of P.Oprocha and J.Boroński :

Minimal system  $\equiv$  every orbit  $\{f^n(x), n \geq 0\}$  is dense.

## Result of P.Oprocha and J.Boroński :

Minimal system  $\equiv$  every orbit  $\{f^n(x), n \geq 0\}$  is dense.

**Theorem**[Oprocha, Boroński] : Every minimal Cantor system can be embedded in  $\mathbb{R}$  with vanishing derivative.

## Jarník's theorem

**Theorem**[Jarník] : Let  $X \subset \mathbb{R}$  be a Cantor set and  $f : X \rightarrow \mathbb{R}$  differentiable. There exists a differentiable function  $\mathbb{R} \rightarrow \mathbb{R}$  which extends  $f$ .


## Jarník's theorem

**Theorem**[Jarník] : Let  $X \subset \mathbb{R}$  be a Cantor set and  $f : X \rightarrow \mathbb{R}$  differentiable. There exists a differentiable function  $\mathbb{R} \rightarrow \mathbb{R}$  which extends  $f$ .

Thus we only need to characterize the Cantor systems  $(X, f)$  with a function  $\phi : X \rightarrow \mathbb{R}$  injective whose derivative is zero.


## Graph coverings :

Graph morphisms :



## Graph coverings :

Graph morphisms :



**Graph coverings :**  $(G_n)_{n \geq 1} = (V_n, E_n)_{n \geq 1}$  finite simple oriented graphs and  $\pi_n : G_{n+1} \rightarrow G_n$  *graph morphisms*.

## Systems associated with graph coverings :

Consider  $(G_n, \pi_n)$  graph coverings.

## Systems associated with graph coverings :

Consider  $(G_n, \pi_n)$  graph coverings.

Let us denote  $V = \{\mathbf{u} = (u_n)_{n \geq 1} : \forall n, u_n \in G_n, \pi_n(u_{n+1}) = u_n\}$ .



## Systems associated with graph coverings :

Consider  $(G_n, \pi_n)$  graph coverings.

Let us denote  $V = \{\mathbf{u} = (u_n)_{n \geq 1} : \forall n, u_n \in G_n, \pi_n(u_{n+1}) = u_n\}$ .


Condition : for every  $\mathbf{u} \in V$ , there exists a unique  $\mathbf{v}$  such that for all  $n \geq 1$ ,  $(u_n, v_n) \in E_n$ .

## Systems associated with graph coverings :

Consider  $(G_n, \pi_n)$  graph coverings.

Let us denote  $V = \{\mathbf{u} = (u_n)_{n \geq 1} : \forall n, u_n \in G_n, \pi_n(u_{n+1}) = u_n\}$ .

Condition : for every  $\mathbf{u} \in V$ , there exists a unique  $\mathbf{v}$  such that for all  $n \geq 1$ ,  $(u_n, v_n) \in E_n$ .




## Systems associated with graph coverings :

Consider  $(G_n, \pi_n)$  graph coverings.

Let us denote  $V = \{\mathbf{u} = (u_n)_{n \geq 1} : \forall n, u_n \in G_n, \pi_n(u_{n+1}) = u_n\}$ .

Condition : for every  $\mathbf{u} \in V$ , there exists a unique  $\mathbf{v}$  such that for all  $n \geq 1$ ,  $(u_n, v_n) \in E_n$ .



By setting  $f(\mathbf{u}) = \mathbf{v}$ , we define a continuous function  $V \rightarrow V$ , and thus a dynamical system  $(V, f)$ .

## Representation of Cantor systems with graph coverings

**Theorem** : any Cantor system  $(X, f)$  is conjugated to a graph coverings system.

*Proof :*

## Representation of Cantor systems with graph coverings

**Theorem** : any Cantor system  $(X, f)$  is conjugated to a graph coverings system.

*Proof :*

**Input** : sequence of clopen partitions  $(\mathcal{U}_n)_{n \geq 1}$  s.t. for all  $n$ ,  $\mathcal{U}_{n+1} \prec \mathcal{U}_n$ , and  $\text{mesh}(\mathcal{U}_n) \rightarrow 0$ .

## Representation of Cantor systems with graph coverings

**Theorem** : any Cantor system  $(X, f)$  is conjugated to a graph coverings system.

*Proof* :

**Input** : sequence of clopen partitions  $(\mathcal{U}_n)_{n \geq 1}$  s.t. for all  $n$ ,  $\mathcal{U}_{n+1} \prec \mathcal{U}_n$ , and  $\text{mesh}(\mathcal{U}_n) \rightarrow 0$ .

**Graphs** :  $G_n$  given by  $V_n = \mathcal{U}_n$  and  $E_n = \{(u, v) : \exists x \in u, f(x) \in v\}$ .

## Representation of Cantor systems with graph coverings

**Theorem** : any Cantor system  $(X, f)$  is conjugated to a graph coverings system.

*Proof* :

**Input** : sequence of clopen partitions  $(\mathcal{U}_n)_{n \geq 1}$  s.t. for all  $n$ ,  $\mathcal{U}_{n+1} \prec \mathcal{U}_n$ , and  $\text{mesh}(\mathcal{U}_n) \rightarrow 0$ .

**Graphs** :  $G_n$  given by  $V_n = \mathcal{U}_n$  and  $E_n = \{(u, v) : \exists x \in u, f(x) \in v\}$ .

**Morphisms** :  $\pi_n(u) = w$  s.t.  $u \subset w$ .

## Representation of Cantor systems with graph coverings

**Theorem** : any Cantor system  $(X, f)$  is conjugated to a graph coverings system.

*Proof* :

**Input** : sequence of clopen partitions  $(\mathcal{U}_n)_{n \geq 1}$  s.t. for all  $n$ ,  $\mathcal{U}_{n+1} \prec \mathcal{U}_n$ , and  $\text{mesh}(\mathcal{U}_n) \rightarrow 0$ .

**Graphs** :  $G_n$  given by  $V_n = \mathcal{U}_n$  and  $E_n = \{(u, v) : \exists x \in u, f(x) \in v\}$ .


**Morphisms** :  $\pi_n(u) = w$  s.t.  $u \subset w$ .

**Conjugacy** :  $x \in X$  is associated with a unique sequence  $(u_n)$ .




## Gambeaudau-Martens representation of minimal systems :

If  $(X, f)$  is minimal, the graphs  $G_n$  can be taken under the following form :



## Gambeaudau-Martens representation of minimal systems :


If  $(X, f)$  is minimal, the graphs  $G_n$  can be taken under the following form :



→ used in the proof of P.Oprocha and J.Boroński.

## Gambeaudau-Martens representation of minimal systems :

If  $(X, f)$  is minimal, the graphs  $G_n$  can be taken under the following form :



→ used in the proof of P.Oprocha and J.Boroński.

We will expose this proof for particular systems : odometers.

## Odometers :

Let us consider  $X = \prod_{k=1}^{+\infty} \{0, 1\}$ .

## Odometers :

Let us consider  $X = \prod_{k=1}^{+\infty} \{0, 1\}$ .

**Clopen basis :** Cylinders  $[u] = \{x \in X : x_{[1,n]} = u\}$ , for  $u$  word on  $\{0, 1\}$  of length  $n \geq 0$ .

## Odometers :

Let us consider  $X = \prod_{k=1}^{+\infty} \{0, 1\}$ .

**Clopen basis :** Cylinders  $[u] = \{x \in X : x_{[1,n]} = u\}$ , for  $u$  word on  $\{0, 1\}$  of length  $n \geq 0$ .

**Dynamical system :**  $(X, f)$ , with  $f(x)$  defined as :

## Odometers :

Let us consider  $X = \prod_{k=1}^{+\infty} \{0, 1\}$ .

**Clopen basis :** Cylinders  $[u] = \{x \in X : x_{[1,n]} = u\}$ , for  $u$  word on  $\{0, 1\}$  of length  $n \geq 0$ .

**Dynamical system :**  $(X, f)$ , with  $f(x)$  defined as :

$$\begin{array}{rcl} x : & 1 & 1 & 1 & 1 & 0 & 1 & 0 & 1 & 1 & \cdots \\ f(x) : & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & \cdots \end{array}$$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.




**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$




$G_3$



$\vdots$


**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.




**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.



**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.



**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.

$G_1$




---

$G_2$



---

$G_3$



---

$\vdots$



**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.


$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Graph coverings representation and embedding.

Odometers are minimal  $\rightarrow$  Gambaudo-Martens representation.



$G_1$



$G_2$



$G_3$



$\vdots$

**Odometers** : Vanishing derivative of the embedding.

***Shrinking*** rate :  $2^{-k}$  for  $G_k$  + length of intervals  $\ll$  space between them.

**Odometers** : Vanishing derivative of the embedding.

***Shrinking*** rate :  $2^{-k}$  for  $G_k$  + length of intervals  $\ll$  space between them.

Consider  $k \geq 1$ , **u**, **v** sufficiently close :  $u_k = v_k$ .



**Odometers** : Vanishing derivative of the embedding.

***Shrinking*** rate :  $2^{-k}$  for  $G_k$  + length of intervals  $\ll$  space between them.

Consider  $k \geq 1$ ,  $\mathbf{u}$ ,  $\mathbf{v}$  sufficiently close :  $u_k = v_k$ .

Assume that  $k$  large enough s.t.  $u_k$  and  $v_k$  are not  $\bullet$ .

**Odometers** : Vanishing derivative of the embedding.

***Shrinking*** rate :  $2^{-k}$  for  $G_k$  + length of intervals  $\ll$  space between them.

Consider  $k \geq 1$ ,  $\mathbf{u}$ ,  $\mathbf{v}$  sufficiently close :  $u_k = v_k$ .

Assume that  $k$  large enough s.t.  $u_k$  and  $v_k$  are not  $\bullet$ .

By definition of embedding,  $d(\phi(\mathbf{u}), \phi(\mathbf{v})) \leq 2^{-k} d(\mathbf{u}, \mathbf{v})$ .

**Odometers** : Vanishing derivative of the embedding.

**Shrinking rate** :  $2^{-k}$  for  $G_k$  + length of intervals  $\ll$  space between them.

Consider  $k \geq 1$ ,  $\mathbf{u}$ ,  $\mathbf{v}$  sufficiently close :  $u_k = v_k$ .

Assume that  $k$  large enough s.t.  $u_k$  and  $v_k$  are not  $\bullet$ .

By definition of embedding,  $d(\phi(\mathbf{u}), \phi(\mathbf{v})) \leq 2^{-k} d(\mathbf{u}, \mathbf{v})$ .

Since  $|\phi'(\mathbf{u})| \leq 2^{-k}$  for all  $k$ , it is zero.



Complete characterization

## Characterization result :

**Theorem**[Gangloff, Oprocha] : A Cantor system can be embedded in  $\mathbb{R}$  with vanishing derivative if and only if all its finite orbits are attractors.



## Finite orbits and attractors in the graph coverings :

Finite orbit :





## Finite orbits and attractors in the graph coverings :

Finite orbit :



## Finite orbits and attractors in the graph coverings :

Finite orbit :





**Attractor** : closed set  $C$  such that for an open neighborhood  $U$  of  $C$ ,  $\bigcap_n f^n(U) = C$ .




## Finite orbits and attractors in the graph coverings :

Finite orbit :



**Attractor** : closed set  $C$  such that for an open neighborhood  $U$  of  $C$ ,  $\bigcap_n f^n(U) = C$ .



## Embeddable $\Rightarrow$ attracting finite orbits :

1. If  $g : \mathbb{R} \rightarrow \mathbb{R}$  is such that on a finite orbit  $p$ ,  $g'|_p \equiv 0$ ,  $|g'| \leq 1/2$  on a neighborhood of  $p$ .

## Embeddable $\Rightarrow$ attracting finite orbits :

1. If  $g : \mathbb{R} \rightarrow \mathbb{R}$  is such that on a finite orbit  $p$ ,  $g'|_p \equiv 0$ ,  $|g'| \leq 1/2$  on a neighborhood of  $p$ .
2. As a consequence  $p$  is attractor.

## Embeddable $\Rightarrow$ attracting finite orbits :

1. If  $g : \mathbb{R} \rightarrow \mathbb{R}$  is such that on a finite orbit  $p$ ,  $g'|_p \equiv 0$ ,  $|g'| \leq 1/2$  on a neighborhood of  $p$ .
2. As a consequence  $p$  is attractor.
3. This has to be true for embeddable Cantor systems.

## *Supercyclical* partitions :

The partitions  $\mathcal{U}_n$  satisfy the following :

## *Supercyclical partitions :*


The partitions  $\mathcal{U}_n$  satisfy the following :

It is possible to form attractors for all length  $n$  orbits with unions of elements of  $\mathcal{U}_n$ .


## *Supercyclical partitions :*

The partitions  $\mathcal{U}_n$  satisfy the following :

It is possible to form attractors for all length  $n$  orbits with unions of elements of  $\mathcal{U}_n$ .




## Rectification of finite attractors in supercyclical partitions :






## Marking and shrinking processes :


1.




2.



3.



How to deal with finite attractors : distance to the orbit :



How to deal with finite attractors : choosing intervals :

