



# Entropy of quasiperiodic subshifts


Guilhem Gamard

26th October 2018


# Tilings and crystals

# Tiles







# Crystals




# Penrose tiles






# Quasicrystals









What's the physical meaning of Penrose tiles?

# Gummelt's decagon





# Gummelt $\simeq$ Penrose

## Gummelt tiling rule

Tiles may overlap if decorations match; each tile must overlap.

## Theorem (Gummelt, 1996)

*Each Gummelt-tiling is isomorphic to a Penrose-tiling, and vice-versa.*

## Physical interpretation


Gummelt decagon has locally minimal energy.

**Back to subshifts**

Let  $q$  denote a pattern.

## Quasiperiodic


A configuration has *quasiperiod*  $q$  when it is covered with copies of  $q$  (possibly overlapping).



Let  $q$  denote a pattern.

## Quasiperiodic


A configuration has *quasiperiod*  $q$  when it is covered with copies of  $q$  (possibly overlapping).



Let  $q$  denote a pattern.

## Quasiperiodic

A configuration has *quasiperiod*  $q$  when it is covered with copies of  $q$  (possibly overlapping).





### Lemma

*The set of  $q$ -quasiperiodic configurations is an SFT, called  $X_q$ .*

What is the entropy of  $X_q$ ?

## The case of $\mathbb{Z}$ -subshifts

Let  $q$  denote a finite word.

### Remark

Consider two overlapping copies of  $q$ ; the overlap is a prefix and suffix of  $q$ .



Let  $q$  denote a finite word.

### Remark

Consider two overlapping copies of  $q$ ; the overlap is a prefix and suffix of  $q$ .



### Border

A **border** is a proper suffix and prefix.

An **antiborder** is the right-complement of a border.

(Note:  $\varepsilon$  is a border and  $q$  is an antiborder.)

Let  $q$ ,  $w$  denote finite words.

### Theorem (Mouchard, 2000)

*The word  $w$  has quasiperiod  $q$  iff  $w = qu_0 \dots u_k$ , where each  $u_i$  is an antiborder of  $q$ .*

Let  $q, w$  denote finite words.

### Theorem (Mouchard, 2000)

*The word  $w$  has quasiperiod  $q$  iff  $w = qu_0 \dots u_k$ , where each  $u_i$  is an antiborder of  $q$ .*

*This decomposition is unique iff  $q$  is **not** quasiperiodic.*

Let  $q, w$  denote finite words.

### Theorem (Mouchard, 2000)

*The word  $w$  has quasiperiod  $q$  iff  $w = qu_0 \dots u_k$ , where each  $u_i$  is an antiborder of  $q$ .*

*This decomposition is unique iff  $q$  is **not** quasiperiodic.*

### Corollary

The biinfinite word  $\mathbf{w}$  has quasiperiod  $q$  iff  $\mathbf{w} = \dots u_{-2} u_{-1} u_0 u_1 u_2 \dots$  where each  $u_i$  is an antiborder of  $q$ .

This decomposition is unique iff  $q$  is **not** quasiperiodic.

Fix  $q$  a quasiperiod. Let  $\ell(n) = \#$   $q$ -quasiperiodic words of length  $n$ .

Let  $r_0, \dots, r_{k-1}$  be the antiborders of  $q$ .

- If  $n < |q|$ , then  $\ell(n) = 0$ .
- If  $n = |q|$ , then  $\ell(n) = 1$ .
- If  $n > |q|$ , then  $\ell(n) = \sum_{i=0}^{k-1} \ell(n - |r_i|)$ .



Fix  $q$  a quasiperiod. Let  $\ell(n) = \#$   $q$ -quasiperiodic words of length  $n$ .

Let  $r_0, \dots, r_{k-1}$  be the antiborders of  $q$ .

- If  $n < |q|$ , then  $\ell(n) = 0$ .
- If  $n = |q|$ , then  $\ell(n) = 1$ .
- If  $n > |q|$ , then  $\ell(n) = \sum_{i=0}^{k-1} \ell(n - |r_i|)$ .

Let

$$P(x) = x^{|q|} - \sum_{i=0}^{k-1} x^{|q| - |r_i|}$$

and  $\lambda$  the largest real root of  $P$ .

### Lemma

For large  $n$ :  $c_1 \lambda^n \leq \ell(n) \leq c_2 \lambda^n$ .

## Theorem (Polley, Staiger, 2010)

Let  $q$  denote a non-quasiperiodic word with antiborders  $r_0, \dots, r_{k-1}$ , and

$$P(x) = x^{|q|} - \sum_{i=0}^{k-1} x^{|q|-|r_i|}$$

If  $\lambda$  is the largest root of  $P$ , then  $\boxed{\text{Ent}(X_q) = \log(\lambda)}$  .

### Theorem (Polley, Staiger, 2010)

Let  $q$  denote a non-quasiperiodic word with antiborders  $r_0, \dots, r_{k-1}$ , and

$$P(x) = x^{|q|} - \sum_{i=0}^{k-1} x^{|q|-|r_i|}$$

If  $\lambda$  is the largest root of  $P$ , then  $\boxed{Ent(X_q) = \log(\lambda)}$  .

### Remark

$X_q$  is finite iff  $q$  has no border besides  $\varepsilon$ .

### Theorem (Polley, Staiger, 2010)

Let  $q$  denote a non-quasiperiodic word with antiborders  $r_0, \dots, r_{k-1}$ , and

$$P(x) = x^{|q|} - \sum_{i=0}^{k-1} x^{|q|-|r_i|}$$

If  $\lambda$  is the largest root of  $P$ , then  $\boxed{\text{Ent}(X_q) = \log(\lambda)}$  .

### Remark

$X_q$  is finite iff  $q$  has no border besides  $\varepsilon$ . Otherwise,  $\text{Ent}(X_q) > 0$ .

### Theorem (Polley, Staiger, 2010)

Let  $q$  denote a non-quasiperiodic word with antiborders  $r_0, \dots, r_{k-1}$ , and

$$P(x) = x^{|q|} - \sum_{i=0}^{k-1} x^{|q|-|r_i|}$$

If  $\lambda$  is the largest root of  $P$ , then  $\boxed{\text{Ent}(X_q) = \log(\lambda)}$  .

### Remark

$X_q$  is finite iff  $q$  has no border besides  $\varepsilon$ . Otherwise,  $\text{Ent}(X_q) > 0$ .

### Remark


$\text{Ent}(X_q)$  is maximal for  $q = 010$ .

## The case of $\mathbb{Z}^2$ -subshifts

We consider **rectangular** patterns only.

## Root


The block  $r$  is a **root** of  $q$  iff  $q = r^{m \times n}$  for  $m, n \in \mathbb{N}$ .



We consider **rectangular** patterns only.

## Root

The block  $r$  is a **root** of  $q$  iff  $q = r^{m \times n}$  for  $m, n \in \mathbb{N}$ .






We consider **rectangular** patterns only.

## Root

The block  $r$  is a **root** of  $q$  iff  $q = r^{m \times n}$  for  $m, n \in \mathbb{N}$ .




## Primitive

The block  $q$  is **primitive** if it has no roots besides itself.

We consider **rectangular** patterns only.

## Root

The block  $r$  is a **root** of  $q$  iff  $q = r^{m \times n}$  for  $m, n \in \mathbb{N}$ .



## Primitive


The block  $q$  is **primitive** if it has no roots besides itself.

## Lemma (G, Richomme, 2015)

*Each block has a unique primitive root.*


## Border

A **border** of  $q$  is a subblock that appears in two opposite corners.



## Border

A **border** of  $q$  is a subblock that appears in two opposite corners.



## Border

A **border** of  $q$  is a subblock that appears in two opposite corners.

## Remark


$$\{\text{borders of the primitive root}\} \simeq \{\text{overlaps}\}$$

## Border

A **border** of  $q$  is a subblock that appears in two opposite corners.

## Remark

$\{\text{borders of the primitive root}\} \simeq \{\text{overlaps}\}$




## Border

A **border** of  $q$  is a subblock that appears in two opposite corners.

## Remark

$\{\text{borders of the primitive root}\} \simeq \{\text{overlaps}\}$




## Border

A **border** of  $q$  is a subblock that appears in two opposite corners.

## Remark

$\{\text{borders of the primitive root}\} \simeq \{\text{overlaps}\}$





Let  $q$  denote a block and  $r$  its primitive root.

Theorem (G, Richomme, 2015)

*The shift  $X_q$  is finite iff the primitive root of  $q$  has no border besides  $\varepsilon$ .*

Let  $q$  denote a block and  $r$  its primitive root.

**Theorem (G, Richomme, 2015)**

*The shift  $X_q$  is finite iff the primitive root of  $q$  has no border besides  $\varepsilon$ .*

- If  $r$  has no border besides  $\varepsilon$ , then  $X_q = r^{\mathbb{Z}^2}$ .

Let  $q$  denote a block and  $r$  its primitive root.

**Theorem (G, Richomme, 2015)**

*The shift  $X_q$  is finite iff the primitive root of  $q$  has no border besides  $\varepsilon$ .*


- If  $r$  has no border besides  $\varepsilon$ , then  $X_q = r^{\mathbb{Z}^2}$ .
- If  $r$  has a nonempty border, then build tiles.


Let  $q$  denote a block and  $r$  its primitive root.

### Theorem (G, Richomme, 2015)


*The shift  $X_q$  is finite iff the primitive root of  $q$  has no border besides  $\varepsilon$ .*

- If  $r$  has no border besides  $\varepsilon$ , then  $X_q = r^{\mathbb{Z}^2}$ .
- If  $r$  has a nonempty border, then build tiles.






Tiles have local constraints:



## Problem

Those constraints do not allow positive entropy!

We can prove that  $X_q$  is infinite, but not that  $\text{Ent}(X_q) > 0$ .

Is it bad? Consider the block  $q =$  .

### Proposition


$X_q$  is infinite, but has zero entropy.

Is it bad? Consider the block  $q =$




### Proposition

$X_q$  is infinite, but has zero entropy.









Is it bad? Consider the block  $q =$ 
.

## Proposition

$X_q$  is infinite, but has zero entropy.




Is it bad? Consider the block  $q =$




## Proposition

$X_q$  is infinite, but has zero entropy.



## Mindmap so far



## Interchangeable pairs

An **interchangeable pair** is a pair of  $q$ -quasiperiodic patterns, with the same shape, but different.




## Valid

A pair is **valid** if it appears in a configuration of  $X_q$ .

## Lemma

*If there is an interchangeable pair whose size is a  $k \times k$  square, then*

$$Ent(X_q) \geq 1/k$$



## Lemma

*If there is no interchangeable pair for  $q$ , then  $\text{Ent}(X_q) = 0$ .*

**Idea:** if there are no interchangeable pair, then the shape of a pattern defines the contents of the pattern.

### Lemma

*If there is no interchangeable pair for  $q$ , then  $\text{Ent}(X_q) = 0$ .*

**Idea:** if there are no interchangeable pair, then the shape of a pattern defines the contents of the pattern.

### Theorem

*We have  $\text{Ent}(X_q) > 0$  if and only if there is a valid pair for  $q$ .*

(Note: this is also true for Wang tiles.)

## Proposition


If  $q$  is of the form 

$a$		$b$
$b$		$a$

 or 

$a$	$b$
$b$	$a$

 then  $\text{Ent}(X_q) > 0$ .





## Proposition


If  $q$  is of the form 


$a$		$b$
$b$		$a$

 or 


$a$	$b$
$b$	$a$

 then  $\text{Ent}(X_q) > 0$ .






## Proposition

$\text{Ent}(X_q) = 0$  unless  $q$  is of the form .

**Idea:** on the blackboard.

## Proposition


$\text{Ent}(X_q) = 0$  unless  $q$  is of the form .

**Idea:** on the blackboard.

## Special case


If all borders of  $q$  are in the same two corners, then  $\text{Ent}(X_q) = 0$ .

## Recap




nonzero entropy

**Not**




zero entropy

## Recap



nonzero entropy

**Not**



zero entropy

Rather frustrating. (Help appreciated.)

**What's next?**

## What's next?

- Finish the characterization
- Exact values (or more precise bounds)
- Extend to other shapes
- Relax the definition (pre-quasiperiodic?)



## What's next?

- Finish the characterization
- Exact values (or more precise bounds)
- Extend to other shapes
- Relax the definition (pre-quasiperiodic?)

Thank you!

# Bibliography

- D. Shechtman, I. Blech, D. Gratias, J. Cahn.  
Metallic Phase with Long-Range Orientational Order and No Translational Symmetry.  
Physical Review Letters 53(20), pp. 1951–1953, 1984.  
<https://link.aps.org/doi/10.1103/PhysRevLett.53.1951>
- Petra Gummelt.  
Penrose tilings as coverings of congruent decagons.  
Geometriae Dedicata 62, pp. 1–17, 1996.  
<https://link.springer.com/article/10.1007/BF00239998>
- Thierry Monteil, Solomon Marcus.  
Quasiperiodic infinite words: multi-scale case and dynamical properties.  
<https://arxiv.org/abs/math/06033542006>
- Ronney Polley, Ludwig Staiger.  
Quasiperiods, Subword Complexity and the Smallest Pisot Number.  
Journal of Automata, Languages and Combinatorics 21(1-2), pp. 93–106, 2016.  
[http://jal.c.de/issues/2016/issue\\_21\\_1-2/jalc-2016-093-106.php](http://jal.c.de/issues/2016/issue_21_1-2/jalc-2016-093-106.php)
- Guilhem Gamard, Gwenaél Richomme.  
Coverability and Multi-Scale Coverability on infinite pictures.  
To appear in Journal of Computer and System Sciences.  
<https://arxiv.org/abs/1506.08375>