

On exact computation of square ice entropy


Silvère Gangloff

LIP, ENS Lyon


March 11, 2019

I. Representations of square ice


Square ice model [Pauling-Lieb]:




Wang tiles representation [Six-vertex model]




Wang tiles representation [Six-vertex model]




Wang tiles representation [Six-vertex model]





Wang tiles representation [Six-vertex model]



Discrete curves subshift $[X^s]$:



Discrete curves subshift $[X^S]$:



Example of apparition in symbolic dynamics:

S.Gangloff, M. Sablik, *Quantified block gluing, aperiodicity and entropy of multidimensional SFT* , 2017.

Example of apparition in symbolic dynamics:

S.Gangloff, M. Sablik, *Quantified block gluing, aperiodicity and entropy of multidimensional SFT* , 2017.

Entropy value ?

E.H. Lieb, *Residual entropy of square ice*, Physical Review, 1967.

Example of apparition in symbolic dynamics:

S.Gangloff, M. Sablik, *Quantified block gluing, aperiodicity and entropy of multidimensional SFT*, 2017.

Entropy value ?

E.H. Lieb, *Residual entropy of square ice*, Physical Review, 1967.

→ **Proof under some hypothesis**

Example of apparition in symbolic dynamics:

S.Gangloff, M. Sablik, *Quantified block gluing, aperiodicity and entropy of multidimensional SFT*, 2017.

Entropy value ?

E.H. Lieb, *Residual entropy of square ice*, Physical Review, 1967.

→ **Proof under some hypothesis**

S. Gangloff, *A proof that square ice entropy is $\frac{3}{2} \log_2(4/3)$* , 2019.

II. Subshifts of finite type and entropy

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

Ex: **Hard square shift**, or hard core model.

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

Ex: **Hard square shift**, or hard core model.

Forbidden patterns $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ et $\begin{bmatrix} 1 & 1 \end{bmatrix}$.

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

Ex: **Hard square shift**, or hard core model.

Forbidden patterns $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ et $\begin{bmatrix} 1 & 1 \end{bmatrix}$.

0	0	0	0	0
0	0	0	1	0
0	0	0	0	1
0	0	0	1	0
1	0	0	0	0

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

Ex: **Hard square shift**, or hard core model.

Forbidden patterns $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ et $\begin{bmatrix} 1 & 1 \end{bmatrix}$.

0	0	0	0	0
0	0	0	1	0
0	0	0	0	1
1	0	0	1	0
0	0	0	0	0

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

Ex: **Hard square shift**, or hard core model.

Forbidden patterns $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ et $\begin{bmatrix} 1 & 1 \end{bmatrix}$.

0	0	0	0	0
0	0	0	1	0
0	0	0	0	1
1	0	1	0	0
0	0	0	0	0

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

Ex: **Hard square shift**, or hard core model.

Forbidden patterns $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ et $\begin{bmatrix} 1 & 1 \end{bmatrix}$.

0	0	0	0	0	
0	0	0	0	0	
0	0	0	1	1	oops
1	0	1	0	0	
0	0	0	0	0	

SFT (subshift of finite type): subset of $\mathcal{A}^{\mathbb{Z}^2}$, defined by a finite set of **forbidden patterns**.

Ex: **Hard square shift**, or hard core model.

Forbidden patterns $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ et $\begin{bmatrix} 1 & 1 \end{bmatrix}$.

0	0	0	0	0
0	0	0	1	0
0	0	0	0	1
1	0	1	0	0
0	0	0	0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	0
0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	0
1	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	0
0	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	0
0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	1
0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	1
1	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	0
0	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	0
1	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	1
0	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	0
1	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	0
1	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

0	1
1	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
0	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

0	0
0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

0	0
1	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

0	0
0	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

1	0
0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

0	1
0	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

0	1
1	0

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

1	0
0	1

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

1	0
0	1

$$\mathcal{N}_2(X) = 7$$

Entropy: "quantity of possible states of the system".

$\mathcal{N}_N(X)$: number of size N square patterns observable in the system.

Free tiles

1	1
1	1

$$\mathcal{N}_2(X) = 2^{2^2}$$

$$\mathcal{N}_N(X) = 2^{N^2}$$

Hard core

1	0
0	1

$$\mathcal{N}_2(X) = 7$$

$$\mathcal{N}_N(X) = 2^{N^2(h(X)+o(1))}$$

Entropy of a SFT X :

$$h(X) = \inf_N \frac{\log_2(\mathcal{N}_N(X))}{N^2}$$

Free tiles

$$h = 1$$

Hard core

$$h \geq 1/2$$

Square ice [Lieb 67]

$$h = \frac{3}{2} \log_2(4/3)$$

Entropy of a SFT X :

$$h(X) = \inf_N \frac{\log_2(\mathcal{N}_N(X))}{N^2} = \inf_N \frac{\log_2(\mathcal{N}_N^{loc}(X))}{N^2}$$

Free tiles

$$h = 1$$

Hard core

$$h \geq 1/2$$

Square ice [Lieb 67]

$$h = \frac{3}{2} \log_2(4/3)$$

Entropy of a SFT X :

$$h(X) = \inf_N \frac{\log_2(\mathcal{N}_N(X))}{N^2} = \inf_N \frac{\log_2(\mathcal{N}_N^{loc}(X))}{N^2}$$

Free tiles

$$h = 1$$


Hard core

$$h \geq 1/2$$

Square ice [Lieb 67]

$$h = \frac{3}{2} \log_2(4/3)$$

Computable from above:



Entropy of a SFT X :

$$h(X) = \inf_N \frac{\log_2(\mathcal{N}_N(X))}{N^2} = \inf_N \frac{\log_2(\mathcal{N}_N^{loc}(X))}{N^2}$$

Free tiles

$$h = 1$$


Hard core

$$h \geq 1/2$$

Square ice [Lieb 67]

$$h = \frac{3}{2} \log_2(4/3)$$

Computable from above:



Entropy of a SFT X :

$$h(X) = \inf_N \frac{\log_2(\mathcal{N}_N(X))}{N^2} = \inf_N \frac{\log_2(\mathcal{N}_N^{loc}(X))}{N^2}$$

Free tiles

$$h = 1$$


Hard core

$$h \geq 1/2$$

Square ice [Lieb 67]

$$h = \frac{3}{2} \log_2(4/3)$$

Computable from above:



Entropy of a SFT X :

$$h(X) = \inf_N \frac{\log_2(\mathcal{N}_N(X))}{N^2} = \inf_N \frac{\log_2(\mathcal{N}_N^{loc}(X))}{N^2}$$

Free tiles

$$h = 1$$


Hard core

$$h \geq 1/2$$

Square ice [Lieb 67]

$$h = \frac{3}{2} \log_2(4/3)$$

Computable from above:




II. Lieb transfer matrices approach

Entropy of square ice:


$$h(X^s) = \lim_{M,N} \frac{\log_2(\mathcal{N}_{M,N}(X^s))}{MN}.$$

Stripes subshifts:


$$h(X^s) = \lim_N \frac{h(X_N^s)}{N}$$



Cylindric stripes subshifts:




X_N^s



\overline{X}_N^s


Cylindric stripes subshifts:




Symmetry properties of square ice imply:

$$h(X^s) = \lim_N \frac{h(\overline{X}_N^s)}{N}$$


Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Symmetry properties of square ice:




Lieb transfer matrices:




Lieb transfer matrices:




Lieb transfer matrices:




Lieb transfer matrices:




Lieb transfer matrices:



Lieb transfer matrices:




Lieb transfer matrices:



$$V_N(t)[\mathbf{u}, \mathbf{v}] = \sum_{\mathbf{u} \mathcal{R}[w] \mathbf{v}} t^{|w|}.$$

where $|w| = \#$ of $\begin{smallmatrix} \square \\ \square \end{smallmatrix}$ and $\begin{smallmatrix} \square \\ \square \end{smallmatrix}$

Lieb transfer matrices:



$$V_N(t)[\mathbf{u}, \mathbf{v}] = \sum_{\mathbf{u} \mathcal{R}[w] \mathbf{v}} t^{|w|}.$$


where $|w| = \#$ of $\begin{smallmatrix} \square \\ \square \end{smallmatrix}$ and $\begin{smallmatrix} \square \\ \square \end{smallmatrix}$

$$h(X^s) = \lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N}$$


Computing maximal eigenvalue of $V_N(1)$, strategy:

$$V_N(1) \quad \bullet$$


Computing maximal eigenvalue of $V_N(1)$, strategy:




Computing maximal eigenvalue of $V_N(1)$, strategy:




Computing maximal eigenvalue of $V_N(1)$, strategy:




Computing maximal eigenvalue of $V_N(1)$, strategy:



Computing maximal eigenvalue of $V_N(1)$, strategy:



Computing maximal eigenvalue of $V_N(1)$, strategy:



III. Yang-Baxter transfer matrices


R-matrices and monodromy matrices:

R-matrices and monodromy matrices: \rightarrow : input/output:

R-matrices and monodromy matrices: \rightarrow : input/output:




R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \left(\begin{array}{c} \\ \end{array} \right)$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \\ & \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ & 1 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 1 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:





$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:





$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:





$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:





$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:





$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$

R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$


R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$



R-matrices and monodromy matrices: \rightarrow : input/output:




$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$




$$M_N(1,0)[\epsilon, \eta]$$

R-matrices and monodromy matrices: \rightarrow : input/output:



$$R(0,1) = \begin{pmatrix} 0 & \lambda \\ 0 & 0 \end{pmatrix}$$





$$M_N(1,0)[\epsilon, \eta]$$


Yang-Baxter transfer matrices:

$$T_N[\epsilon, \eta] = \sum_{u \in \{0,1\}} M_N(u, u)[\epsilon, \eta].$$


Composition of these matrices and condition for commutation:



Composition of these matrices and condition for commutation:



Yang-Baxter equation:



Trigonometric R -matrices: for $t \in (0, \sqrt{2}]$, $2 - t^2 = -\cos(\mu_t)$:

Trigonometric R -matrices: for $t \in (0, \sqrt{2}]$, $2 - t^2 = -\cos(\mu_t)$:

$$R_{\mu_t}^x = \frac{1}{\sin(\mu_t/2)} \begin{pmatrix} \sin(\mu_t - x) & 0 & 0 & 0 \\ 0 & \sin(x) & \sin(\mu_t) & 0 \\ 0 & \sin(\mu_t) & \sin(x) & 0 \\ 0 & 0 & 0 & \sin(\mu_t - x) \end{pmatrix}.$$

Trigonometric R -matrices: for $t \in (0, \sqrt{2}]$, $2 - t^2 = -\cos(\mu_t)$:

$$R_{\mu_t}^x = \frac{1}{\sin(\mu_t/2)} \begin{pmatrix} \sin(\mu_t - x) & 0 & 0 & 0 \\ 0 & \sin(x) & \sin(\mu_t) & 0 \\ 0 & \sin(\mu_t) & \sin(x) & 0 \\ 0 & 0 & 0 & \sin(\mu_t - x) \end{pmatrix}.$$

Bethe ansatz: if $(p_j)_j$ is solution of:

$$Np_j = 2\pi j - (n+1)\pi - \sum_{k=1}^n \Theta_t(p_j, p_k)$$

Trigonometric R -matrices: for $t \in (0, \sqrt{2}]$, $2 - t^2 = -\cos(\mu_t)$:

$$R_{\mu_t}^x = \frac{1}{\sin(\mu_t/2)} \begin{pmatrix} \sin(\mu_t - x) & 0 & 0 & 0 \\ 0 & \sin(x) & \sin(\mu_t) & 0 \\ 0 & \sin(\mu_t) & \sin(x) & 0 \\ 0 & 0 & 0 & \sin(\mu_t - x) \end{pmatrix}.$$

Bethe ansatz: if $(p_j)_j$ is solution of:

$$Np_j = 2\pi j - (n+1)\pi - \sum_{k=1}^n \Theta_t(p_j, p_k)$$

then we have a candidate eigenvector for the eigenvalue:

$$\prod_{k=1}^n L_t(e^{ip_k}) + \prod_{k=1}^n M_t(e^{ip_k}).$$

Trigonometric R -matrices: for $t \in (0, \sqrt{2}]$, $2 - t^2 = -\cos(\mu_t)$:

$$R_{\mu_t}^x = \frac{1}{\sin(\mu_t/2)} \begin{pmatrix} \sin(\mu_t - x) & 0 & 0 & 0 \\ 0 & \sin(x) & \sin(\mu_t) & 0 \\ 0 & \sin(\mu_t) & \sin(x) & 0 \\ 0 & 0 & 0 & \sin(\mu_t - x) \end{pmatrix}.$$

Bethe ansatz: if $(p_j)_j$ is solution of:

$$Np_j = 2\pi j - (n+1)\pi - \sum_{k=1}^n \Theta_t(p_j, p_k)$$

then we have a candidate eigenvector for the eigenvalue:

$$\prod_{k=1}^n L_t(e^{ip_k}) + \prod_{k=1}^n M_t(e^{ip_k}).$$

→ **Known:** existence and analyticity in t .

Identification to $\lambda_{\max}(V_N(t))$: ingredients:

Identification to $\lambda_{\max}(V_N(t))$: ingredients:

- ① Simplification of equations when $t = \sqrt{2}$.

Identification to $\lambda_{\max}(V_N(t))$: ingredients:

- ① Simplification of equations when $t = \sqrt{2}$.
- ② For some H_N diagonalised, $V_N(\sqrt{2})H_N = H_N V_N(\sqrt{2})$.

Identification to $\lambda_{\max}(V_N(t))$: ingredients:

- ① Simplification of equations when $t = \sqrt{2}$.
- ② For some H_N diagonalised, $V_N(\sqrt{2})H_N = H_N V_N(\sqrt{2})$.
- ③ Perron-Frobenius theorem.

Identification to $\lambda_{\max}(V_N(t))$: ingredients:

- ① Simplification of equations when $t = \sqrt{2}$.
- ② For some H_N diagonalised, $V_N(\sqrt{2})H_N = H_N V_N(\sqrt{2})$.
- ③ Perron-Frobenius theorem.
- ④ Identification around $\sqrt{2}$, on $(0, \sqrt{2})$ by analycity.

IV. Asymptotics

Asymptotics of counting functions:

Asymptotics of counting functions: $n_k/N_k \rightarrow 1/2$.


Asymptotics of counting functions: $n_k/N_k \rightarrow 1/2$. Solution of the equations for N_k, n_k : $(\mathbf{p}_j^{(k)})_j = (\kappa_t(\boldsymbol{\alpha}_j^{(k)}))_j$.

Asymptotics of counting functions: $n_k/N_k \rightarrow 1/2$. Solution of the equations for N_k, n_k : $(\mathbf{p}_j^{(k)})_j = (\kappa_t(\boldsymbol{\alpha}_j^{(k)}))_j$.

$$\xi_t^{(k)} : \alpha \mapsto \frac{1}{2\pi} \kappa_t(\alpha) + \frac{n_k + 1}{2N_k} + \frac{1}{2\pi N_k} \sum_{j=1}^{n_k} \theta_t(\alpha, \boldsymbol{\alpha}_j^{(k)})$$


Asymptotics of counting functions: $n_k/N_k \rightarrow 1/2$. Solution of the equations for N_k, n_k : $(\mathbf{p}_j^{(k)})_j = (\kappa_t(\boldsymbol{\alpha}_j^{(k)}))_j$.

$$\xi_t^{(k)} : \alpha \mapsto \frac{1}{2\pi} \kappa_t(\alpha) + \frac{n_k + 1}{2N_k} + \frac{1}{2\pi N_k} \sum_{j=1}^{n_k} \theta_t(\alpha, \alpha_j^{(k)})$$



Asymptotics of counting functions: $n_k/N_k \rightarrow 1/2$. Solution of the equations for N_k, n_k : $(\mathbf{p}_j^{(k)})_j = (\kappa_t(\boldsymbol{\alpha}_j^{(k)}))_j$.

$$\xi_t^{(k)} : \alpha \mapsto \frac{1}{2\pi} \kappa_t(\alpha) + \frac{n_k + 1}{2N_k} + \frac{1}{2\pi N_k} \sum_{j=1}^{n_k} \theta_t(\alpha, \boldsymbol{\alpha}_j^{(k)})$$



$$\lim_N \frac{\log_2(\lambda_{\max}(V_N(1)))}{N} = \lim_k \frac{1}{N_k} \sum_{j=1}^{n_k} f(\boldsymbol{\alpha}_j^{(k)}) = \int_{\mathbb{R}} f(\alpha) \rho_t(\alpha) d\alpha.$$

Strategy:

- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :


Strategy:

- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :




Strategy:

- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :




Strategy:

- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :



Strategy:


- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :



- 2 Assume $(\xi_t)^{\nu(k)} \rightarrow \xi_t^{(+\infty)}$ on any compact K .

Strategy:


- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :



- 2 Assume $(\xi_t)^{\nu(k)} \rightarrow \xi_t^{(+\infty)}$ on any compact K .

Strategy:


- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :



- 2 Assume $(\xi_t)^{\nu(k)} \rightarrow \xi_t^{(+\infty)}$ on any compact K .

Strategy:


- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :



- 2 Assume $(\xi_t)^{\nu(k)} \rightarrow \xi_t^{(+\infty)}$ on any compact K .

Strategy:


- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :



- 2 Assume $(\xi_t)^{\nu(k)} \rightarrow \xi_t^{(+\infty)}$ on any compact K .

Strategy:


- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :



- 2 Assume $(\xi_t)^{\nu(k)} \rightarrow \xi_t^{(+\infty)}$ on any compact K .
- 3 $\xi_t^{(+\infty)}$ verifies an integral equation with unique solution ρ_t .


Strategy:

- 1 Extend $\xi_t^{(k)}$ on a stripe including \mathbb{R} :




- 2 Assume $(\xi_t)^{\nu(k)} \rightarrow \xi_t^{(+\infty)}$ on any compact K .
- 3 $\xi_t^{(+\infty)}$ verifies an integral equation with unique solution ρ_t .
- 4 Thus, $\xi_t^{(k)} \rightarrow \xi_t^{(\infty)}$.


Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:




Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:




Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:




Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:




Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:




Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:



Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:




Rarefaction of the roots and $\xi_t^{(k)}$ biholomorphisms: $\epsilon > 0$:




The functions have distinct values on \mathcal{V} and Γ . Thus they are biholomorphisms onto \mathcal{V} .

Lace integral expression of $\xi_t^{(k)}$:



Lace integral expression of $\xi_t^{(k)}$:



By residues theorem:

$$\xi_t^{(k)}(\alpha) = \frac{1}{2\pi} \kappa_t(\alpha) + \frac{n_k + 1}{2N_k} + \oint_{\Gamma} \theta_t \left(\left(\xi_t^{(k)} \right)^{-1}(\alpha) \right) \frac{e^{2i\pi s N_k}}{e^{2i\pi s N_k} - 1} ds + O(\epsilon).$$

Fredholm integral equation: Limit and change of variable:

$$\xi_t^{(\infty)}(\alpha) = \frac{1}{2\pi} \kappa_t(\alpha) + \frac{1}{4} + \int_0^{+\infty} \theta_t(\alpha) \left(\xi_t^{(\infty)} \right)'(\alpha) d\alpha.$$

Fredholm integral equation: Limit and change of variable:

$$\xi_t^{(\infty)}(\alpha) = \frac{1}{2\pi} \kappa_t(\alpha) + \frac{1}{4} + \int_0^{+\infty} \theta_t(\alpha) \left(\xi_t^{(\infty)} \right)'(\alpha) d\alpha.$$

Solution by Fourier transforms.

Final computation:

Final computation:

$$h(X^s) = \int_{\mathbb{R}} \log_2(2|\sin(\kappa_t(\alpha))/2|) \cdot \rho_t(\alpha) d\alpha.$$

Final computation:


$$h(X^s) = \int_{\mathbb{R}} \log_2(2|\sin(\kappa_t(\alpha))/2|) \cdot \rho_t(\alpha) d\alpha.$$

Through an expression of $\rho_t = \left(\xi_t^{(\infty)}\right)'$ and lace integrals computations:

$$h(X^s) = \frac{3}{2} \log_2(4/3).$$

V. Comments

Why mathematical physics are hard to read for mathematicians?



Archaeology of Knowledge, 1969

Concept of *discursive formation*

Mathematics and mathematical physics are distinct discursive formations ;
different conceptions of units of meaning, etc.

Further research:

- 1 Extensions: eight-vertex model [Baxter], dimer model [Lieb]..



Further research:

- 1 Extensions: eight-vertex model [Baxter], dimer model [Lieb]..



- 2 **Hard core model** ? Tridimensional ice ? Kari-Culik tilings ?



Further research:

- 1 Extensions: eight-vertex model [Baxter], dimer model [Lieb]..



- 2 **Hard core model** ? Tridimensional ice ? Kari-Culik tilings ?



- 3 Transformation of entropy by subshifts operators ?

