

# Exact Inference in Bayesian Networks

---

## Outline

- I. Probabilistic query using a BN
- II. Variable Elimination
- III. Variable ordering and relevance

# I. Probabilistic Query

---

$X$ : query variable

$E = \{E_1, \dots, E_m\}$ : evidence variables

$e = \{e_1, \dots, e_m\}$ : an observed event

$Y = \{Y_1, \dots, Y_l\}$ : hidden variables

# I. Probabilistic Query

---

$X$ : query variable

$E = \{E_1, \dots, E_m\}$ : evidence variables

$e = \{e_1, \dots, e_m\}$ : an observed event

$Y = \{Y_1, \dots, Y_l\}$ : hidden variables

Complete variable set:  $\{X\} \cup E \cup Y$

# I. Probabilistic Query

---

$X$ : query variable

$E = \{E_1, \dots, E_m\}$ : evidence variables

$e = \{e_1, \dots, e_m\}$ : an observed event

$Y = \{Y_1, \dots, Y_l\}$ : hidden variables

Complete variable set:  $\{X\} \cup E \cup Y$

**Query**  $P(X \mid e)$ ?

# I. Probabilistic Query

---

$X$ : query variable

$E = \{E_1, \dots, E_m\}$ : evidence variables

$e = \{e_1, \dots, e_m\}$ : an observed event

$Y = \{Y_1, \dots, Y_l\}$ : hidden variables

Complete variable set:  $\{X\} \cup E \cup Y$

**Query**  $P(X \mid e)$ ?

//  $P(\text{Burglary} \mid j, m)$

$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true}) = \langle 0.284, 0.716 \rangle$

# I. Probabilistic Query

---

$X$ : query variable

$E = \{E_1, \dots, E_m\}$ : evidence variables

$e = \{e_1, \dots, e_m\}$ : an observed event

$Y = \{Y_1, \dots, Y_l\}$ : hidden variables

Complete variable set:  $\{X\} \cup E \cup Y$

**Query**  $P(X \mid e)$ ?

//  $P(\text{Burglary} \mid j, m)$

$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true}) = \langle 0.284, 0.716 \rangle$

We will discuss exact algorithms for posterior probability computation.

# Inference by Enumeration

---

$$P(X | e) = \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$

# Inference by Enumeration

---

$$P(X | e) = \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{Parents}(X_i))$$



# Inference by Enumeration

---

$$P(X | e) = \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{Parents}(X_i))$$



computable as products of  
conditional probabilities from the BN.

# Inference by Enumeration

---

$$P(X | e) = \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$


$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{Parents}(X_i))$$



computable as products of  
conditional probabilities from the BN.

- ♦ Answer the query  $P(X | e)$  using a BN.


# Burglary Example (revisited)



**$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true})?$**

Hidden variables:  $Y = \{\text{Earthquake}, \text{Alarm}\}$

# Burglary Example (revisited)




$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true})?$

Hidden variables:  $Y = \{\text{Earthquake}, \text{Alarm}\}$

$$P(B \mid j, m) = \alpha P(B, j, m) = \alpha \sum_{E \in \{e, \neg e\}} \sum_{A \in \{a, \neg a\}} P(B, j, m, E, A)$$

# Burglary Example (revisited)




$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true})?$

Hidden variables:  $Y = \{\text{Earthquake}, \text{Alarm}\}$

$$P(B \mid j, m) = \alpha P(B, j, m) = \alpha \sum_{E \in \{e, \neg e\}} \sum_{A \in \{a, \neg a\}} P(B, j, m, E, A)$$

// the textbook uses  $e, a$  instead of  $E, A$  in the summation.  
 // this is **incorrect** because  $e$  is a constant value standing  
 // for  $\text{Earthquake} = \text{true}$ . the random variable  $E$  can take  
 // on both  $e$  and  $\neg e$  needed for the summation. you may  
 // also call it an abuse of the notation  $e$  and  $a$ .

# Burglary Example (revisited)



$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true})?$

Hidden variables:  $Y = \{\text{Earthquake}, \text{Alarm}\}$


$$P(B \mid j, m) = \alpha P(B, j, m) = \alpha \sum_{E \in \{e, \neg e\}} \sum_{A \in \{a, \neg a\}} P(B, j, m, E, A)$$

evaluated for  $b$  ( $\text{Burglary} = \text{true}$ ) based on

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$$

// the textbook uses  $e, a$  instead of  $E, A$  in the summation.  
 // this is **incorrect** because  $e$  is a constant value standing  
 // for  $\text{Earthquake} = \text{true}$ . the random variable  $E$  can take  
 // on both  $e$  and  $\neg e$  needed for the summation. you may  
 // also call it an abuse of the notation  $e$  and  $a$ .

# Burglary Example (revisited)



$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true})?$

Hidden variables:  $Y = \{\text{Earthquake}, \text{Alarm}\}$

$$P(B \mid j, m) = \alpha P(B, j, m) = \alpha \sum_{E \in \{e, \neg e\}} \sum_{A \in \{a, \neg a\}} P(B, j, m, E, A)$$

evaluated for  $b$  ( $\text{Burglary} = \text{true}$ ) based on


$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$$



// the textbook uses  $e, a$  instead of  $E, A$  in the summation.  
 // this is **incorrect** because  $e$  is a constant value standing  
 // for  $\text{Earthquake} = \text{true}$ . the random variable  $E$  can take  
 // on both  $e$  and  $\neg e$  needed for the summation. you may  
 // also call it an abuse of the notation  $e$  and  $a$ .

$$P(b \mid j, m) = \alpha \sum_E \sum_A P(b)P(E)P(A \mid b, E)P(j \mid A)P(m \mid A)$$

# Burglary Example (revisited)



$P(\text{Burglary} \mid \text{JohnCalls} = \text{true}, \text{MaryCalls} = \text{true})?$

Hidden variables:  $Y = \{\text{Earthquake}, \text{Alarm}\}$

$$P(B \mid j, m) = \alpha P(B, j, m) = \alpha \sum_{E \in \{e, \neg e\}} \sum_{A \in \{a, \neg a\}} P(B, j, m, E, A)$$

evaluated for  $b$  ( $\text{Burglary} = \text{true}$ ) based on

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$$



// the textbook uses  $e, a$  instead of  $E, A$  in the summation.  
 // this is **incorrect** because  $e$  is a constant value standing  
 // for  $\text{Earthquake} = \text{true}$ . the random variable  $E$  can take  
 // on both  $e$  and  $\neg e$  needed for the summation. you may  
 // also call it an abuse of the notation  $e$  and  $a$ .

$$P(b \mid j, m) = \alpha \sum_E \sum_A P(b)P(E)P(A \mid b, E)P(j \mid A)P(m \mid A)$$

In the general case with  $n$  variables, there are  $2^n$  summands, each as a product requires  $O(n)$  computation time.



# Expression Tree

---


Take advantage of the nested structure to move summations inwards as far as possible.

$$P(b \mid j, m) = \alpha \sum_E \sum_A P(b)P(E)P(A \mid b, E)P(j \mid A)P(m \mid A)$$

# Expression Tree

---


Take advantage of the nested structure to move summations inwards as far as possible.

$$P(b \mid j, m) = \alpha \sum_E \sum_A P(b)P(E)P(A \mid b, E)P(j \mid A)P(m \mid A)$$


# Expression Tree

---

Take advantage of the nested structure to move summations inwards as far as possible.

$$P(b \mid j, m) = \alpha \sum_E \sum_A P(b)P(E)P(A \mid b, E)P(j \mid A)P(m \mid A)$$



# Expression Tree

---

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$



# Expression Tree

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$




# Expression Tree

$$P(b | j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A | b, E) P(j | A) P(m | A)$$




# Expression Tree

$$P(b | j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A | b, E) P(j | A) P(m | A)$$



# Expression Tree

$$P(b | j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A | b, E) P(j | A) P(m | A)$$




$$P(b | j, m) = \alpha \times 0.00059224$$



# Expression Tree

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$




$$P(b \mid j, m) = \alpha \times 0.00059224$$

$$P(\neg b \mid j, m) = \alpha \times 0.0014919$$

# Expression Tree

$$P(b | j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A | b, E) P(j | A) P(m | A)$$



$$\left. \begin{array}{l} P(b | j, m) = \alpha \times 0.00059224 \\ P(\neg b | j, m) = \alpha \times 0.0014919 \end{array} \right\} \Rightarrow P(B | j, m) = \alpha \langle 0.00059224, 0.0014919 \rangle \approx \langle 0.284, 0.716 \rangle$$

# Enumeration Algorithm

**function** ENUMERATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayes net with variables  $vars$

$\mathbf{Q}(X) \leftarrow$  a distribution over  $X$ , initially empty

**for each** value  $x_i$  of  $X$  **do**

$\mathbf{Q}(x_i) \leftarrow$  ENUMERATE-ALL( $vars, \mathbf{e}_{x_i}$ )

where  $\mathbf{e}_{x_i}$  is  $\mathbf{e}$  extended with  $X = x_i$

**return** NORMALIZE( $\mathbf{Q}(X)$ )

**function** ENUMERATE-ALL( $vars, \mathbf{e}$ ) **returns** a real number

**if** EMPTY?( $vars$ ) **then return** 1.0


$V \leftarrow$  FIRST( $vars$ )

**if**  $V$  is an evidence variable with value  $v$  in  $\mathbf{e}$

**then return**  $P(v | \text{parents}(V)) \times \text{ENUMERATE-ALL}(\text{REST}(vars), \mathbf{e})$

**else return**  $\sum_v P(v | \text{parents}(V)) \times \text{ENUMERATE-ALL}(\text{REST}(vars), \mathbf{e}_v)$

where  $\mathbf{e}_v$  is  $\mathbf{e}$  extended with  $V = v$



# Enumeration Algorithm

**function** ENUMERATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayes net with variables  $vars$

$Q(X) \leftarrow$  a distribution over  $X$ , initially empty

**for each** value  $x_i$  of  $X$  **do**

$Q(x_i) \leftarrow$  ENUMERATE-ALL( $vars, \mathbf{e}_{x_i}$ )

where  $\mathbf{e}_{x_i}$  is  $\mathbf{e}$  extended with  $X = x_i$

**return** NORMALIZE( $Q(X)$ )

**function** ENUMERATE-ALL( $vars, \mathbf{e}$ ) **returns** a real number

**if** EMPTY?( $vars$ ) **then return** 1.0

$V \leftarrow$  FIRST( $vars$ )


**if**  $V$  is an evidence variable with value  $v$  in  $\mathbf{e}$

**then return**  $P(v | \text{parents}(V)) \times \text{ENUMERATE-ALL}(\text{REST}(vars), \mathbf{e})$

**else return**  $\sum_v P(v | \text{parents}(V)) \times \text{ENUMERATE-ALL}(\text{REST}(vars), \mathbf{e}_v)$

where  $\mathbf{e}_v$  is  $\mathbf{e}$  extended with  $V = v$

- Depth-first recursion



# Enumeration Algorithm

**function** ENUMERATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayes net with variables  $vars$

$Q(X) \leftarrow$  a distribution over  $X$ , initially empty

**for each** value  $x_i$  of  $X$  **do**

$Q(x_i) \leftarrow$  ENUMERATE-ALL( $vars, \mathbf{e}_{x_i}$ )

where  $\mathbf{e}_{x_i}$  is  $\mathbf{e}$  extended with  $X = x_i$

**return** NORMALIZE( $Q(X)$ )

**function** ENUMERATE-ALL( $vars, \mathbf{e}$ ) **returns** a real number

**if** EMPTY?( $vars$ ) **then return** 1.0

$V \leftarrow$  FIRST( $vars$ )


**if**  $V$  is an evidence variable with value  $v$  in  $\mathbf{e}$

**then return**  $P(v | \text{parents}(V)) \times \text{ENUMERATE-ALL}(\text{REST}(vars), \mathbf{e})$

**else return**  $\sum_v P(v | \text{parents}(V)) \times \text{ENUMERATE-ALL}(\text{REST}(vars), \mathbf{e}_v)$

where  $\mathbf{e}_v$  is  $\mathbf{e}$  extended with  $V = v$

- Depth-first recursion
- $O(2^n)$  time for  $n$  variables



# Enumeration Algorithm

**function** ENUMERATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayes net with variables  $vars$

$Q(X) \leftarrow$  a distribution over  $X$ , initially empty

**for each** value  $x_i$  of  $X$  **do**

$Q(x_i) \leftarrow$  ENUMERATE-ALL( $vars, \mathbf{e}_{x_i}$ )

where  $\mathbf{e}_{x_i}$  is  $\mathbf{e}$  extended with  $X = x_i$

**return** NORMALIZE( $Q(X)$ )

**function** ENUMERATE-ALL( $vars, \mathbf{e}$ ) **returns** a real number

**if** EMPTY?( $vars$ ) **then return** 1.0

$V \leftarrow$  FIRST( $vars$ )


**if**  $V$  is an evidence variable with value  $v$  in  $\mathbf{e}$

**then return**  $P(v | \text{parents}(V)) \times$  ENUMERATE-ALL(REST( $vars$ ),  $\mathbf{e}$ )

**else return**  $\sum_v P(v | \text{parents}(V)) \times$  ENUMERATE-ALL(REST( $vars$ ),  $\mathbf{e}_v$ )

where  $\mathbf{e}_v$  is  $\mathbf{e}$  extended with  $V = v$

- Depth-first recursion
- $O(2^n)$  time for  $n$  variables
- Repeated evaluations of the same subexpressions



# Enumeration Algorithm

**function** ENUMERATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayes net with variables  $vars$

$Q(X) \leftarrow$  a distribution over  $X$ , initially empty

**for each** value  $x_i$  of  $X$  **do**

$Q(x_i) \leftarrow$  ENUMERATE-ALL( $vars, \mathbf{e}_{x_i}$ )

where  $\mathbf{e}_{x_i}$  is  $\mathbf{e}$  extended with  $X = x_i$

**return** NORMALIZE( $Q(X)$ )

**function** ENUMERATE-ALL( $vars, \mathbf{e}$ ) **returns** a real number

**if** EMPTY?( $vars$ ) **then return** 1.0

$V \leftarrow$  FIRST( $vars$ )


**if**  $V$  is an evidence variable with value  $v$  in  $\mathbf{e}$

**then return**  $P(v | \text{parents}(V)) \times$  ENUMERATE-ALL(REST( $vars$ ),  $\mathbf{e}$ )

**else return**  $\sum_v P(v | \text{parents}(V)) \times$  ENUMERATE-ALL(REST( $vars$ ),  $\mathbf{e}_v$ )

where  $\mathbf{e}_v$  is  $\mathbf{e}$  extended with  $V = v$

- Depth-first recursion
- $O(2^n)$  time for  $n$  variables
- Repeated evaluations of the same subexpressions



# Enumeration Algorithm

**function** ENUMERATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayes net with variables  $vars$

$Q(X) \leftarrow$  a distribution over  $X$ , initially empty

**for each** value  $x_i$  of  $X$  **do**

$Q(x_i) \leftarrow$  ENUMERATE-ALL( $vars, \mathbf{e}_{x_i}$ )

where  $\mathbf{e}_{x_i}$  is  $\mathbf{e}$  extended with  $X = x_i$

**return** NORMALIZE( $Q(X)$ )

**function** ENUMERATE-ALL( $vars, \mathbf{e}$ ) **returns** a real number

**if** EMPTY?( $vars$ ) **then return** 1.0

$V \leftarrow$  FIRST( $vars$ )


**if**  $V$  is an evidence variable with value  $v$  in  $\mathbf{e}$

**then return**  $P(v | \text{parents}(V)) \times$  ENUMERATE-ALL(REST( $vars$ ),  $\mathbf{e}$ )

**else return**  $\sum_v P(v | \text{parents}(V)) \times$  ENUMERATE-ALL(REST( $vars$ ),  $\mathbf{e}_v$ )

where  $\mathbf{e}_v$  is  $\mathbf{e}$  extended with  $V = v$

- Depth-first recursion
- $O(2^n)$  time for  $n$  variables
- Repeated evaluations of the same subexpressions





## II. Variable Elimination

---

**Idea** Do the calculation once and save the results for later use (like dynamic programming).

## II. Variable Elimination

---

**Idea** Do the calculation once and save the results for later use (like dynamic programming).

- ◆ Evaluate an expression from right to left (i.e., bottom up in the expression tree).

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$

## II. Variable Elimination

---

**Idea** Do the calculation once and save the results for later use (like dynamic programming).

- ◆ Evaluate an expression from right to left (i.e., bottom up in the expression tree).

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$



## II. Variable Elimination

---

**Idea** Do the calculation once and save the results for later use (like dynamic programming).

- ◆ Evaluate an expression from right to left (i.e., bottom up in the expression tree).

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$



$$P(B \mid j, m) = \alpha P(B) \sum_E P(E) \sum_A P(A \mid B, E) P(j \mid A) P(m \mid A)$$

## II. Variable Elimination

**Idea** Do the calculation once and save the results for later use (like dynamic programming).

- ◆ Evaluate an expression from right to left (i.e., bottom up in the expression tree).

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$



$$P(B \mid j, m) = \alpha \underbrace{P(B)} \sum_E \underbrace{P(E)} \sum_A \underbrace{P(A \mid B, E)} \underbrace{P(j \mid A)} \underbrace{P(m \mid A)}$$

**factors:**  $f_1(B)$   $f_2(E)$   $f_3(A, B, E)$   $f_4(A)$   $f_5(A)$

## II. Variable Elimination

**Idea** Do the calculation once and save the results for later use (like dynamic programming).

- ◆ Evaluate an expression from right to left (i.e., bottom up in the expression tree).

$$P(b \mid j, m) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(j \mid A) P(m \mid A)$$



$$P(B \mid j, m) = \alpha \underbrace{P(B)} \sum_E \underbrace{P(E)} \sum_A \underbrace{P(A \mid B, E)} \underbrace{P(j \mid A)} \underbrace{P(m \mid A)}$$

**factors:**  $f_1(B)$   $f_2(E)$   $f_3(A, B, E)$   $f_4(A)$   $f_5(A)$

**dimensions:**  $2 \times 1$   $2 \times 1$   $2 \times 2 \times 2$   $2 \times 1$   $2 \times 1$

# Evaluation Example

---

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A)$$

# Evaluation Example

---

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A)$$

- Sum out  $A$  from the product of  $f_3, f_4, f_5$ .

$$f_6(B, E) = \sum_{A \in \{a, \neg a\}} f_3(A, B, E) \times f_4(A) \times f_5(A)$$

$2 \times 2$



# Evaluation Example

---

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A)$$

- Sum out  $A$  from the product of  $f_3, f_4, f_5$ .

$$\begin{aligned} f_6(B, E) &= \sum_{A \in \{a, \neg a\}} f_3(A, B, E) \times f_4(A) \times f_5(A) \\ 2 \times 2 & \\ &= \left( f_3(a, B, E) \times P(a) \times P(a) \right) + \left( f_3(\neg a, B, E) \times P(\neg a) \times P(\neg a) \right) \\ &\quad 2 \times 2 \qquad \qquad \qquad 2 \times 2 \end{aligned}$$

# Evaluation Example

$$\mathbf{P}(B \mid j, m) = \alpha \mathbf{f}_1(B) \times \sum_E \mathbf{f}_2(E) \times \sum_A \mathbf{f}_3(A, B, E) \times \mathbf{f}_4(A) \times \mathbf{f}_5(A)$$

- Sum out  $A$  from the product of  $\mathbf{f}_3, \mathbf{f}_4, \mathbf{f}_5$ .

$$\begin{aligned} \mathbf{f}_6(B, E) &= \sum_{A \in \{a, \neg a\}} \mathbf{f}_3(A, B, E) \times \mathbf{f}_4(A) \times \mathbf{f}_5(A) \\ 2 \times 2 &= \left( \mathbf{f}_3(a, B, E) \times \textcolor{red}{P}(a) \times \textcolor{red}{P}(a) \right) + \left( \mathbf{f}_3(\neg a, B, E) \times P(\neg a) \times P(\neg a) \right) \\ &\quad 2 \times 2 \qquad \qquad \qquad 2 \times 2 \end{aligned}$$

$$\mathbf{P}(B \mid j, m) = \alpha \mathbf{f}_1(B) \times \sum_E \mathbf{f}_2(E) \times \mathbf{f}_6(B, E)$$

# Evaluation Example

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A)$$

- Sum out  $A$  from the product of  $f_3, f_4, f_5$ .

$$\begin{aligned} f_6(B, E) &= \sum_{A \in \{a, \neg a\}} f_3(A, B, E) \times f_4(A) \times f_5(A) \\ 2 \times 2 & \\ &= \left( f_3(a, B, E) \times P(a) \times P(a) \right) + \left( f_3(\neg a, B, E) \times P(\neg a) \times P(\neg a) \right) \\ &\quad 2 \times 2 \qquad \qquad \qquad 2 \times 2 \end{aligned}$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times f_6(B, E)$$

- Sum out  $E$  from the product of  $f_2$  and  $f_6$ .

$$f_7(B) = \sum_{E \in \{e, \neg e\}} f_2(E) \times f_6(B, E) = P(e) \times f_6(B, e) + P(\neg e) \times f_6(B, \neg e)$$

$2 \times 1 \qquad \qquad \qquad 2 \times 1$

# Evaluation Example

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A)$$

- Sum out  $A$  from the product of  $f_3, f_4, f_5$ .

$$\begin{aligned} f_6(B, E) &= \sum_{A \in \{a, \neg a\}} f_3(A, B, E) \times f_4(A) \times f_5(A) \\ 2 \times 2 & \\ &= \left( f_3(a, B, E) \times P(a) \times P(a) \right) + \left( f_3(\neg a, B, E) \times P(\neg a) \times P(\neg a) \right) \\ &\quad 2 \times 2 \qquad \qquad \qquad 2 \times 2 \end{aligned}$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times f_6(B, E)$$

- Sum out  $E$  from the product of  $f_2$  and  $f_6$ .

$$f_7(B) = \sum_{E \in \{e, \neg e\}} f_2(E) \times f_6(B, E) = P(e) \times f_6(B, e) + P(\neg e) \times f_6(B, \neg e)$$

$2 \times 1 \qquad \qquad \qquad 2 \times 1$

- Finally, carry out the following pointwise product:

$$P(B \mid j, m) = \alpha f_1(B) \times f_7(B)$$

# Pointwise Product of Two Factors


---

$$f(X_1, \dots, X_j, Y_1, \dots, Y_k) \times g(Y_1, \dots, Y_k, Z_1, \dots, Z_l) = h(X_1, \dots, X_j, Y_1, \dots, Y_k, Z_1, \dots, Z_l)$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Pointwise Product of Two Factors


$$f(X_1, \dots, X_j, Y_1, \dots, Y_k) \times g(Y_1, \dots, Y_k, Z_1, \dots, Z_l) = h(X_1, \dots, X_j, Y_1, \dots, Y_k, Z_1, \dots, Z_l)$$


  
common variables

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Pointwise Product of Two Factors


$$f(X_1, \dots, X_j, Y_1, \dots, Y_k) \times g(Y_1, \dots, Y_k, Z_1, \dots, Z_l) = h(X_1, \dots, X_j, Y_1, \dots, Y_k, Z_1, \dots, Z_l)$$


  
common variables

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Pointwise Product of Two Factors

$$f(X_1, \dots, X_j, Y_1, \dots, Y_k) \times g(Y_1, \dots, Y_k, Z_1, \dots, Z_l) = h(X_1, \dots, X_j, Y_1, \dots, Y_k, Z_1, \dots, Z_l)$$



  
common variables

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$



# Pointwise Product of Two Factors

$$f(X_1, \dots, X_j, Y_1, \dots, Y_k) \times g(Y_1, \dots, Y_k, Z_1, \dots, Z_l) = h(X_1, \dots, X_j, Y_1, \dots, Y_k, Z_1, \dots, Z_l)$$


  
common variables

$X$	$Y$	$f(X, Y)$	$Y$	$Z$	$g(Y, Z)$	$X$	$Y$	$Z$	$h(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X \mathbf{h}(X, Y, Z) = \mathbf{h}(x, Y, Z) + \mathbf{h}(\neg x, Y, Z)$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X h(X, Y, Z) = h(x, Y, Z) + h(\neg x, Y, Z)$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X h(X, Y, Z) = h(x, Y, Z) + h(\neg x, Y, Z)$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X h(X, Y, Z) = h(x, Y, Z) + h(\neg x, Y, Z)$$

$$= \begin{pmatrix} .06 & .24 \\ .42 & .28 \end{pmatrix}$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X h(X, Y, Z) = h(x, Y, Z) + h(\neg x, Y, Z)$$

$$= \begin{pmatrix} .06 & .24 \\ .42 & .28 \end{pmatrix}$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X h(X, Y, Z) = h(x, Y, Z) + h(\neg x, Y, Z)$$

$$= \begin{pmatrix} .06 & .24 \\ .42 & .28 \end{pmatrix}$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X h(X, Y, Z) = h(x, Y, Z) + h(\neg x, Y, Z)$$

$$= \begin{pmatrix} .06 & .24 \\ .42 & .28 \end{pmatrix} + \begin{pmatrix} .18 & .72 \\ .06 & .04 \end{pmatrix}$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$



# Summing Out a Variable

This operation over a product of factors is carried out by adding up the submatrices, each for one value of the same variable.

$$l(Y, Z) = \sum_X h(X, Y, Z) = h(x, Y, Z) + h(\neg x, Y, Z)$$

$$= \begin{pmatrix} .06 & .24 \\ .42 & .28 \end{pmatrix} + \begin{pmatrix} .18 & .72 \\ .06 & .04 \end{pmatrix} = \begin{pmatrix} .24 & .96 \\ .48 & .32 \end{pmatrix}$$

$X$	$Y$	$\mathbf{f}(X, Y)$	$Y$	$Z$	$\mathbf{g}(Y, Z)$	$X$	$Y$	$Z$	$\mathbf{h}(X, Y, Z)$
$t$	$t$	.3	$t$	$t$	.2	$t$	$t$	$t$	$.3 \times .2 = .06$
$t$	$f$	.7	$t$	$f$	.8	$t$	$t$	$f$	$.3 \times .8 = .24$
$f$	$t$	.9	$f$	$t$	.6	$t$	$f$	$t$	$.7 \times .6 = .42$
$f$	$f$	.1	$f$	$f$	.4	$t$	$f$	$f$	$.7 \times .4 = .28$
						$f$	$t$	$t$	$.9 \times .2 = .18$
						$f$	$t$	$f$	$.9 \times .8 = .72$
						$f$	$f$	$t$	$.1 \times .6 = .06$
						$f$	$f$	$f$	$.1 \times .4 = .04$

# Variable Elimination Algorithm

---

Move outside the summation any factor independent of the variable to be summed out.

# Variable Elimination Algorithm

---

Move outside the summation any factor independent of the variable to be summed out.

$$\sum_X f(X, Y) \times g(Y, Z) = g(Y, Z) \times \sum_X f(X, Y)$$

# Variable Elimination Algorithm

---

Move outside the summation any factor independent of the variable to be summed out.

$$\sum_X f(X, Y) \times g(Y, Z) = g(Y, Z) \times \sum_X f(X, Y)$$

**function** ELIMINATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayesian network with variables  $vars$

$factors \leftarrow []$

**for each**  $V$  **in** ORDER( $vars$ ) **do**

$factors \leftarrow [\text{MAKE-FACTOR}(V, \mathbf{e})] + factors$

**if**  $V$  is a hidden variable **then**  $factors \leftarrow \text{SUM-OUT}(V, factors)$

**return** NORMALIZE(POINTWISE-PRODUCT( $factors$ ))

# Variable Elimination Algorithm

---

Move outside the summation any factor independent of the variable to be summed out.

$$\sum_X f(X, Y) \times g(Y, Z) = g(Y, Z) \times \sum_X f(X, Y)$$

**function** ELIMINATION-ASK( $X, \mathbf{e}, bn$ ) **returns** a distribution over  $X$

**inputs:**  $X$ , the query variable

$\mathbf{e}$ , observed values for variables  $\mathbf{E}$

$bn$ , a Bayesian network with variables  $vars$

$factors \leftarrow []$  ?

**for each**  $V$  **in** ORDER( $vars$ ) **do**

$factors \leftarrow [\text{MAKE-FACTOR}(V, \mathbf{e})] + factors$

**if**  $V$  is a hidden variable **then**  $factors \leftarrow \text{SUM-OUT}(V, factors)$

**return** NORMALIZE(Pointwise-Product( $factors$ ))

# III. Ordering Variables to Eliminate

---

$$P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$$

# III. Ordering Variables to Eliminate

---

♣ Every choice of ordering yields a valid algorithm.  $\Leftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

# III. Ordering Variables to Eliminate

---

♣ Every choice of ordering yields a valid algorithm.  $\Longleftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A) \quad (\text{Order: } A, E)$$



# III. Ordering Variables to Eliminate

♣ Every choice of ordering yields a valid algorithm.  $\Longleftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A) \quad (\text{Order: } A, E)$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_A f_4(A) \times f_5(A) \times \sum_E f_2(E) \times f_3(A, B, E) \quad (\text{Order: } E, A)$$

# III. Ordering Variables to Eliminate

♣ Every choice of ordering yields a valid algorithm.  $\Longleftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A) \quad (\text{Order: } A, E)$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_A f_4(A) \times f_5(A) \times \sum_E f_2(E) \times f_3(A, B, E) \quad (\text{Order: } E, A)$$

♣ Different orderings generates different intermediate factors.

# III. Ordering Variables to Eliminate

♣ Every choice of ordering yields a valid algorithm.  $\Longleftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A) \quad (\text{Order: } A, E)$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_A f_4(A) \times f_5(A) \times \sum_E f_2(E) \times f_3(A, B, E) \quad (\text{Order: } E, A)$$

♣ Different orderings generates different intermediate factors.

♣ Time and space are dominated by the size of the largest factor constructed, which is subject to two factors:

# III. Ordering Variables to Eliminate

♣ Every choice of ordering yields a valid algorithm.  $\Longleftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A) \quad (\text{Order: } A, E)$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_A f_4(A) \times f_5(A) \times \sum_E f_2(E) \times f_3(A, B, E) \quad (\text{Order: } E, A)$$

♣ Different orderings generates different intermediate factors.

♣ Time and space are dominated by the size of the largest factor constructed, which is subject to two factors:

- ordering of variables
- structure of the network

# III. Ordering Variables to Eliminate

♣ Every choice of ordering yields a valid algorithm.  $\Longleftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A) \quad (\text{Order: } A, E)$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_A f_4(A) \times f_5(A) \times \sum_E f_2(E) \times f_3(A, B, E) \quad (\text{Order: } E, A)$$

- ♣ Different orderings generates different intermediate factors.
- ♣ Time and space are dominated by the size of the largest factor constructed, which is subject to two factors:
  - ordering of variables
  - structure of the network
- ♣ It is intractable to determine the optimal order.

# III. Ordering Variables to Eliminate

♣ Every choice of ordering yields a valid algorithm.  $\Longleftarrow P(x_1, \dots, x_n) = \prod_{i=1}^n P(x_i \mid \text{Parents}(X_i))$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_E f_2(E) \times \sum_A f_3(A, B, E) \times f_4(A) \times f_5(A) \quad (\text{Order: } A, E)$$

$$P(B \mid j, m) = \alpha f_1(B) \times \sum_A f_4(A) \times f_5(A) \times \sum_E f_2(E) \times f_3(A, B, E) \quad (\text{Order: } E, A)$$

- ♣ Different orderings generates different intermediate factors.
- ♣ Time and space are dominated by the size of the largest factor constructed, which is subject to two factors:
  - ordering of variables
  - structure of the network
- ♣ It is intractable to determine the optimal order.
- ♣ Use a greedy heuristic: eliminate whichever variable minimizes the size of the next factor to be constructed.


# Irrelevant Variable

---

$P(\text{JohnCalls} \mid \text{Burglary} = \text{true})?$

# Irrelevant Variable

$P(\text{JohnCalls} \mid \text{Burglary} = \text{true})?$






# Irrelevant Variable

$P(\text{JohnCalls} \mid \text{Burglary} = \text{true})?$

$$P(J \mid b) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(J \mid A) \sum_M P(M \mid A)$$




# Irrelevant Variable

$P(\text{JohnCalls} \mid \text{Burglary} = \text{true})?$

$$P(J \mid b) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(J \mid A) \sum_M P(M \mid A)$$

$$\sum_M P(M \mid A) = 1$$



# Irrelevant Variable


$P(\text{JohnCalls} \mid \text{Burglary} = \text{true})?$

$$P(J \mid b) = \alpha P(b) \sum_E P(E) \sum_A P(A \mid b, E) P(J \mid A) \sum_M P(M \mid A)$$

$$\sum_M P(M \mid A) = 1$$



The variable  $M$  is irrelevant to the query.



# Inference with Elimination

---

- We can remove a leaf node (e.g.,  $M$ ) that is neither a query variable nor an evidence variable.
- After the removal, there may be more leaf nodes that are irrelevant. Remove them as well, and so on.

# Inference with Elimination

---

- We can remove a leaf node (e.g.,  $M$ ) that is neither a query variable nor an evidence variable.
- After the removal, there may be more leaf nodes that are irrelevant. Remove them as well, and so on.

Every variable that is not an ancestor of a query variable or evidence variable is irrelevant to the query.

# Inference with Elimination

---

- We can remove a leaf node (e.g.,  $M$ ) that is neither a query variable nor an evidence variable.
- After the removal, there may be more leaf nodes that are irrelevant. Remove them as well, and so on.

Every variable that is not an ancestor of a query variable or evidence variable is irrelevant to the query.

- ♦ Using *reverse topological order* for variables, exact inference with elimination can be 1,000 times faster than the enumeration algorithm.

# Inference with Elimination

---

- We can remove a leaf node (e.g.,  $M$ ) that is neither a query variable nor an evidence variable.
- After the removal, there may be more leaf nodes that are irrelevant. Remove them as well, and so on.

Every variable that is not an ancestor of a query variable or evidence variable is irrelevant to the query.

- ♦ Using *reverse topological order* for variables, exact inference with elimination can be 1,000 times faster than the enumeration algorithm.
- ♦ If we want to compute posterior probabilities for all the variables rather than answer individual queries, we can use clustering algorithms (i.e., *join tree algorithms*).