

Probabilistic Inference

Outline

I. Probability for continuous variables

II. Inference by enumeration

I. Probability Density Function

A *continuous random variable* has its outcome take on a continuous set of values.

- the high temperature tomorrow
- the reading from a noisy force sensor
- the actual arrival time of a flight, etc.

I. Probability Density Function

A *continuous random variable* has its outcome take on a continuous set of values.

- the high temperature tomorrow
- the reading from a noisy force sensor
- the actual arrival time of a flight, etc.

The *cumulative distribution function* of a continuous random variable X is

$$D(x) = P(X \leq x)$$

I. Probability Density Function

A *continuous random variable* has its outcome take on a continuous set of values.

- the high temperature tomorrow
- the reading from a noisy force sensor
- the actual arrival time of a flight, etc.

The *cumulative distribution function* of a continuous random variable X is

$$D(x) = P(X \leq x)$$

Its derivative is the *probability density function (p.d.f.)*:

I. Probability Density Function

A *continuous random variable* has its outcome take on a continuous set of values.

- the high temperature tomorrow
- the reading from a noisy force sensor
- the actual arrival time of a flight, etc.

The *cumulative distribution function* of a continuous random variable X is

$$D(x) = P(X \leq x)$$

Its derivative is the *probability density function (p.d.f.)*:

$$p(x) = \frac{d}{dx} D(x)$$

I. Probability Density Function

A *continuous random variable* has its outcome take on a continuous set of values.

- the high temperature tomorrow
- the reading from a noisy force sensor
- the actual arrival time of a flight, etc.

The *cumulative distribution function* of a continuous random variable X is

$$D(x) = P(X \leq x)$$

Its derivative is the *probability density function (p.d.f.)*:

$$p(x) = \frac{d}{dx} D(x)$$

$$= \lim_{dx \rightarrow 0} \frac{P(X \leq x + dx) - P(X \leq x)}{dx}$$

$$\lim_{dx \rightarrow 0} \frac{P(X \leq x + dx) - P(X \leq x)}{dx}$$

I. Probability Density Function

A *continuous random variable* has its outcome take on a continuous set of values.

- the high temperature tomorrow
- the reading from a noisy force sensor
- the actual arrival time of a flight, etc.

The *cumulative distribution function* of a continuous random variable X is

$$D(x) = P(X \leq x)$$

Its derivative is the *probability density function (p.d.f.)*:

$$p(x) = \frac{d}{dx} D(x)$$

$$= \lim_{dx \rightarrow 0} \frac{P(X \leq x + dx) - P(X \leq x)}{dx}$$


$$= \lim_{dx \rightarrow 0} \frac{P(x \leq X \leq x + dx)}{dx}$$

Uniform Distribution

A *uniform distribution* has constant p.d.f.

Example Continuous uniform distribution over the interval $[a, b]$.

$$p(x) = \begin{cases} 0, & \text{for } x < a, \\ \frac{1}{b-a}, & \text{for } a \leq x < b, \\ 0, & \text{for } x > b, \end{cases} \quad D(x) = \begin{cases} 0, & \text{for } x < a, \\ \frac{x-a}{b-a}, & \text{for } a \leq x < b, \\ 1, & \text{for } x \geq b, \end{cases}$$



Expected Value

The *expected value* (or *mean*) $E(X)$ of a random variable X is its average value over a large number of chance experiments.

Expected Value

The *expected value* (or *mean*) $E(X)$ of a random variable X is its average value over a large number of chance experiments.

- n experiments

Expected Value

The *expected value* (or *mean*) $E(X)$ of a random variable X is its average value over a large number of chance experiments.

- n experiments
- m different values: x_1, \dots, x_m

Expected Value

The *expected value* (or *mean*) $E(X)$ of a random variable X is its average value over a large number of chance experiments.

- n experiments
- m different values: x_1, \dots, x_m
- For $1 \leq i \leq m$, the value x_i occurs n_i times. $// n = n_1 + \dots + n_m$

Expected Value

The *expected value* (or *mean*) $E(X)$ of a random variable X is its average value over a large number of chance experiments.

- n experiments
- m different values: x_1, \dots, x_m
- For $1 \leq i \leq m$, the value x_i occurs n_i times. // $n = n_1 + \dots + n_m$

$$E(X) = \frac{1}{n} \sum_{i=1}^m x_i \cdot n_i$$

Expected Value

The *expected value* (or *mean*) $E(X)$ of a random variable X is its average value over a large number of chance experiments.

- n experiments
- m different values: x_1, \dots, x_m
- For $1 \leq i \leq m$, the value x_i occurs n_i times. // $n = n_1 + \dots + n_m$

$$E(X) = \frac{1}{n} \sum_{i=1}^m x_i \cdot n_i$$

Example Roll a die an infinite number of times. Each number appears 1/6 of the time.

$$E(X) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^6 i \cdot \frac{n}{6} = \frac{7}{2}$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

The *variance* of X measures the dispersion of the values x_1, \dots, x_n around μ .

$$\text{var}(X) = E((X - E(X))^2)$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

The *variance* of X measures the dispersion of the values x_1, \dots, x_n around μ .

$$\text{var}(X) = E((X - E(X))^2) = \sum_{i=1}^m p_i(x_i - \mu)^2$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

The *variance* of X measures the dispersion of the values x_1, \dots, x_n around μ .

$$\begin{aligned} \text{var}(X) &= E((X - E(X))^2) = \sum_{i=1}^m p_i(x_i - \mu)^2 \\ &= E((X - \mu)(X - \mu)) \end{aligned}$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

The *variance* of X measures the dispersion of the values x_1, \dots, x_n around μ .

$$\begin{aligned}\text{var}(X) &= E((X - E(X))^2) = \sum_{i=1}^m p_i(x_i - \mu)^2 \\ &= E((X - \mu)(X - \mu)) \\ &= E(X^2 - 2X\mu + \mu^2)\end{aligned}$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

The *variance* of X measures the dispersion of the values x_1, \dots, x_n around μ .

$$\text{var}(X) = E((X - E(X))^2) = \sum_{i=1}^m p_i(x_i - \mu)^2$$

$$= E((X - \mu)(X - \mu))$$

$$= E(X^2 - 2X\mu + \mu^2)$$

$$= E(X^2) - 2\mu^2 + \mu^2$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

The *variance* of X measures the dispersion of the values x_1, \dots, x_n around μ .

$$\begin{aligned}\text{var}(X) &= E((X - E(X))^2) = \sum_{i=1}^m p_i(x_i - \mu)^2 \\ &= E((X - \mu)(X - \mu)) \\ &= E(X^2 - 2X\mu + \mu^2) \\ &= E(X^2) - 2\mu^2 + \mu^2 \\ &= E(X^2) - \mu^2\end{aligned}$$

Variance and Standard Deviation

Suppose p_i is the probability of the value x_i , for $1 \leq i \leq m$.

The mean μ of X is

$$\mu = E(X) = p_1x_1 + \cdots + p_mx_m.$$

The *variance* of X measures the dispersion of the values x_1, \dots, x_n around μ .

$$\begin{aligned}\text{var}(X) &= E((X - E(X))^2) = \sum_{i=1}^m p_i(x_i - \mu)^2 \\ &= E((X - \mu)(X - \mu)) \\ &= E(X^2 - 2X\mu + \mu^2) \\ &= E(X^2) - 2\mu^2 + \mu^2 \\ &= E(X^2) - \mu^2\end{aligned}$$

The *standard deviation* of X is $\sigma = \sqrt{\text{var}(X)}$.

Mean and Variance of a Continuous Variable

If X is a continuous variable with p.d.f. $p(X)$, its expected value is

$$\mu = E(X) = \int_{-\infty}^{\infty} x p(x) dx$$

Mean and Variance of a Continuous Variable

If X is a continuous variable with p.d.f. $p(X)$, its expected value is

$$\mu = E(X) = \int_{-\infty}^{\infty} x p(x) dx$$

Its variance is given as

$$\text{var}(X) = \int_{-\infty}^{\infty} p(x)(x - \mu)^2 dx$$

Gaussian Distribution


A continuous random variable X with mean μ and variance σ^2 has *Gaussian distribution* (or *normal distribution*) if its p.d.f. is

$$p(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)}$$

Gaussian Distribution

A continuous random variable X with mean μ and variance σ^2 has *Gaussian distribution* (or *normal distribution*) if its p.d.f. is


$$p(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)}$$



Gaussian Distribution

A continuous random variable X with mean μ and variance σ^2 has *Gaussian distribution* (or *normal distribution*) if its p.d.f. is

$$p(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)}$$



$$P(\mu - \sigma \leq X \leq \mu + \sigma) \approx 0.6826895$$

$$P(\mu - 2\sigma \leq X \leq \mu + 2\sigma) \approx 0.9544997$$

$$P(\mu - 3\sigma \leq X \leq \mu + 3\sigma) \approx 0.9973002$$


$$P(\mu - 4\sigma \leq X \leq \mu + 4\sigma) \approx 0.9999366$$

$$P(\mu - 5\sigma \leq X \leq \mu + 5\sigma) \approx 0.9999994$$

Gaussian Distribution

A continuous random variable X with mean μ and variance σ^2 has *Gaussian distribution* (or *normal distribution*) if its p.d.f. is

$$p(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)}$$



$$\begin{aligned} P(\mu - \sigma \leq X \leq \mu + \sigma) &\approx 0.6826895 \\ P(\mu - 2\sigma \leq X \leq \mu + 2\sigma) &\approx 0.9544997 \\ P(\mu - 3\sigma \leq X \leq \mu + 3\sigma) &\approx 0.9973002 \\ P(\mu - 4\sigma \leq X \leq \mu + 4\sigma) &\approx 0.9999366 \\ P(\mu - 5\sigma \leq X \leq \mu + 5\sigma) &\approx 0.9999994 \end{aligned}$$

Three-sigma rule (in practice): Consider population within $(\mu - 3\sigma, \mu + 3\sigma)$.

Why Gaussian Distribution?

- ◆ It is the **most important distribution** because it fits many natural phenomena (e.g., human characteristics such as weight, height, body temperature, etc.)
- ◆ It is the limiting distribution of $X_1 + \dots + X_n$ of n independent random variables X_1, \dots, X_n , as $n \rightarrow \infty$, explaining a characteristic impacted by numerous independent factors (by the central limit theorem).
- ◆ It is the foundation for important methods such as least-squares, Kalman filters that are used in statistics and engineering.

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

$$P(\phi) = \sum_{\omega \in \phi} P(\omega)$$

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

$$P(\phi) = \sum_{\omega \in \phi} P(\omega)$$

Joint distribution table:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

$$P(\phi) = \sum_{\omega \in \phi} P(\omega)$$

Joint distribution table:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

$P(\text{cavity} \vee \text{toothache})$

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

$$P(\phi) = \sum_{\omega \in \phi} P(\omega)$$

Joint distribution table:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

$P(\text{cavity} \vee \text{toothache})$

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

$$P(\phi) = \sum_{\omega \in \phi} P(\omega)$$

Joint distribution table:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

$$P(\text{cavity} \vee \text{toothache}) = 0.108 + 0.012 + 0.072 + 0.008$$

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

$$P(\phi) = \sum_{\omega \in \phi} P(\omega)$$

Joint distribution table:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

$$P(\text{cavity} \vee \text{toothache}) = 0.108 + 0.012 + 0.072 + 0.008$$

II. Inference by Enumeration

Probabilistic inference: compute posterior probabilities for query propositions given observed evidence.

Calculate the probability of a proposition ϕ :

$$P(\phi) = \sum_{\omega \in \phi} P(\omega)$$

Joint distribution table:

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

$$P(\text{cavity} \vee \text{toothache}) = 0.108 + 0.012 + 0.072 + 0.008 + 0.016 + 0.064 = 0.28$$

Marginalization

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

The unconditional (or marginal) probability of \neg *toothache*:

$$P(\neg\textit{toothache})$$

Marginalization

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

The unconditional (or marginal) probability of \neg *toothache*:

$$P(\neg\textit{toothache})$$

Marginalization

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

The unconditional (or marginal) probability of \neg *toothache*:

$$P(\neg\textit{toothache}) = 0.072 + 0.144 + 0.008 + 0.576 = 0.8$$

Marginalization

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

The unconditional (or marginal) probability of \neg *toothache*:

$$P(\neg\textit{toothache}) = 0.072 + 0.144 + 0.008 + 0.576 = 0.8$$

Marginalization: Sum up the probabilities for each possible value of the other variables.

$$P(Y) = \sum_z P(Y, Z = z)$$

Marginalization

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

The unconditional (or marginal) probability of \neg *toothache*:

$$P(\neg\textit{toothache}) = 0.072 + 0.144 + 0.008 + 0.576 = 0.8$$

Marginalization: Sum up the probabilities for each possible value of the other variables.

$$P(Y) = \sum_{\mathbf{Z}} P(Y, \mathbf{Z} = \mathbf{z})$$

|
Sums over all possible combinations
of the values of the set of variables \mathbf{Z} .

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

$$\begin{aligned} P(\text{Cavity}) = & P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch}) \\ & + P(\text{Cavity}, \neg \text{toothache}, \text{catch}) + P(\text{Cavity}, \neg \text{toothache}, \neg \text{catch}) \end{aligned}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

$$P(\text{Cavity}) = P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch}) \\ + P(\text{Cavity}, \neg \text{toothache}, \text{catch}) + P(\text{Cavity}, \neg \text{toothache}, \neg \text{catch})$$

random variable
beginning with
an uppercase
letter.

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

$$\begin{aligned} P(\text{Cavity}) &= P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch}) \\ &\quad + P(\text{Cavity}, \neg \text{toothache}, \text{catch}) + P(\text{Cavity}, \neg \text{toothache}, \neg \text{catch}) \\ &= \langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle + \langle 0.072, 0.144 \rangle + \langle 0.008, 0.576 \rangle \end{aligned}$$

random variable
beginning with
an uppercase
letter.

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

$$\begin{aligned} P(\text{Cavity}) &= P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch}) \\ &\quad + P(\text{Cavity}, \neg \text{toothache}, \text{catch}) + P(\text{Cavity}, \neg \text{toothache}, \neg \text{catch}) \\ &= \langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle + \langle 0.072, 0.144 \rangle + \langle 0.008, 0.576 \rangle \\ &= \langle 0.2, 0.8 \rangle \end{aligned}$$

random variable
beginning with
an uppercase
letter.

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

$$\begin{aligned} P(\text{Cavity}) &= P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch}) \\ &\quad + P(\text{Cavity}, \neg \text{toothache}, \text{catch}) + P(\text{Cavity}, \neg \text{toothache}, \neg \text{catch}) \\ &= \langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle + \langle 0.072, 0.144 \rangle + \langle 0.008, 0.576 \rangle \\ &= \langle 0.2, 0.8 \rangle \end{aligned}$$

random variable
beginning with
an uppercase
letter.

Rule of *conditioning*:

$$P(Y) = \sum_z P(Y | z) P(z)$$

	<i>toothache</i>		$\neg \text{toothache}$	
	<i>catch</i>	$\neg \text{catch}$	<i>catch</i>	$\neg \text{catch}$
<i>cavity</i>	0.108	0.012	0.072	0.008
$\neg \text{cavity}$	0.016	0.064	0.144	0.576

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

$$\begin{aligned}
 P(\text{Cavity}) &= P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch}) \\
 &\quad + P(\text{Cavity}, \neg \text{toothache}, \text{catch}) + P(\text{Cavity}, \neg \text{toothache}, \neg \text{catch}) \\
 &= \langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle + \langle 0.072, 0.144 \rangle + \langle 0.008, 0.576 \rangle \\
 &= \langle 0.2, 0.8 \rangle
 \end{aligned}$$

random variable
beginning with
an uppercase
letter.

not bold-faced

Rule of *conditioning*:

$$P(Y) = \sum_z P(Y | z) P(z)$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Conditioning

Abbreviate $P(Y, Z = z)$ as $P(Y, z)$.

$$\begin{aligned}
 P(\text{Cavity}) &= P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch}) \\
 &\quad + P(\text{Cavity}, \neg \text{toothache}, \text{catch}) + P(\text{Cavity}, \neg \text{toothache}, \neg \text{catch}) \\
 &= \langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle + \langle 0.072, 0.144 \rangle + \langle 0.008, 0.576 \rangle \\
 &= \langle 0.2, 0.8 \rangle
 \end{aligned}$$

random variable
beginning with
an uppercase
letter.

Rule of *conditioning*:

$$\begin{aligned}
 P(Y) &= \sum_z \underbrace{P(Y | z)P(z)}_{\text{not bold-faced}} \\
 &= P(Y, z)
 \end{aligned}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Computing Conditional Probabilities

1. $P(a \mid b) = \frac{P(a \wedge b)}{P(b)}$ \Rightarrow an expression in terms of unconditional probabilities.
2. Evaluate the expression from full joint distribution.

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Computing Conditional Probabilities

1. $P(a | b) = \frac{P(a \wedge b)}{P(b)}$ \Rightarrow an expression in terms of unconditional probabilities.
2. Evaluate the expression from full joint distribution.

$$P(\text{cavity} | \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Computing Conditional Probabilities

1. $P(a | b) = \frac{P(a \wedge b)}{P(b)}$ \Rightarrow an expression in terms of unconditional probabilities.
2. Evaluate the expression from full joint distribution.

$$P(\text{cavity} | \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Computing Conditional Probabilities

1. $P(a | b) = \frac{P(a \wedge b)}{P(b)}$ \Rightarrow an expression in terms of unconditional probabilities.
2. Evaluate the expression from full joint distribution.

$$\begin{aligned} P(\text{cavity} | \text{toothache}) &= \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\ &= \frac{0.108 + 0.012}{0.108 + 0.012 + 0.016 + 0.064} = 0.6 \end{aligned}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Computing Conditional Probabilities

1. $P(a | b) = \frac{P(a \wedge b)}{P(b)} \Rightarrow$ an expression in terms of unconditional probabilities.
2. Evaluate the expression from full joint distribution.

$$\begin{aligned} P(\text{cavity} | \text{toothache}) &= \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\ &= \frac{0.108 + 0.012}{0.108 + 0.012 + 0.016 + 0.064} = 0.6 \end{aligned}$$

$$P(\neg \text{cavity} | \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Computing Conditional Probabilities

1. $P(a | b) = \frac{P(a \wedge b)}{P(b)} \Rightarrow$ an expression in terms of unconditional probabilities.
2. Evaluate the expression from full joint distribution.

$$\begin{aligned} P(\text{cavity} | \text{toothache}) &= \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\ &= \frac{0.108 + 0.012}{0.108 + 0.012 + 0.016 + 0.064} = 0.6 \end{aligned}$$

$$P(\neg \text{cavity} | \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Computing Conditional Probabilities

1. $P(a | b) = \frac{P(a \wedge b)}{P(b)} \Rightarrow$ an expression in terms of unconditional probabilities.
2. Evaluate the expression from full joint distribution.

$$\begin{aligned} P(\text{cavity} | \text{toothache}) &= \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\ &= \frac{0.108 + 0.012}{0.108 + 0.012 + 0.016 + 0.064} = 0.6 \end{aligned}$$

$$\begin{aligned} P(\neg \text{cavity} | \text{toothache}) &= \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\ &= \frac{0.012 + 0.064}{0.108 + 0.012 + 0.016 + 0.064} = 0.4 \end{aligned}$$

	<i>toothache</i>		<i>¬toothache</i>	
	<i>catch</i>	<i>¬catch</i>	<i>catch</i>	<i>¬catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
<i>¬cavity</i>	0.016	0.064	0.144	0.576

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$


- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.
- ◆ We can calculate $P(\text{Cavity} \mid \text{toothache})$ without $P(\text{toothache})$ as follows.

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.
- ◆ We can calculate $P(\text{Cavity} \mid \text{toothache})$ without $P(\text{toothache})$ as follows.



$$P(\text{Cavity} \mid \text{toothache}) = \alpha P(\text{Cavity}, \text{toothache})$$

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.
- ◆ We can calculate $P(\text{Cavity} \mid \text{toothache})$ without $P(\text{toothache})$ as follows.



$$\begin{aligned} P(\text{Cavity} \mid \text{toothache}) &= \alpha P(\text{Cavity}, \text{toothache}) \\ &= \alpha [P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch})] \end{aligned}$$

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.
- ◆ We can calculate $P(\text{Cavity} \mid \text{toothache})$ without $P(\text{toothache})$ as follows.



$$\begin{aligned} P(\text{Cavity} \mid \text{toothache}) &= \alpha P(\text{Cavity}, \text{toothache}) \\ &= \alpha [P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch})] \\ &= \alpha [\langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle] \end{aligned}$$

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.
- ◆ We can calculate $P(\text{Cavity} \mid \text{toothache})$ without $P(\text{toothache})$ as follows.



$$\begin{aligned} P(\text{Cavity} \mid \text{toothache}) &= \alpha P(\text{Cavity}, \text{toothache}) \\ &= \alpha [P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch})] \\ &= \alpha [\langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle] \\ &= \alpha \langle 0.12, 0.008 \rangle \end{aligned}$$

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.
- ◆ We can calculate $P(\text{Cavity} \mid \text{toothache})$ without $P(\text{toothache})$ as follows.


$$\begin{aligned} P(\text{Cavity} \mid \text{toothache}) &= \alpha P(\text{Cavity}, \text{toothache}) \\ &= \alpha [P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch})] \\ &= \alpha [\langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle] \\ &= \alpha \langle 0.12, 0.008 \rangle \\ &= \langle 0.6, 0.4 \rangle \end{aligned}$$

Normalization

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.6$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = 0.4$$

- ◆ $P(\text{toothache})$ is a normalization constant for the distribution $P(\text{Cavity} \mid \text{toothache})$ so it adds up to 1.
- ◆ We can calculate $P(\text{Cavity} \mid \text{toothache})$ without $P(\text{toothache})$ as follows.

$$P(\text{Cavity} \mid \text{toothache}) = \alpha P(\text{Cavity}, \text{toothache})$$

$$= \alpha [P(\text{Cavity}, \text{toothache}, \text{catch}) + P(\text{Cavity}, \text{toothache}, \neg \text{catch})]$$

$$= \alpha [\langle 0.108, 0.016 \rangle + \langle 0.012, 0.064 \rangle]$$

$$= \alpha \langle 0.12, 0.008 \rangle$$

$$= \langle 0.6, 0.4 \rangle \quad // \alpha = 5 = 1/P(\text{toothache}).$$

Inference Procedure

- X : single query variable (e.g., *Cavity*).
- E : evidence variables (e.g., *Toothache*).
- e : their observed values.
- Y : unobserved variables (e.g., *Catch*).

Inference Procedure

- X : single query variable (e.g., *Cavity*).
- E : evidence variables (e.g., *Toothache*).
- e : their observed values.
- Y : unobserved variables (e.g., *Catch*).

$$P(X | e) \leftarrow \alpha P(X, e) = \alpha \sum_y P(X, e, y)$$

- ♣ Summation is over all possible combinations y of values of variables in Y .
- ♣ $P(X, e, y)$ is a subset of probabilities from the full joint distribution.
- ♣ The full joint distribution has size exponential in # variables and is rarely computed.

Independence

Add a fourth variable *Weather* with domain { *sun*, *rain*, *cloud*, *snow* }.

$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather})$$


	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576

Independence

Add a fourth variable *Weather* with domain { *sun*, *rain*, *cloud*, *snow* }.

$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather})$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576



A 32-element table.


Independence

Add a fourth variable *Weather* with domain { *sun*, *rain*, *cloud*, *snow* }.

$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather})$

$P(\textit{toothache}, \textit{catch}, \textit{cavity}, \textit{cloud})$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576



A 32-element table.

Independence


Add a fourth variable *Weather* with domain { *sun*, *rain*, *cloud*, *snow* }.

$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather})$$

$$P(\textit{toothache}, \textit{catch}, \textit{cavity}, \textit{cloud})$$

$$= P(\textit{cloud} \mid \textit{toothache}, \textit{catch}, \textit{cavity}) P(\textit{toothache}, \textit{catch}, \textit{cavity})$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576



A 32-element table.

Independence

Add a fourth variable *Weather* with domain { *sun*, *rain*, *cloud*, *snow* }.

$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather})$$


$$P(\textit{toothache}, \textit{catch}, \textit{cavity}, \textit{cloud})$$

$$= P(\textit{cloud} \mid \textit{toothache}, \textit{catch}, \textit{cavity}) P(\textit{toothache}, \textit{catch}, \textit{cavity})$$

$$\Downarrow P(\textit{cloud} \mid \textit{toothache}, \textit{catch}, \textit{cavity}) = P(\textit{cloud})$$

$$= P(\textit{cloud}) P(\textit{toothache}, \textit{catch}, \textit{cavity})$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576



A 32-element table.

Independence

Add a fourth variable *Weather* with domain { *sun*, *rain*, *cloud*, *snow* }.

$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather})$$

$$P(\textit{toothache}, \textit{catch}, \textit{cavity}, \textit{cloud})$$


$$= P(\textit{cloud} \mid \textit{toothache}, \textit{catch}, \textit{cavity}) P(\textit{toothache}, \textit{catch}, \textit{cavity})$$

$$\Downarrow P(\textit{cloud} \mid \textit{toothache}, \textit{catch}, \textit{cavity}) = P(\textit{cloud})$$

$$= P(\textit{cloud}) P(\textit{toothache}, \textit{catch}, \textit{cavity})$$

$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather}) = P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}) P(\textit{Weather})$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576



A 32-element
table.

Independence

Add a fourth variable *Weather* with domain { *sun*, *rain*, *cloud*, *snow* }.

$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather})$$

$$P(\textit{toothache}, \textit{catch}, \textit{cavity}, \textit{cloud})$$


$$= P(\textit{cloud} \mid \textit{toothache}, \textit{catch}, \textit{cavity}) P(\textit{toothache}, \textit{catch}, \textit{cavity})$$

$$\Downarrow P(\textit{cloud} \mid \textit{toothache}, \textit{catch}, \textit{cavity}) = P(\textit{cloud})$$

$$= P(\textit{cloud}) P(\textit{toothache}, \textit{catch}, \textit{cavity})$$

$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather}) = \underbrace{P(\textit{Toothache}, \textit{Catch}, \textit{Cavity})}_{\text{8-element table}} \underbrace{P(\textit{Weather})}_{\text{4-element table}}$$

	<i>toothache</i>		\neg <i>toothache</i>	
	<i>catch</i>	\neg <i>catch</i>	<i>catch</i>	\neg <i>catch</i>
<i>cavity</i>	0.108	0.012	0.072	0.008
\neg <i>cavity</i>	0.016	0.064	0.144	0.576



A 32-element
table.

Independent Variables

- ♣ Two propositions a and b are *independent* if

$$P(a \mid b) = P(a) \quad \text{or} \quad P(b \mid a) = P(b) \quad \text{or} \quad P(a \wedge b) = P(a)P(b)$$

Independent Variables

- ♣ Two propositions a and b are *independent* if

$$P(a \mid b) = P(a) \quad \text{or} \quad P(b \mid a) = P(b) \quad \text{or} \quad P(a \wedge b) = P(a)P(b)$$

- ♣ Two variables X and Y are *independent* if

$$\mathbf{P}(X \mid Y) = \mathbf{P}(X) \quad \text{or} \quad \mathbf{P}(Y \mid X) = \mathbf{P}(Y) \quad \text{or} \quad \mathbf{P}(X \wedge Y) = \mathbf{P}(X)P(Y)$$

Independent Variables

- ♣ Two propositions a and b are *independent* if

$$P(a \mid b) = P(a) \quad \text{or} \quad P(b \mid a) = P(b) \quad \text{or} \quad P(a \wedge b) = P(a)P(b)$$

- ♣ Two variables X and Y are *independent* if

$$P(X \mid Y) = P(X) \quad \text{or} \quad P(Y \mid X) = P(Y) \quad \text{or} \quad P(X \wedge Y) = P(X)P(Y)$$

- ♣ Two continuous variables X and Y are *independent* if, for all $x, y \in \mathbb{R}$,

$$P(X \leq x \wedge Y \leq y) = P(X \leq x) \cdot P(Y \leq y)$$

Independent Variables

- ♣ Two propositions a and b are *independent* if

$$P(a \mid b) = P(a) \quad \text{or} \quad P(b \mid a) = P(b) \quad \text{or} \quad P(a \wedge b) = P(a)P(b)$$

- ♣ Two variables X and Y are *independent* if

$$P(X \mid Y) = P(X) \quad \text{or} \quad P(Y \mid X) = P(Y) \quad \text{or} \quad P(X \wedge Y) = P(X)P(Y)$$

- ♣ Two continuous variables X and Y are *independent* if, for all $x, y \in \mathbb{R}$,


$$P(X \leq x \wedge Y \leq y) = P(X \leq x) \cdot P(Y \leq y)$$

The joint probability density function $p(x, y)$ satisfies


$$p(x, y) = \int_{-\infty}^{\infty} p(x, y) \, dy \int_{-\infty}^{\infty} p(x, y) \, dx$$

Factoring a Joint Distribution

The full joint distribution can be factored into *separate* joint distributions on subsets of variables that are *independent*.



Weather and dental problems are independent.



Coin flips are independent.