

Proof Using Resolution

Outline

I. Rule of resolution

II. Resolution refutation

I. Resolution

An inference algorithm i is

sound if $KB \models \alpha$ whenever $KB \vdash_i \alpha$

complete if $KB \vdash_i \alpha$ whenever $KB \models \alpha$

I. Resolution

An inference algorithm i is

sound if $KB \models \alpha$ whenever $KB \vdash_i \alpha$

complete if $KB \vdash_i \alpha$ whenever $KB \models \alpha$

- ◆ Inference rules covered so far are sound.
- ◆ The inference algorithms using them may not be complete.

I. Resolution

An inference algorithm i is

sound if $KB \models \alpha$ whenever $KB \vdash_i \alpha$

complete if $KB \vdash_i \alpha$ whenever $KB \models \alpha$

- ◆ Inference rules covered so far are sound.
- ◆ The inference algorithms using them may not be complete.

resolution + a complete search algorithm = a complete inference algorithm

I. Resolution

An inference algorithm i is

sound if $KB \models \alpha$ whenever $KB \vdash_i \alpha$

complete if $KB \vdash_i \alpha$ whenever $KB \models \alpha$

- ◆ Inference rules covered so far are sound.
- ◆ The inference algorithms using them may not be complete.

resolution + a complete search algorithm = a complete inference algorithm

|

single inference rule

Wumpus World Revisited

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

KB:

$$R_1: \neg P_{1,1}$$

$$R_2: B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})$$

$$R_3: B_{2,1} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{3,1})$$

$$R_4: \neg B_{1,1}$$

$$R_5: B_{2,1}$$

Rules

Wumpus World Revisited

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

KB:

$$R_1: \neg P_{1,1}$$

$$R_2: B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})$$

$$R_3: B_{2,1} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{3,1})$$

$$R_4: \neg B_{1,1}$$

$$R_5: B_{2,1}$$

Rules

Agent: [1,1] → [2,1] → [1,1]

Wumpus World Revisited

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

Agent: [1,1] → [2,1] → [1,1]

KB:

$$R_1: \neg P_{1,1}$$

$$R_2: B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1})$$

$$R_3: B_{2,1} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{3,1})$$

Rules

$$R_4: \neg B_{1,1}$$

$$R_5: B_{2,1}$$

$$R_6: (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1})$$

$$R_7: (P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}$$

$$R_8: \neg B_{1,1} \Rightarrow \neg(P_{1,2} \vee P_{2,1})$$


$$R_9: \neg(P_{1,2} \vee P_{2,1}) \quad // R_4, R_8$$

$$R_{10}: \neg P_{1,2} \wedge \neg P_{2,1}$$

Added to KB via inferences


(cont'd)

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1




(cont'd)

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1



(cont'd)


1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1



[1,1] → [1,2]: stench but no breeze

(cont'd)

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1



$[1,1] \rightarrow [1,2]$: stench but no breeze

Add to KB:

$$R_{11}: \neg B_{1,2}$$

$$R_{12}: B_{1,2} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{1,3})$$

(cont'd)

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

$[1,1] \rightarrow [1,2]$: stench but no breeze

Add to KB:

$$R_{11}: \neg B_{1,2}$$

$$R_{12}: B_{1,2} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{1,3})$$




Similarly, as in deriving R_{10}

$$R_{13}: \neg P_{2,2}$$

$$R_{14}: \neg P_{1,3}$$

(cont'd)

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1



$[1,1] \rightarrow [1,2]$: stench but no breeze

Add to KB:

$$R_{11}: \neg B_{1,2}$$

$$R_{12}: B_{1,2} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{1,3})$$



Similarly, as in deriving R_{10}


$$R_{13}: \neg P_{2,2}$$

$$R_{14}: \neg P_{1,3}$$

$$R_3: B_{2,1} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{3,1})$$

(cont'd)

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1



$[1,1] \rightarrow [1,2]$: stench but no breeze

Add to KB:

$$R_{11}: \neg B_{1,2}$$

$$R_{12}: B_{1,2} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{1,3})$$



Similarly, as in deriving R_{10}

$$R_{13}: \neg P_{2,2}$$

$$R_{14}: \neg P_{1,3}$$

$$R_3: B_{2,1} \Leftrightarrow (P_{1,1} \vee P_{2,2} \vee P_{3,1})$$



biconditional elimination

$$R_5: B_{2,1}$$

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$

Resolvent

$$R_{13}: \neg P_{2,2}$$

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$

Resolvent

$R_{13}: \neg P_{2,2}$

$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$




resolving the two literals that are
negations of each other

Resolvent

$R_{13}: \neg P_{2,2}$

$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$




resolving the two literals that are
negations of each other

$R_{16}: P_{1,1} \vee P_{3,1}$ (*resolvent*)

Resolvent

$R_{13}: \neg P_{2,2}$

$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$



resolving the two literals that are
negations of each other


$R_{16}: P_{1,1} \vee P_{3,1}$ (*resolvent*)

If there's a pit in one of [1,1], [2,2], and [3,1] and it's not in [2,2], then it's in [1,1] or [3,1].

Resolvent

$R_{13}: \neg P_{2,2}$

$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$



resolving the two literals that are
negations of each other


$R_{16}: P_{1,1} \vee P_{3,1}$ (*resolvent*)

If there's a pit in one of [1,1], [2,2], and [3,1] and it's not in [2,2], then it's in [1,1] or [3,1].

Resolvent

$$R_{13}: \neg P_{2,2}$$

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$



$$R_{16}: P_{1,1} \vee P_{3,1} \text{ (*resolvent*)}$$

If there's a pit in one of [1,1], [2,2], and [3,1] and it's not in [2,2], then it's in [1,1] or [3,1].

$$R_1: \neg P_{1,1}$$

$$R_{16}: P_{1,1} \vee P_{3,1}$$

Resolvent

$$R_{13}: \neg P_{2,2}$$

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$




resolving the two literals that are
negations of each other

$$R_{16}: P_{1,1} \vee P_{3,1} \text{ (*resolvent*)}$$

If there's a pit in one of [1,1], [2,2], and [3,1] and it's not in [2,2], then it's in [1,1] or [3,1].

$$R_1: \neg P_{1,1}$$


$$R_{16}: P_{1,1} \vee P_{3,1}$$



Resolvent

$R_{13}: \neg P_{2,2}$

$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$




$R_{16}: P_{1,1} \vee P_{3,1}$ (*resolvent*)

If there's a pit in one of [1,1], [2,2], and [3,1] and it's not in [2,2], then it's in [1,1] or [3,1].

$R_1: \neg P_{1,1}$

$R_{16}: P_{1,1} \vee P_{3,1}$



Simple Resolution Rule

$$\frac{l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m}{l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k}$$

(l_i and m are complementary literals, i.e., $l_i = \neg m$ or $m = \neg l_i$.)

Simple Resolution Rule

$$\frac{l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m}{l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k}$$

(l_i and m are complementary literals, i.e., $l_i = \neg m$ or $m = \neg l_i$.)

Since m is true, then l_i must be false. But one of l_1, \dots, l_k must be true. Therefore, we can exclude l_i and assert that one of the remaining $k - 1$ literals must be true.

Simple Resolution Rule

$$\frac{l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m}{l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k} \quad (l_i \text{ and } m \text{ are complementary literals, i.e., } l_i = \neg m \text{ or } m = \neg l_i.)$$

Since m is true, then l_i must be false. But one of l_1, \dots, l_k must be true. Therefore, we can exclude l_i and assert that one of the remaining $k - 1$ literals must be true.

Clause: a disjunction of literals.

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$

Simple Resolution Rule

$$\frac{l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m}{l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k} \quad (l_i \text{ and } m \text{ are complementary literals, i.e., } l_i = \neg m \text{ or } m = \neg l_i.)$$

Since m is true, then l_i must be false. But one of l_1, \dots, l_k must be true. Therefore, we can exclude l_i and assert that one of the remaining $k - 1$ literals must be true.

Clause: a disjunction of literals.

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$

Simple Resolution Rule

$$\frac{l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m}{l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k} \quad (l_i \text{ and } m \text{ are complementary literals, i.e., } l_i = \neg m \text{ or } m = \neg l_i.)$$

Since m is true, then l_i must be false. But one of l_1, \dots, l_k must be true. Therefore, we can exclude l_i and assert that one of the remaining $k - 1$ literals must be true.

Clause: a disjunction of literals.

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$

Unit clause: a single literal.

Simple Resolution Rule

$$\frac{l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m}{l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k} \quad (l_i \text{ and } m \text{ are complementary literals, i.e., } l_i = \neg m \text{ or } m = \neg l_i.)$$

Since m is true, then l_i must be false. But one of l_1, \dots, l_k must be true. Therefore, we can exclude l_i and assert that one of the remaining $k - 1$ literals must be true.

Clause: a disjunction of literals.

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$

Unit clause: a single literal.

$$R_1: \neg P_{2,2} \qquad R_5: B_{2,1}$$

Simple Resolution Rule

$$\frac{l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m}{l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k} \quad (l_i \text{ and } m \text{ are complementary literals, i.e., } l_i = \neg m \text{ or } m = \neg l_i.)$$

Since m is true, then l_i must be false. But one of l_1, \dots, l_k must be true. Therefore, we can exclude l_i and assert that one of the remaining $k - 1$ literals must be true.

Clause: a disjunction of literals.

$$R_{15}: P_{1,1} \vee P_{2,2} \vee P_{3,1}$$

$$\frac{P_{1,1} \vee P_{2,2} \vee P_{3,1}, \quad \neg P_{2,2}}{P_{1,1} \vee P_{3,1}}$$

Unit clause: a single literal.

$$R_1: \neg P_{2,2}$$

$$R_5: B_{2,1}$$

Full Resolution Rule

l_i and m_j are complementary literals:

$$l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m_1 \vee \cdots \vee m_j \vee \cdots \vee m_k$$

$$l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k \vee m_1 \vee \cdots \vee m_{j-1} \vee m_{j+1} \vee \cdots \vee m_n$$

Full Resolution Rule

l_i and m_j are complementary literals:

$$l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m_1 \vee \cdots \vee m_j \vee \cdots \vee m_k$$

$$l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k \vee m_1 \vee \cdots \vee m_{j-1} \vee m_{j+1} \vee \cdots \vee m_n$$

If l_i is true, then m_j is false. Hence $m_1 \vee \cdots \vee m_{j-1} \vee m_{j+1} \vee \cdots \vee m_n$ must be true.
If l_i is false, then $l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k$ must be true.

Full Resolution Rule

l_i and m_j are complementary literals:

$$l_1 \vee \cdots \vee l_i \vee \cdots \vee l_k, \quad m_1 \vee \cdots \vee m_j \vee \cdots \vee m_k$$

$$l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k \vee m_1 \vee \cdots \vee m_{j-1} \vee m_{j+1} \vee \cdots \vee m_n$$

If l_i is true, then m_j is false. Hence $m_1 \vee \cdots \vee m_{j-1} \vee m_{j+1} \vee \cdots \vee m_n$ must be true.
If l_i is false, then $l_1 \vee \cdots \vee l_{i-1} \vee l_{i+1} \vee \cdots \vee l_k$ must be true.

$$\frac{P_{1,1} \vee P_{3,1}, \quad \neg P_{1,1} \vee \neg P_{2,2}}{P_{3,1} \vee \neg P_{2,2}}$$

One Pair at a Time

Only one pair of complementary literals can be resolved at each step.

$$P \vee \neg Q \vee R, \quad \neg P \vee Q$$

One Pair at a Time

Only one pair of complementary literals can be resolved at each step.

$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{\neg Q \vee R \vee Q}$$

One Pair at a Time

Only one pair of complementary literals can be resolved at each step.

$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{\neg Q \vee R \vee Q \equiv \text{true}}$$

One Pair at a Time

Only one pair of complementary literals can be resolved at each step.

$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{\neg Q \vee R \vee Q \equiv \text{true}}$$



One Pair at a Time

Only one pair of complementary literals can be resolved at each step.

$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{\neg Q \vee R \vee Q \equiv \text{true}}$$



$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{R}$$

One Pair at a Time

Only one pair of complementary literals can be resolved at each step.

$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{\neg Q \vee R \vee Q \equiv \text{true}}$$



$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{R}$$

Incorrect conclusion!

One Pair at a Time

Only one pair of complementary literals can be resolved at each step.

$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{\neg Q \vee R \vee Q \equiv \text{true}}$$



$$\frac{P \vee \neg Q \vee R, \quad \neg P \vee Q}{R}$$



Incorrect conclusion!

Conjunctive Normal Form

The resolution rule applies to clauses only.

Conjunctive normal form (CNF): a conjunction of clauses

Conjunctive Normal Form

The resolution rule applies to clauses only.

Conjunctive normal form (CNF): a conjunction of clauses

$$CNFSentence \rightarrow Clause_1 \wedge \dots \wedge Clause_n$$
$$Clause \rightarrow Literal_1 \vee \dots \vee Literal_m$$
$$Fact \rightarrow Symbol$$
$$Literal \rightarrow Symbol \mid \neg Symbol$$
$$Symbol \rightarrow P \mid Q \mid R \mid \dots$$

Conjunctive Normal Form

The resolution rule applies to clauses only.

Conjunctive normal form (CNF): a conjunction of clauses

$$CNFSentence \rightarrow Clause_1 \wedge \dots \wedge Clause_n$$

$$Clause \rightarrow Literal_1 \vee \dots \vee Literal_m$$

$$Fact \rightarrow Symbol$$

$$Literal \rightarrow Symbol \mid \neg Symbol$$

$$Symbol \rightarrow P \mid Q \mid R \mid \dots$$

Every sentence of propositional logic is equivalent to a CNF.

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\alpha \Leftrightarrow \beta$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\alpha \Leftrightarrow \beta$$

replaced with \downarrow

$$(\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha)$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with } \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \end{array}$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with } \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \end{array}$$

2. Eliminate \Rightarrow .

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with } \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \end{array}$$

2. Eliminate \Rightarrow .

$$\begin{array}{c} \alpha \Rightarrow \beta \\ \downarrow \\ \neg \alpha \vee \beta \end{array}$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with } \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

2. Eliminate \Rightarrow .

$$\begin{array}{c} \alpha \Rightarrow \beta \\ \downarrow \\ \neg \alpha \vee \beta \end{array}$$

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \\ \vdots \\ (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg (P_{1,2} \vee P_{2,1}) \vee B_{1,1}) \end{array}$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with } \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \\ \vdots \\ (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg(P_{1,2} \vee P_{2,1}) \vee B_{1,1}) \end{array}$$

2. Eliminate \Rightarrow .

$$\begin{array}{c} \alpha \Rightarrow \beta \\ \downarrow \\ \neg \alpha \vee \beta \end{array}$$

3. Move \neg inwards, repeatedly applying

$$\begin{array}{l} \neg(\neg \alpha) \equiv \alpha \\ \neg(\alpha \vee \beta) \equiv (\neg \alpha \wedge \neg \beta) \\ \neg(\alpha \wedge \beta) \equiv (\neg \alpha \vee \neg \beta) \end{array}$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with} \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

2. Eliminate \Rightarrow .

$$\begin{array}{c} \alpha \Rightarrow \beta \\ \downarrow \\ \neg \alpha \vee \beta \end{array}$$

3. Move \neg inwards, repeatedly applying

$$\begin{array}{l} \neg(\neg \alpha) \equiv \alpha \\ \neg(\alpha \vee \beta) \equiv (\neg \alpha \wedge \neg \beta) \\ \neg(\alpha \wedge \beta) \equiv (\neg \alpha \vee \neg \beta) \end{array}$$

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \end{array}$$

$$(\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg(P_{1,2} \vee P_{2,1}) \vee B_{1,1})$$

$$(\neg B_{1,1} \vee \neg P_{1,2} \vee P_{2,1}) \wedge ((\neg P_{1,2} \wedge \neg P_{2,1}) \vee B_{1,1})$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with} \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

2. Eliminate \Rightarrow .

$$\begin{array}{c} \alpha \Rightarrow \beta \\ \downarrow \\ \neg \alpha \vee \beta \end{array}$$

3. Move \neg inwards, repeatedly applying

$$\begin{array}{l} \neg(\neg \alpha) \equiv \alpha \\ \neg(\alpha \vee \beta) \equiv (\neg \alpha \wedge \neg \beta) \\ \neg(\alpha \wedge \beta) \equiv (\neg \alpha \vee \neg \beta) \end{array}$$

4. Apply the distributivity law

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \\ \downarrow \\ (\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg(P_{1,2} \vee P_{2,1}) \vee B_{1,1}) \\ \downarrow \\ (\neg B_{1,1} \vee \neg P_{1,2} \vee P_{2,1}) \wedge ((\neg P_{1,2} \wedge \neg P_{2,1}) \vee B_{1,1}) \end{array}$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with} \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

2. Eliminate \Rightarrow .

$$\begin{array}{c} \alpha \Rightarrow \beta \\ \downarrow \\ \neg \alpha \vee \beta \end{array}$$

3. Move \neg inwards, repeatedly applying

$$\begin{array}{l} \neg(\neg \alpha) \equiv \alpha \\ \neg(\alpha \vee \beta) \equiv (\neg \alpha \wedge \neg \beta) \\ \neg(\alpha \wedge \beta) \equiv (\neg \alpha \vee \neg \beta) \end{array}$$

4. Apply the distributivity law

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \end{array}$$

$$(\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg(P_{1,2} \vee P_{2,1}) \vee B_{1,1})$$

$$(\neg B_{1,1} \vee \neg P_{1,2} \vee P_{2,1}) \wedge ((\neg P_{1,2} \wedge \neg P_{2,1}) \vee B_{1,1})$$

$$(\neg B_{1,1} \vee \neg P_{1,2} \vee P_{2,1}) \wedge (\neg P_{1,2} \vee B_{1,1}) \wedge (\neg P_{2,1} \vee B_{1,1})$$

Converting to CNF

1. Eliminate \Leftrightarrow .

$$\begin{array}{c} \alpha \Leftrightarrow \beta \\ \text{replaced with} \downarrow \\ (\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha) \end{array}$$

2. Eliminate \Rightarrow .

$$\begin{array}{c} \alpha \Rightarrow \beta \\ \downarrow \\ \neg \alpha \vee \beta \end{array}$$

3. Move \neg inwards, repeatedly applying

$$\begin{array}{l} \neg(\neg \alpha) \equiv \alpha \\ \neg(\alpha \vee \beta) \equiv (\neg \alpha \wedge \neg \beta) \\ \neg(\alpha \wedge \beta) \equiv (\neg \alpha \vee \neg \beta) \end{array}$$

4. Apply the distributivity law

$$\begin{array}{c} B_{1,1} \Leftrightarrow (P_{1,2} \vee P_{2,1}) \\ \downarrow \\ (B_{1,1} \Rightarrow (P_{1,2} \vee P_{2,1})) \wedge ((P_{1,2} \vee P_{2,1}) \Rightarrow B_{1,1}) \end{array}$$

$$(\neg B_{1,1} \vee P_{1,2} \vee P_{2,1}) \wedge (\neg(P_{1,2} \vee P_{2,1}) \vee B_{1,1})$$

$$(\neg B_{1,1} \vee \neg P_{1,2} \vee P_{2,1}) \wedge ((\neg P_{1,2} \wedge \neg P_{2,1}) \vee B_{1,1})$$

$$(\neg B_{1,1} \vee \neg P_{1,2} \vee P_{2,1}) \wedge (\neg P_{1,2} \vee B_{1,1}) \wedge (\neg P_{2,1} \vee B_{1,1})$$

II. Proof by Resolution – An Example

KB:

P
 $P \rightarrow (Q \vee R)$
 $Q \rightarrow S$
 $R \rightarrow (S \wedge T)$

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S$

$R \rightarrow (S \wedge T)$

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T)$

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T) \quad \text{-----} \rightarrow \neg R \vee (S \wedge T)$

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T) \quad \text{-----} \rightarrow \neg R \vee (S \wedge T)$

$\text{-----} \rightarrow (\neg R \vee S) \wedge (\neg R \vee T)$

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T) \quad \text{-----} \rightarrow \neg R \vee (S \wedge T)$

$\text{-----} \rightarrow (\neg R \vee S) \wedge (\neg R \vee T)$

Q: $KB \vdash S$?

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T) \quad \text{-----} \rightarrow \neg R \vee (S \wedge T)$

$\text{-----} \rightarrow (\neg R \vee S) \wedge (\neg R \vee T)$

Q: $KB \vdash S$?

1. Converting sentences to CNF

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T) \quad \text{-----} \rightarrow \neg R \vee (S \wedge T)$

$\text{-----} \rightarrow (\neg R \vee S) \wedge (\neg R \vee T)$

Q: $KB \vdash S$?

1. Converting sentences to CNF

KB: P

$\neg P \vee Q \vee R$

$\neg Q \vee S$

$(\neg R \vee S) \wedge (\neg R \vee T)$

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T) \quad \text{-----} \rightarrow \neg R \vee (S \wedge T)$

$\text{-----} \rightarrow (\neg R \vee S) \wedge (\neg R \vee T)$

Q: $KB \vdash S$?

1. Converting sentences to CNF
2. Spilt each conjunction into clauses.

KB: P

$\neg P \vee Q \vee R$

$\neg Q \vee S$

$(\neg R \vee S) \wedge (\neg R \vee T)$

II. Proof by Resolution – An Example

KB:

P

$P \rightarrow (Q \vee R) \quad \text{-----} \rightarrow \neg P \vee Q \vee R$

$Q \rightarrow S \quad \text{-----} \rightarrow \neg Q \vee S$

$R \rightarrow (S \wedge T) \quad \text{-----} \rightarrow \neg R \vee (S \wedge T)$

$\text{-----} \rightarrow (\neg R \vee S) \wedge (\neg R \vee T)$

Q: $KB \vdash S$?

1. Converting sentences to CNF
2. Spilt each conjunction into clauses.

KB: P

$\neg P \vee Q \vee R$

$\neg Q \vee S$

$(\neg R \vee S) \wedge (\neg R \vee T)$

$\left\{ \begin{array}{l} \neg R \vee S \\ \neg R \vee T \end{array} \right.$

II. Proof by Resolution – An Example

KB:

P	
$P \rightarrow (Q \vee R)$	$\dashv\vdash \neg P \vee Q \vee R$
$Q \rightarrow S$	$\dashv\vdash \neg Q \vee S$
$R \rightarrow (S \wedge T)$	$\dashv\vdash \neg R \vee (S \wedge T)$
	$\dashv\vdash (\neg R \vee S) \wedge (\neg R \vee T)$

Q: $KB \vdash S$?

1. Converting sentences to CNF
2. Split each conjunction into clauses.

KB: P
 $\neg P \vee Q \vee R$
 $\neg Q \vee S$

$(\neg R \vee S) \wedge (\neg R \vee T)$

KB: P
 $\neg P \vee Q \vee R$
 $\neg Q \vee S$

$\left[\begin{array}{l} \neg R \vee S \\ \neg R \vee T \end{array} \right.$

Proof by Resolution

KB (updated):

- (1) P
- (2) $\neg P \vee Q \vee R$
- (3) $\neg Q \vee S$
- (4) $\neg R \vee S$
- (5) $\neg R \vee T$


$$(1) \quad P$$

$$(2) \quad \neg P \vee Q \vee R$$

Proof by Resolution

KB (updated):


- (1) P
- (2) $\neg P \vee Q \vee R$
- (3) $\neg Q \vee S$
- (4) $\neg R \vee S$
- (5) $\neg R \vee T$



Proof by Resolution

KB (updated):


- (1) P
- (2) $\neg P \vee Q \vee R$
- (3) $\neg Q \vee S$
- (4) $\neg R \vee S$
- (5) $\neg R \vee T$



Proof by Resolution

KB (updated):


- (1) P
- (2) $\neg P \vee Q \vee R$
- (3) $\neg Q \vee S$
- (4) $\neg R \vee S$
- (5) $\neg R \vee T$



Proof by Resolution

KB (updated):


- (1) P
- (2) $\neg P \vee Q \vee R$
- (3) $\neg Q \vee S$
- (4) $\neg R \vee S$
- (5) $\neg R \vee T$



Proof by Resolution

KB (updated):


- (1) P
- (2) $\neg P \vee Q \vee R$
- (3) $\neg Q \vee S$
- (4) $\neg R \vee S$
- (5) $\neg R \vee T$



Proof by Resolution

KB (updated):

- (1) P
- (2) $\neg P \vee Q \vee R$
- (3) $\neg Q \vee S$
- (4) $\neg R \vee S$
- (5) $\neg R \vee T$



Resolution tree

Resolution Refutation

(Proof by contradiction)

To show that $KB \models \alpha$, we show that $KB \wedge \neg\alpha$ is unsatisfiable. .

Resolution Refutation

(Proof by contradiction)

To show that $KB \models \alpha$, we show that $KB \wedge \neg\alpha$ is unsatisfiable. .

KB (about a summer day):

- (1) If it is raining and you are outside then you will get wet.
- (2) If it is warm and there is no rain then it is a pleasant day.
- (3) You are not wet.
- (4) You are outside.
- (5) It is a warm day.

Resolution Refutation

(Proof by contradiction)

To show that $KB \models \alpha$, we show that $KB \wedge \neg\alpha$ is unsatisfiable. .

KB (about a summer day):

- (1) If it is raining and you are outside then you will get wet.
- (2) If it is warm and there is no rain then it is a pleasant day.
- (3) You are not wet.
- (4) You are outside.
- (5) It is a warm day.

Prove

It is a pleasant day.

KB in Propositional Sentences

KB (rewritten):

- (1) $(\text{rain} \wedge \text{outside}) \Rightarrow \text{wet}$
- (2) $(\text{warm} \wedge \neg \text{rain}) \Rightarrow \text{pleasant}$
- (3) $\neg \text{wet}$
- (4) outside
- (5) warm

KB in Propositional Sentences

KB (rewritten):

```
(1) ( rain  $\wedge$  outside )  $\Rightarrow$  wet  
(2) ( warm  $\wedge$   $\neg$ rain )  $\Rightarrow$  pleasant  
(3)  $\neg$ wet  
(4) outside  
(5) warm
```



converted into clauses

```
(1)  $\neg$ rain  $\vee$   $\neg$ outside  $\vee$  wet  
(2)  $\neg$ warm  $\vee$  rain  $\vee$  pleasant  
(3)  $\neg$ wet  
(4) outside  
(5) warm
```

KB in Propositional Sentences

KB (rewritten):

```
(1) ( rain  $\wedge$  outside )  $\Rightarrow$  wet  
(2) ( warm  $\wedge$   $\neg$ rain )  $\Rightarrow$  pleasant  
(3)  $\neg$ wet  
(4) outside  
(5) warm
```



converted into clauses

```
(1)  $\neg$ rain  $\vee$   $\neg$ outside  $\vee$  wet  
(2)  $\neg$ warm  $\vee$  rain  $\vee$  pleasant  
(3)  $\neg$ wet  
(4) outside  
(5) warm
```


We add \neg pleasant to KB and try to derive false.

Resolution Refutation Tree


$\neg \text{pleasant}$

(2) $\neg \text{warm} \vee \text{rain} \vee \text{pleasant}$


Resolution Refutation Tree




Resolution Refutation Tree




Resolution Refutation Tree




Resolution Refutation Tree




Resolution Refutation Tree




Resolution Refutation Tree




Resolution Refutation Tree




Resolution Refutation Tree




Resolution Refutation Tree



Resolution Refutation Tree



Proving $\neg P_{1,2}$ in the Wumpus World



1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

Resolution Algorithm

```
function PL-RESOLUTION( $KB, \alpha$ ) returns true or false
  inputs:  $KB$ , the knowledge base, a sentence in propositional logic
            $\alpha$ , the query, a sentence in propositional logic

   $clauses \leftarrow$  the set of clauses in the CNF representation of  $KB \wedge \neg\alpha$ 
   $new \leftarrow \{ \}$ 
  while true do
    for each pair of clauses  $C_i, C_j$  in  $clauses$  do
       $resolvents \leftarrow$  PL-RESOLVE( $C_i, C_j$ )
      if  $resolvents$  contains the empty clause then return true
       $new \leftarrow new \cup resolvents$ 
    if  $new \subseteq clauses$  then return false // no new clauses can be added.
   $clauses \leftarrow clauses \cup new$ 
```

The process ends in one of two situations below:

- ◆ No new clauses can be added, in which case KB does not entail α ;
- ◆ Two clauses resolve to yield the empty clause, in which case KB entails α .

Completeness of Resolution

Given a set of clauses S , its *resolution closure* $RC(S)$ includes all the clauses in S as well as all the resolvents from repeated applications of the resolution rule.

Completeness of Resolution

Given a set of clauses S , its *resolution closure* $RC(S)$ includes all the clauses in S as well as all the resolvents from repeated applications of the resolution rule.

$RC(S)$ is *finite* because only 3^n distinct clauses can be constructed out of n propositional symbols appearing in S .

Completeness of Resolution

Given a set of clauses S , its *resolution closure* $RC(S)$ includes all the clauses in S as well as all the resolvents from repeated applications of the resolution rule.

$RC(S)$ is *finite* because only 3^n distinct clauses can be constructed out of n propositional symbols appearing in S .

Ground Resolution Theorem: If S is unsatisfiable, then $RC(S)$ contains the empty clause \emptyset .

Completeness of Resolution

Given a set of clauses S , its *resolution closure* $RC(S)$ includes all the clauses in S as well as all the resolvents from repeated applications of the resolution rule.

$RC(S)$ is *finite* because only 3^n distinct clauses can be constructed out of n propositional symbols appearing in S .

Ground Resolution Theorem: If S is unsatisfiable, then $RC(S)$ contains the empty clause \emptyset .

Constructive proof by explicitly generating an assignment for S if $\emptyset \notin RC(S)$.

Completeness of Resolution

Given a set of clauses S , its *resolution closure* $RC(S)$ includes all the clauses in S as well as all the resolvents from repeated applications of the resolution rule.

$RC(S)$ is *finite* because only 3^n distinct clauses can be constructed out of n propositional symbols appearing in S .

Ground Resolution Theorem: If S is unsatisfiable, then $RC(S)$ contains the empty clause \emptyset .

Constructive proof by explicitly generating an assignment for S if $\emptyset \notin RC(S)$.

Completeness of Resolution

Given a set of clauses S , its *resolution closure* $RC(S)$ includes all the clauses in S as well as all the resolvents from repeated applications of the resolution rule.

$RC(S)$ is *finite* because only 3^n distinct clauses can be constructed out of n propositional symbols appearing in S .

Ground Resolution Theorem: If S is unsatisfiable, then $RC(S)$ contains the empty clause \emptyset .

Constructive proof by explicitly generating an assignment for S if $\emptyset \notin RC(S)$.