## Stat 330 Homework 2

Sean Gordon

January 27, 2020

$$\begin{array}{l} (b) \ A = \{T \ T \ H, \ T \ H \ T, \ H \ T \ T \} \\ B = \{T \ T \ H, \ T \ H \ T, \ H \ T \ T, \ T \ T \ T \} \\ C = \{T \ H \ H, \ H \ H \ H \} \\ \end{array}$$

(c) 
$$\overline{A} = \{ T \ T \ T, \ T \ H \ H, \ H \ T \ H, \ H \ H \ T, \ H \ H \ H \}$$
  
 $A \cup B = \{ T \ T \ H, \ T \ H \ T, \ H \ T \ T \ T \}$   
 $A \cap B = \{ T \ T \ H, \ T \ H \ T, \ H \ T \ T \}$   
 $A \cap C = \{ \}$ 

2) 
$$A = \{1, 3, 5\}$$

$$B = \{1, 5, 10\}$$

$$\overline{A} = \{2, 4, 6, 7, 8, 9, 10\}$$

$$\overline{B} = \{2, 3, 4, 6, 7, 8, 9\}$$

(a) 
$$\underline{A \cap B} = \{1, 5\}$$
  
 $\underline{\overline{A \cap B}} = \{2, 3, 4, 6, 7, 8, 9, 10\}$   
 $\overline{A} \cup \overline{B} = \{2, 3, 4, 6, 7, 8, 10\}$ 

(b) 
$$\underline{A \cup B} = \{1, 3, 5, 10\}$$
  
 $\underline{\overline{A \cup B}} = \{2, 4, 6, 7, 8, 9\}$   
 $\overline{\overline{A} \cap \overline{B}} = \{2, 4, 6, 7, 8, 9\}$ 

3) Total probability sums to 1.0,  $\therefore$  1k + 2k + 3k + 4k + 5k + 6k = 21k = 1.0  $k = 1.0 / 21 = 0.0476 \Rightarrow$  [1] = 1\*0.0476 = 0.0476 [2] = 2\*0.0476 = 0.0952 [3] = 3\*0.0476 = 0.1429 [4] = 4\*0.0476 = 0.1905 [5] = 5\*0.0476 = 0.2381

4)
(a) 
$$\{(1,1),(1,2),(1,3),(1,4),(1,5),(1,6),(2,1),(2,2),(2,3),(2,4),(2,5),(2,6),(3,1),(3,2),(3,3),(3,4),(3,5),(3,6),(4,1),(4,2),(4,3),(4,4),(4,5),(4,6),(5,1),(5,2),(5,3),(5,4),(5,5),(5,6),(6,1),(6,2),(6,3),(6,4),(6,5),(6,6)\}$$

[6] = 6\*0.0476 = 0.2857

- (b) 6 possible options for '7' out of 36 possible outcomes: 6/36 = 1/6 chance
- (c) 6 possible options for '7' out of 36 possible outcomes, 2 possible options for '11' out of 36 possible outcomes: Cannot get a 7 and an 11, 8/36 = 2/9 chance
- (d) 18 possible options for even first die out of 36 possible outcomes, 2 possible options for '11' out of 36 possible outcomes, with 1 overlap  $\Rightarrow$  1 option: 7/36 chance

- 5) (a)  $P(MB \cup HD) = 35 + 30 20 = 45\%$ 
  - (b) 1 P(MB  $\cup$  HD) = 100 45 = 65%
- 6) (a)  $P(L \cap S) = 40 + 30 50 = 20\%$ 
  - (b)  $P(L \cup S) = 100 50 = 50\%$
  - (c)  $P(L \text{ xor } S) = 100 50 P(L \cap S) = 30\%$