

humans manifests itself as mild symptoms to severe respiratory failure. On binding to epithelial cells in the respiratory tract, SARS-CoV-2 starts replicating and migrating down to the airways and enters alveolar epithelial cells in the lungs. The rapid replication of SARS-CoV-2 in the lungs may trigger a strong immune response. Cytokine storm syndrome causes acute respiratory distress syndrome and respiratory failure, which is considered the main cause of death in patients with COVID-19 (REFS^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}!). Patients of older age (>60 years) and with serious pre-existing diseases have a greater risk of developing acute respiratory distress syndrome and death" (FIG. 4). Multiple organ failure has also been reported in some COVID-19 cases^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}. Histopathological changes in patients with COVID-19 occur mainly in the lungs. Histopathology analyses showed bilateral diffused alveolar damage, hyaline membrane formation, desquamation of pneumocytes and fibrin deposits in lungs of patients with severe COVID-19. Exudative inflammation was also shown in some cases. Immunohistochemistry assays detected SARS-CoV-2 antigen in the upper airway, bronchiolar epithelium and submucosal gland epithelium, as well as in type I and type II pneumocytes, alveolar macrophages and hyaline membranes in the lungs^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}. Animal models used for studying SARS-CoV-2 infection pathogenesis include non-human primates (rhesus macaques, cynomolgus monkeys, marmosets and African green monkeys), mice (wild-type mice (with mouse-adapted virus) and human ACE2-transgenic or human ACE2-knock-in mice), ferrets and golden hamsters^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}. In non-human primate animal models, most species display clinical features similar to those of patients with COVID-19, including virus shedding, virus replication and host responses to SARS-CoV-2 infection^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100}. For example, in the rhesus macaque model, high viral loads were detected in the upper and