

Weak Lensing for Precision Cosmology

Mohamed Shaaban

April 2019

1 Introduction

The most revolutionary discovery in cosmology since Hubble observed that the Universe is expanding is that this expansion is accelerating. A revelation that was awarded the 2011 Nobel Prize for its profound implications. [1]. An accelerating universe implies that either our understanding of gravity is flawed or that a mysterious negative pressure known as Dark Energy is driving the expansion [12]. This Dark Energy accounts for most (over 68%) of the energy density in the observable universe, however its origin and physics are presently unknown [2]. As a result, the nature of Dark Energy is considered one of the greatest mysteries of modern science [3].

One of the most powerful ways to probe Dark Energy, as well as modified theories of gravity, is a technique known as weak gravitational lensing, or weak lensing for short [5, 8]. Gravitational lensing is the phenomenon of light ray deflection by intervening mass. When the deflection is sufficiently weak, this phenomenon manifests in images of galaxies as a shearing effect due to the differential deflection of neighboring light rays [16, 5]. This shearing induces a subtle (sub 1%) change in the ellipticity of the images. Although such a change is negligible in comparison to the 30% dispersion in intrinsic galaxy ellipticities, it can be statistically measured by using the coherence of the lensing shear over the sky [16]. The reason weak lensing is considered very powerful is because it provides a direct measurement of the matter distribution in the universe as a function of redshift independent of any cosmological assumptions. Thus, allowing us to directly probe the growth of cosmic structure with time [5].

In this report I present a general overview of weak lensing in the context of cosmology. I begin by presenting the theory behind the Cosmology we would like to probe, as well as the general theoretical framework on which weak lensing operates. I then present the weak lensing measurement procedure and how cosmological data can be extracted from such measurements. Finally, I end this report by presenting some results from an active experiment and a short discussion on potentially concerning systematics.

2 Background Theory

2.1 Standard Model of Cosmology

The fundamental assumption in cosmology, known as the cosmological principle, is that we live in a homogenous (independent of position) and isotropic (independent of direction) universe [16, 14]. Solving Einstein's equations under the geometric symmetries provided by the cosmological principle and taking into account the expansion of the universe one finds that the dynamics of the universe are governed by the Friedman equation [14, 16],

$$H^2(z) = H_0^2 \left(\Omega_r(1+z)^4 + \Omega_m(1+z)^3 + \Omega_k(1+z)^2 + \Omega_\Lambda \frac{\rho_\Lambda(z)}{\rho_\Lambda(0)} \right) \quad (1)$$

Remember to show the Standard Model parameters! can be found in wikipedia.

Angular diameter distance is the one of relevance in lensing (distance measures in cosmology) Distances in cosmo [6] lol

For current and future surveys, one goal is to use the redshifts of the background galaxies (often approximated using photometric redshifts) to divide the survey into multiple redshift bins. The low-redshift bins will only be lensed by structures very near to us, while the high-redshift bins will be lensed by structures over a wide range of redshift. This technique, dubbed "cosmic tomography", makes it possible to map out the 3D distribution of mass. Because the third dimension involves not only distance but cosmic time, tomographic weak lensing is sensitive not only to the matter power spectrum today, but also to its evolution over the history of the universe, and the expansion history of the universe during that time. This is a much more valuable cosmological probe, and many proposed experiments to measure the properties of dark energy and dark matter have focused on weak lensing, such as the Dark Energy Survey, Pan-STARRS, and Large Synoptic Survey Telescope.

1. Motivation (modified gravity vs Dark energy) is DE constant in space ? and time ?
2. FRW Standard Cosmology
3. Distance Measures In Cosmo
4. Gravitational Evolution of Dark matter
5. Observables and Parameters

2.1.1 Matter Power Spectrum

[15]

2.2 Bending of Light

The fundamental concept on which weak lensing is built is gravity's ability to alter the path of a photon. This phenomenon is explored in full detail in [13, 7, 10]. In this subsection I present a simple overview of the theory behind the bending of light necessary to develop the weak lensing formalism. For more detailed calculations consult [13, 7, 10].

2.2.1 Newtonian Lens

It is a common misconception that the gravitational bending of light is an exclusive property of GR. However, gravity induced alterations to a photon's path are predicted by newtonian mechanics [11]. To illustrate this consider a mass M located at the origin of the cartesian plane and a corpuscle(newtonian photon) propagating along the $x = b$ line (in this context b is known as the impact parameter). Newton's second law predicts that the presence of the point mass will result in a momentum transfer between the two objects. If the corpuscle starts with momentum $(p, 0)$ then it will end up with momentum (p_x, p_y) . Therefore, the particle path is deflected by some angle $\Delta\theta$. The deflection angle is simply given by

$$\sin(\Delta\theta) = \frac{p_y}{\sqrt{p_x^2 + p_y^2}} \quad (2)$$

For very small deflections we have $p \approx p_x \gg p_y$ and $\Delta\theta \ll 1$. Therefore Equation 2 simplifies to $\Delta\theta \approx \frac{p_y}{p_x}$. We now consider the infinitesimal deflection along the entire path of the photon with $d\Delta\theta = \frac{dp_y}{p_x} = \frac{1}{p_x} dx \frac{dp_y}{dx}$. Therefore, we can find the deflection angle by

$$\begin{aligned} \Delta\theta_N &= -\frac{1}{p_x} \int dx \frac{dp_y}{dx} \\ &= -\frac{1}{cp_x} \int dx \frac{dp_y}{dt} \\ &= \frac{2GM}{c^2 b} \end{aligned} \quad (3)$$

We note that the mass of the corpuscle cancels out of the deflection equation. Therefore this equation applies for massless particles i.e. photons. Therefore Equation 3 provides a newtonian description for the bending of light [11].

2.2.2 General Relativistic Bending of Light

The Einstein's field equations in the presence of a charge free static point mass is uniquely solved by the Schwarzschild metric [13]. The Schwarzschild metric is

$$ds^2 = \left(1 - \frac{r_s}{r}\right) dt^2 - \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 - r^2 d\Omega^2 \quad (4)$$

where r_s is the Shewarzchild radius of the system given by $r_s = 2\mu = 2GM/c^2$ and (t, r, Ω) are the standard parameters for 4D space-time in polar coordinates. We can analyze the path of the photon from subsection 2.2.1 by studying the geodesic equations of the metric and finding the conserved quantities of the system. We can then combine the conservation equations with the tangent vector norm condition for a null path to get the shape equation of the system as

$$\frac{d\phi}{dr} = \frac{1}{r^2} \left(\frac{1}{b^2} - \frac{1}{r^2} \left(1 - \frac{2\mu}{r} \right) \right)^{-1/2} \quad (5)$$

where (r, ϕ) are the photons position in 2D polar coordinates and b is the impact parameter. Rewriting this equation under the transformation of $r = 1/u$ and working pertrubatively around $u(\mu = 0) = \frac{1}{b} \sin \phi$ we get

$$u(\phi) \approx \frac{1}{b} \sin \phi + \frac{3\mu}{2b^2} \left(1 + \frac{1}{3} \cos 2\phi \right) \quad (6)$$

in the limit were $\phi \ll 1$ and $u \rightarrow 0$ Equation 6 simplifies to $\phi = \Delta\theta_N = \frac{2GM}{c^2 b}$. Geometrically the deflection is given by $\Delta\theta = 2\phi$ and therefore the deflection angle is

$$\Delta\theta = 2\Delta\theta_N = \frac{4GM}{c^2 b} \quad (7)$$

We conclude that general relativity predicts a factor of 2 greater deflection form a point mass than is predicted by newtonian mechanics. This relationship greatly simplifies the formalism developed for weak lensing.

2.3 Weak Lensing Formalism

Now that we have a theoretical understanding of the Cosmological parameters we would like to measure, as well as an understanding of how gravitational fields impact the trajectory of light. We are ready to develop the theoretical formalism on which all weak lensing applications are built.

In order to develop our formalism let us consider a general lensing system as seen in Figure 1. For astronomical applications on cosmological scales the distance from the observer to the lens D_L , the distance from the lens to the source D_{SL} , and the distance from the observer to the source D_S are much greater than the thickness of the lens along the optical axis. Therefore we can treat the lens as a "thin lens", i.e. it lives on a planar slice along the line of sight. We project the mass and potential of the lens onto the lensing plane by defining the projected surface density Σ and projected potential Ψ as

$$\begin{aligned} \Sigma(x, y) &= \int \rho(x, y, z) dz \\ \Psi &= \int \psi dz \end{aligned} \quad (8)$$




Figure 1: Sketch of a thin lens system highlighting the parameters of relevance in the weak lensing formalism. It is conventional to assume the planes are orthogonal to the z axis.

were ρ and ψ are the

Talk about thin lens and how the formalism revolves around projecting newtonian potential onto the lens 2D surface.[7]

3 Measuring Shear / Weak Lensing Pipeline

4 Getting Cosmology from Weak Lensing

[11] [8] [5]

5 Weak Lensing Results

[4]




Figure 2: The effects of the convergence κ and the shear γ on a circular image of a galaxy. The black line represents the nominal image, orange represents positive parameter values, and green represents negative parameter values.

6 Problems With Weak Lensing

[9]

1. Current observations imply $w=-1$ however that is no constraint on its evolution
2. The initial fluctuation in matter density produced by random quantum events therefore CLT tells us that is a gaussian random field therefore power spectrum is enough to get all statistics

References

- [1] The nobel prize in physics 2011 - advanced information. nobelprize.org., 2014.
- [2] P. A. R. Ade et al. Planck 2013 results. I. Overview of products and scientific results. *Astron. Astrophys.*, 571:A1, 2014.
- [3] K. Bandura, G. E. Addison, M. Amiri, J. R. Bond, D. Campbell-Wilson, L. Connor, J.-F. Cliche, G. Davis, M. Deng, N. Denman, M. Dobbs, M. Fandino, K. Gibbs, A. Gilbert, M. Halpern, D. Hanna, A. D. Hincks, G. Hinshaw, C. Höfer, P. Klages, T. L. Landecker, K. Masui, J. Mena Parra, L. B. Newburgh, U.-l. Pen, J. B. Peterson, A. Recnik, J. R. Shaw, K. Sigurdson, M. Sitwell, G. Smecher, R. Smegal, K. Vanderlinde, and D. Wiebe. Canadian Hydrogen Intensity Mapping Experiment (CHIME) pathfinder. In *Ground-based and Airborne Telescopes V*, volume 9145, page 914522, July 2014.

- [4] Chiaki Hikage, Masamune Oguri, Takashi Hamana, Surhud More, Rachel Mandelbaum, Masahiro Takada, Fabian Köhlinger, Hironao Miyatake, Atsushi J. Nishizawa, Hiroaki Aihara, Robert Armstrong, James Bosch, Jean Coupon, Anne Ducout, Paul Ho, Bau-Ching Hsieh, Yutaka Komiyama, François Lanusse, Alexie Leauthaud, Robert H. Lupton, Elinor Medezinski, Sogo Mineo, Shoken Miyama, Satoshi Miyazaki, Ryoma Murata, Hitoshi Murayama, Masato Shirasaki, Cristóbal Sifón, Melanie Simet, Joshua Speagle, David N. Spergel, Michael A. Strauss, Naoshi Sugiyama, Masayuki Tanaka, Yousuke Utsumi, Shiang-Yu Wang, and Yoshihiko Yamada. Cosmology from cosmic shear power spectra with Subaru Hyper Suprime-Cam first-year data. *Publications of the Astronomical Society of Japan*, page 22, Mar 2019.
- [5] H. Hoekstra and B. Jain. Weak Gravitational Lensing and Its Cosmological Applications. *Annual Review of Nuclear and Particle Science*, 58:99–123, November 2008.
- [6] David W. Hogg. Distance measures in cosmology. 1999.
- [7] Konrad Kuijken. The Basics of Lensing. *arXiv e-prints*, pages astro-ph/0304438, Apr 2003.
- [8] Rachel Mandelbaum. Weak Lensing for Precision Cosmology. *Annual Review of Astronomy and Astrophysics*, 56:393–433, Sep 2018.
- [9] Richard Massey, Henk Hoekstra, Thomas Kitching, Jason Rhodes, Mark Cropper, Jérôme Amiaux, David Harvey, Yannick Mellier, Massimo Meneghetti, Lance Miller, Stéphane Paulin-Henriksson, Sandrine Pires, Roberto Scaramella, and Tim Schrabback. Origins of weak lensing systematics, and requirements on future instrumentation (or knowledge of instrumentation). *mnras*, 429:661–678, Feb 2013.
- [10] Yannick Mellier. Probing the universe with weak lensing. *Ann. Rev. Astron. Astrophys.*, 37:127–189, 1999.
- [11] G. Meylan, P. Jetzer, P. North, P. Schneider, C. S. Kochanek, and J. Wambsganss, editors. *Gravitational Lensing: Strong, Weak and Micro*, 2006.
- [12] P. J. E. Peebles and Bharat Ratra. The Cosmological constant and dark energy. *Rev. Mod. Phys.*, 75:559–606, 2003.
- [13] A.W. Peet. General Relativity Notes. page <https://ap.io/483f/files/gr1.pdf>, 2018.
- [14] B. Ryden. *Introduction to Cosmology*. Cambridge University Press, 2016.
- [15] P. Schneider. *Extragalactic Astronomy and Cosmology*. 2006.
- [16] David H. Weinberg, Michael J. Mortonson, Daniel J. Eisenstein, Christopher Hirata, Adam G. Riess, and Eduardo Roza. Observational probes of cosmic acceleration. *physrep*, 530:87–255, Sep 2013.