

```
In [4]: import random as rnd

import matplotlib.pyplot as plt

import numpy as np

import math

from scipy.stats import norm

from PIL import Image
from IPython.display import display
```

I worked with the wonderful \_\_!~!!Michael Scutari!!! on this project

And also the great Peter Banyas \_\_ ^^

## Question 1

```
In [5]: img = Image.open('q1.png')

display(img)
```

### Airline Overbooking

Posted on [September 11, 2012](#) by [Jonathan Mattingly](#) | [Comments Off](#)

An airline knows that over the long run, 90% of passengers who reserve seats for a flight show up. On a particular flight with 300 seats, the airline sold 324 reservations.

1. Assuming that passengers show up independently of each other, what is the chance that the flight will be overbooked ?
2. Suppose that people tend to travel in groups. Would that increase or decrease the probability of overbooking ? Explain your answer.
3. Redo the calculation in the first question assuming that passengers always travel in pairs. Are your answers to all three questions consistent ?

### Q1, part 1

```
In [6]: img = Image.open('q1_part1.png')

display(img)
```

1) 90 % passengers. 300 seats 324 reservations.

$$\therefore P(\text{person appears}) = p = 0.9$$

$$\therefore P(\text{More than 300 people appear}) = P(\text{overbooked})$$

$$\text{let } X \sim \text{Binomial}(\overset{\uparrow}{324}, \overset{\uparrow}{0.9})$$

$$\therefore P = P(X > 300) = P(X \geq 301)$$

$$= \sum_{k=301}^{324} \binom{324}{k} \cdot (0.9)^k \cdot (0.1)^{324-k}$$

Normal approximation:

$$\mu = np$$

$$\sigma = \sqrt{npq} = \sqrt{np(1-p)}$$

$$\therefore \alpha = \frac{a - 1/2 - \mu}{\sigma}, \quad \beta = \frac{b + 1/2 - \mu}{\sigma}$$

$$\therefore \int_{\alpha}^{\beta} \frac{1}{\sqrt{2\pi}} e^{-1/2 z^2} dz = \Phi(\beta) - \Phi(\alpha)$$

$$\approx 0.0497 \quad (3sf)$$

$$\therefore P(X > 300) \approx \underline{\underline{0.0497}}$$

In [7]: `#Q1 normal approx.`

```
def normalApproxWithContinuityCorrection(n, p, a, b):  
  
    mu = n * p  
    sigma = (n * p * (1 - p)) ** 0.5  
  
    #shifts  
    a = a - 0.5  
    b = b + 0.5  
  
    return norm.cdf(b, loc=mu, scale=sigma) - norm.cdf(a, loc=mu, scale=sigma)
```

In [8]: `res = normalApproxWithContinuityCorrection(n=324, p=0.9, a=301, b=324)`  
`print(str(res))`

0.049661136484298374

## Q1, part 2

In [9]: `img = Image.open('q1_part2.png')`  
`display(img)`

2) Another binomial  $Y \sim B\left(\frac{324}{9}, 0.9\right)$

$g$  = group size (only valid for values  
of  $g$  that go perfectly into 0.9)

$$\begin{aligned} P(Y > \frac{300}{9}) & \quad \text{let } N_{\text{groups}} = \frac{324}{9} \\ &= P(Y_g > 300) = 1 - P(Y_g \leq 300) \\ &= \sum_{k=\frac{300}{9}+1}^{\frac{324}{9}} \binom{\frac{324}{9}}{k} \cdot (0.9)^k \cdot (0.1)^{\frac{324}{9}-k} \end{aligned}$$

Binomial approximation:

$$\mu = \frac{324}{9} \times 0.9$$

$$\sigma = \sqrt{\frac{324}{g} \times 0.9 \times 0.1}$$

$$\alpha = \frac{a - \frac{1}{2} - \mu}{\sigma}$$

$$\int_{\beta}^{\alpha} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2} dz$$

$$\beta = \frac{b + \frac{1}{2} - \mu}{\sigma}$$

$\therefore$  I expect the range of the bounds  $(\alpha, \beta)$  to increase ( $\because \sigma$  gets smaller as  $g$  increases).

$\therefore$  as  $g$  gets larger, I expect it to be more and more probable that the flight is overbooked.

```

In [10]: ### image of part two

#group size values that can divide 324 exactly
g_vals = [1, 2, 3, 4, 6, 9, 12, 18, 27, 36, 54, 81, 108, 162, 324]

#this array will contain the probabilities of being overbooked for each group
probs = []

for g in g_vals:

    n_in = 324//g
    p_in = 0.9


    a_in = (300//g) + 1
    b_in = 324//g

    probs.append(normalApproxWithContinuityCorrection(n=n_in, p=p_in, a=a_in, b=b_in))

# Plot the result
plt.figure(figsize=(12, 6))
plt.title("Probability of Overbooking against Group Size")
plt.plot(g_vals, probs, marker='o')
plt.ylabel("Probability of Overbooking")
plt.xlabel("Group Size")
plt.grid(True)
plt.show()

# Print the result as a rounded percentage with 2 decimal places
print("Group Size | Probability of Overbooking")
print("-----")
for i in range(len(g_vals)):
    print(f"{g_vals[i]:<10} | {probs[i]*100:.2f}%")

```





Group Size	Probability of Overbooking
1	4.97%
2	10.92%
3	14.48%
4	16.71%
6	19.07%
9	25.92%
12	20.07%
18	37.15%
27	19.94%
36	26.85%
54	37.87%
81	49.94%
108	58.80%
162	71.08%
324	88.60%

### Q1, part 3

```
In [11]: img = Image.open('q1_part3.png')
display(img)
```

3) As can be seen in code above  
for case group size,  $g=2$

the probability of being overbooked = 10.92%.

$\therefore$  It agrees with my observation in part 2

### Question 2

```
In [12]: img = Image.open('q2.png')
display(img)
```

## Leukemia Test

Posted on August 13, 2012 by Jonathan Mattingly | [Leave a comment](#)

A new drug for leukemia works 25% of the time in patients 55 and older, and 50% of the time in patients younger than 55. A test group has 17 patients 55 and older and 12 patients younger than 55.

1. A uniformly random patient is chosen from the test group, and the drug is administered and it is a success. What is the probability the patient was 55 and older?
2. A subgroup of 4 patients are chosen and the drug is administered to each. What is the probability that the drug works in all of them?

```
In [13]: img = Image.open('q2_part1.png')
display(img)
```

1)

	12 younger than 55	17 older than 55
Drug works	$\frac{1}{2}$	$\frac{1}{4}$
Drug does not work	$\frac{1}{2}$	$\frac{3}{4}$

$$\begin{aligned} P(\text{older 55} \mid \text{success}) &= \frac{P(A)}{P(B)} P(B|A) \\ &= \frac{P(\text{older 55})}{P(\text{success})} \cdot P(\text{success} \mid \text{older 55}) \end{aligned}$$

$$\begin{aligned} P(\text{success}) &= P(\text{success} \mid \text{older 55}) P(\text{older 55}) \\ &\quad + P(\text{success} \mid \text{younger 55}) P(\text{younger 55}) \\ &= \frac{1}{4} \cdot \frac{17}{29} + \frac{1}{2} \cdot \frac{12}{29} = \frac{41}{116} \end{aligned}$$

$$\therefore P(\text{older 55} \mid \text{success}) = \frac{17}{29} \cdot \frac{116}{41} \cdot \frac{1}{4} = \underline{\underline{\frac{17}{41}}} \quad (\approx 0.415)$$

```
In [14]: img = Image.open('q2_part2.png')
display(img)
```

2)  $P(\text{drug works on all of them})$ .

$$= P(\text{drug works on } P_1) \cdot P(\text{drug works on } P_2) \\ \cdot P(\text{drug works on } P_3) \cdot P(\text{drug works on } P_4)$$

Assuming we know nothing about which group each person is in, have to assume independent.

$$\therefore = P(\text{drug works on a person})^4$$

$$= \left( \frac{41}{116} \right)^4 \approx \underline{\underline{0.0156}}$$

from previous q.  $\uparrow$

NOTE: Prof Mattingly mentioned that we have not learnt appropriate tools to do this  $\therefore$  this may be incorrect.

## Question 3

```
In [15]: img = Image.open('q3.png')
display(img)
```



### Exercise 5.1: Using the Normal Approximation to Binomial

Write code to solve each of the following problems concerning rolls of a fair die. Organize your results in a nice table.

1. What is the exact probability of rolling a 6-sided die an even number more than 70% of the time in 10 rolls, 100 rolls, 1000 rolls.
2. What is the exact probability of rolling the number 1 more than 20% of the time in 10 rolls, 100 rolls, 1000 rolls of a 6-sided die.
3. What is the exact probability of rolling the number 1 more than 9% of the time in 10 rolls, 100 rolls, 1000 rolls of a 20-sided die.
4. Approximate each of the above probabilities using a normal approximation with and without the continuity correction. (You should use the continuity correction at the level of the number of rolls and not the percentages).

Discuss any trends you see. How does the approximation behave as  $n$  increases or as the probabilities of the events decrease? How does the region you are integrating the normal curve over change as  $n$  increase? Does this have any implications for the approximation by a normal?

```
In [16]: img = Image.open('q3_parts1-3.png')
display(img)
```

$$1) \quad X \sim B(n, p)$$

$$P(X=k) = \binom{n}{k} \cdot (p)^k \cdot (q)^{n-k}$$

$$p = 0.5 \quad \therefore P(\text{even}) = \frac{3}{6} = \frac{1}{2}.$$

$$P(X > 0.7n) = \sum_{k=0.7n+1}^n \binom{n}{k} (p)^k (q)^{n-k}$$

$$2) \quad X \sim B(n, \frac{1}{6})$$

$$P(X > 0.2n) = \sum_{k=0.2n+1}^n \binom{n}{k} \left(\frac{1}{6}\right)^k \left(\frac{5}{6}\right)^{n-k}$$

$$3) \quad X \sim B(n, \frac{1}{20})$$

$$P(X > 0.09n) = \sum_{k=0.09n+1}^n \binom{n}{k} \left(\frac{1}{20}\right)^k \left(\frac{19}{20}\right)^{n-k}$$

### Q3, part 1

```
In [17]: def numEven(n):
# n = num trials
# p = probability of success on each trial

running_prob = 0

for i in range(math.ceil((0.7*n) + 1), n+1):
    running_prob += math.comb(n, i) * ( (1/2)**i ) * ( (1/2)**(n-i) )

return running_prob
```



```
In [21]: def numTwentySidedOnes(n):  
         # n = num trials  
         running_prob = 0  
  
         for k in range(math.ceil((0.09*n)), n+1):  
  
             running_prob += math.comb(n, k) * ( (1/20)**k ) * ( (19/20)**(n-k) )  
  
         return running_prob
```

```
In [22]: print("What is the exact probability of rolling the number 1 (20-sided dice)  
print()  
print("  
print()  
print("  
print()  
print("  
print("
```

What is the exact probability of rolling the number 1 (20-sided dice) more than 9% of the time in,

in 10 rolls: 40.126306076162095%

in 100 rolls: 6.308959062744848%

in 1000 rolls: 9.743965305003579e-06%

## Q3, part 4

```
In [23]: img = Image.open('q3_part4.png')  
display(img)
```

4)

i)

$$n = 10, 100, 1000$$

$$p = \frac{1}{2}$$

$$a = 0.7n + 1$$

$$b = n$$

ii)

$$n = 10, 100, 1000$$

$$p = \frac{1}{6}$$

$$a = 0.2n + 1$$

$$b = n$$

iii)

$$n = 10, 100, 1000$$

$$p = \frac{1}{20}$$

$$a = 0.09n + 1$$

$$b = n$$



```
In [24]: def binomialApproximationWContinuity(n, p, a, b):
```

```
    mu = n*p
    sigma = math.sqrt(n*p*(1-p))

    alpha = (a - 0.5 - mu) / sigma

    beta = (b + 0.5 - mu) / sigma

    return norm.cdf(beta) - norm.cdf(alpha)
```

```
def binomialApproximationNoContinuity(n, p, a, b):
```

```
    mu = n*p
    sigma = math.sqrt(n*p*(1-p))

    alpha = (a - mu) / sigma

    beta = (b - mu) / sigma

    return norm.cdf(beta) - norm.cdf(alpha)
```

## Q3, part 4-1

```
In [25]: rolls = [10,100,1000]
```

```
print("For 1. What is the exact probability of rolling a 6-sided die an even")
print()
```

```
for roll in rolls:
```

```
    print("Normal approximation for " + str(roll) + " rolls w/ continuity co")
    print("                                " +
```

```
print()
```

```
for roll in rolls:
```

```
    print("Normal approximation for " + str(roll) + " rolls w/ no continuity")
    print("                                " +
```

For 1. What is the exact probability of rolling a 6-sided die an even number more than 70% of the time in 10 rolls, 100 rolls, 1000 rolls.

Normal approximation for 10 rolls w/ continuity correction:

5.667103988860

456%

Normal approximation for 100 rolls w/ continuity correction:

0.002065750691

2491463%

Normal approximation for 1000 rolls w/ continuity correction:

0.0%

Normal approximation for 10 rolls w/ no continuity correction:

2.810708443279

7413%

Normal approximation for 100 rolls w/ no continuity correction:

0.001334574901

5902797%

Normal approximation for 1000 rolls w/ no continuity correction:

0.0%

## Q3, part 4-2

```
In [26]: rolls = [10,100,1000]

print("For 2. What is the exact probability of rolling the number 1 more than")
print()

for roll in rolls:
    print("Normal approximation for " + str(roll) + " rolls w/ continuity co")
    print(" ")

print()

for roll in rolls:
    print("Normal approximation for " + str(roll) + " rolls w/ no continuity")
    print(" ")
```

For 2. What is the exact probability of rolling the number 1 more than 20% of the time in 10 rolls, 100 rolls, 1000 rolls of a 6-sided die.

Normal approximation for 10 rolls w/ continuity correction:

23.97500610934

4356%

Normal approximation for 100 rolls w/ continuity correction:

15.18358910818

507%

Normal approximation for 1000 rolls w/ continuity correction:

0.204682577804

3937%

Normal approximation for 10 rolls w/ no continuity correction:

12.89495176454

0096%

Normal approximation for 100 rolls w/ no continuity correction:

12.24643891180

1505%

Normal approximation for 1000 rolls w/ no continuity correction:

0.178826910515

20627%

### Q3, part 4-3

```
In [27]: rolls = [10,100,1000]

print("For 3. What is the exact probability of rolling the number 1 more than")
print()

for roll in rolls:
    print("Normal approximation for " + str(roll) + " rolls w/ continuity correction")
    print(" ")
print()

for roll in rolls:
    print("Normal approximation for " + str(roll) + " rolls w/ no continuity correction")
    print(" ")
```

For 3. What is the exact probability of rolling the number 1 more than 9% of the time in 10 rolls, 100 rolls, 1000 rolls of a 20-sided die.

Normal approximation for 10 rolls w/ continuity correction:

55.76821657313

194%

Normal approximation for 100 rolls w/ continuity correction:

5.414682794950

454%

Normal approximation for 1000 rolls w/ continuity correction:

4.984296420040

835e-07%

Normal approximation for 10 rolls w/ no continuity correction:

28.08288575065

212%

Normal approximation for 100 rolls w/ no continuity correction:

3.322871000846

561%

Normal approximation for 1000 rolls w/ no continuity correction:

3.241236501416

722e-07%

## Question 4

```

In [28]: def monteCarlo(f, boxX, boxY, area, numberSamples=10000, showPlot=True):
    samplesIn=[]
    samplesOut=[]

    for k in range(numberSamples):
        x = boxX()
        y = boxY()

        if f(x) > y: # Check to see if point is below curve y=f(x)
            samplesIn.append([x,y]) # point below
        else:
            samplesOut.append([x,y]) # point above

    numIn=len(samplesIn) # number of points below
    numOut=len(samplesOut) # number of points above

    ratioIn = numIn/numberSamples # = P(R)
    totalArea = area()

    areaRegion = ratioIn * totalArea

    if (showPlot):
        print("Out of {:,} samples, {:,} are in the blue region under \n \t")
        print("Hence fraction of the samples below the curve is {:,} \n \t a")
        print()
        print("Hence our esitmate of the area under the curve is {:}.".format(ratioIn))

        x_samplesIn=[ p[0] for p in samplesIn]
        y_samplesIn=[ p[1] for p in samplesIn]

        x_samplesOut=[ p[0] for p in samplesOut]
        y_samplesOut=[ p[1] for p in samplesOut]

        plt.scatter(x_samplesOut,y_samplesOut,color='black',s=3)
        plt.scatter(x_samplesIn,y_samplesIn,color='lightblue',s=3)
        plt.xlabel("x")
        plt.ylabel("y")
        plt.show()

    #monteCarlo estimate of region for num samples

    return areaRegion

```

## Q4, part 1

My working for this question is below, and the calculations are below

```

In [29]: img = Image.open('q4_part1-1.png')
display(img)
img = Image.open('q4_part1-2.png')
display(img)
img = Image.open('q4_part1-3.png')
display(img)

```



1)  $\hat{p}_n$  = prob after  $n$  independent samples of point from  $R$  falling in  $A$ .  $p$  = true probability

$$A = \{ (x, y) \in R : \cos(x) \sin(2x) + 1 \leq y \}$$

How big does  $n$  need to be to

$$\text{ensure } P(|\hat{p}_n - p| > 0.05) < 0.01$$

$$P(|\hat{p}_n - p| > 0.05) = P(-0.05 < \hat{p}_n - p < 0.05)$$

$$\text{let } \delta = 0.05$$

$$\text{lower bound: } -\delta = \hat{p} - p = \frac{a}{n} - \frac{np}{n} = \frac{a - np}{n}$$

$$\Rightarrow a = np - n\delta, \quad b = np + n\delta$$

$np$  = expected successes

$n\hat{p}_n$  = observed successes

$a, b$  represent bounds  $\rightarrow$  on  $n$ .


such that  $|\hat{p}_n - p| > 0.05$

is true w/  $< 0.01$  probability,

$\delta = \frac{a - np}{n}$  transforms for use on standard normal.

$$P(|\hat{p}_n - p| > 0.05) = P(\hat{p}_n - p > 0.05 \text{ or } \hat{p}_n - p < -0.05) < 0.01$$

$$P(\hat{p}_n - p > 0.05) + P(\hat{p}_n - p < -0.05) < 0.01$$



$$\Rightarrow 2 \cdot P(\hat{p}_n - p < -0.05) < 0.01$$

$$z = \frac{x - \mu}{\sigma} = \frac{n\hat{p}_n - np}{\sqrt{npq}} = \frac{n(\hat{p}_n - p)}{\sqrt{npq}}$$

$$= \frac{\sqrt{n}(\hat{p}_n - p)}{\sqrt{pq}} = \frac{\delta\sqrt{n}}{\sqrt{pq}}$$

$$= 2 \cdot \Phi\left(\frac{\delta\sqrt{n}}{\sqrt{pq}}\right) - 0.01 < 0$$

$$\Rightarrow \Phi\left(\frac{\delta\sqrt{n}}{\sqrt{pq}}\right) = 0.005 \quad \delta = -0.05$$

$$\frac{\delta\sqrt{n}}{\sqrt{pq}} = \Phi^{-1}(0.005)$$

$$\Rightarrow \sqrt{n} = \frac{\sqrt{pq}}{\delta} \Phi^{-1}(0.005)$$

$$\Rightarrow n = \frac{pq}{\delta^2} (\Phi^{-1}(0.005))^2$$

As we assume we do not

know  $I$ , we cannot know true  $p$

$\therefore$  Must take worst case  $p = \frac{1}{2}$

from code: 663.49

$$\Rightarrow n \geq \underline{\underline{664}}$$

Q4, part 2

```
In [30]: #assume worst case scenario
p = 1/2
q = 1-p

delta = -0.05

confidence = 0.01 ## this is the confidence for which we are trying to get n

n = ((p*q) / ((delta)** 2)) * (norm.ppf(confidence/2)**2)

print(n)

663.4896601021214
```

```
In [31]: #function to integrate
def function(x):
    return math.cos( x ) * math.sin( 2*x) + 1

#random x plot value is between 0 and 8
def boxX():
    return (8)*rnd.random()

# random y plot value is between 0 and 2
def boxY():
    return (2)*rnd.random()


# box area = 2*8
def trueArea():
    return 2*8
```

```
In [32]: monteCarlo(function, boxX, boxY, trueArea)
```

Out of 10,000 samples, 5,421 are in the blue region under the curve and 4,579 are above.

Hence fraction of the samples below the curve is 0.5421 and the fraction above is 0.4579.

Hence our estimate of the area under the curve is 8.6736.



Out[32]: 8.6736

```
In [33]: img = Image.open('q4_part2-1.png')
display(img)
img = Image.open('q4_part2-2.png')
display(img)
```

$$2) \quad P(|\hat{I} - I| \leq \delta) > 0.9$$

$$P(-\delta \leq \hat{I} - I \leq \delta) > 0.9$$



$$= 1 - \text{[unshaded area in the tails]}$$

$$= 1 - 2 \times \text{[unshaded area in the tails]}$$

$$\rightarrow 1 - 2P(\hat{I} - I \leq \delta) > 0.9$$

$$\Rightarrow 2P(\hat{I} - I \leq \delta) < 0.1$$

$$\Rightarrow P(\hat{I} - I \leq \delta) < 0.05$$


$$\bar{\Phi}(z) < 0.05$$

$$z = \frac{x - \mu}{\sigma} \quad x = n\hat{p}_n$$

$$\hat{I} = \left( \frac{n\hat{p}_n}{n} \right) \text{boxSize} = \hat{p}_n \cdot \text{boxSize} = 16\hat{p}_n$$

$$I = \frac{n p}{n} \text{boxSize} = p \cdot \text{boxSize} = 16 p.$$

$$z = \frac{x - \mu}{\sigma}$$



$$\begin{aligned} \rightarrow \Phi(z) = 0.05 &\Rightarrow z = \Phi^{-1}(0.05) = -1.6449 \quad \checkmark \\ \therefore \frac{\bar{x} - \mu}{\sigma} &= \frac{n\hat{p}_n - np}{\sqrt{npq}} = \frac{\sqrt{n} \left( \frac{\hat{I} - I}{\text{boxSize}} \right)}{\sqrt{pq}} \\ \Rightarrow \hat{I} - I &= \frac{-1.645 \cdot \sqrt{pq} \cdot \text{boxSize}}{\sqrt{n}} = -0.1861 > \delta \\ \therefore \delta &> \underline{\underline{0.1861}} \end{aligned}$$

In [34]: `((-1.645) * (1/2) * 16)/math.sqrt(5000)`

Out[34]: -0.1861105048082993

Q4, part 3

```

In [35]: numSamp = 5000
delta = 0.18611

history = []
average_history = []
error = []
running_probability = []
average_running_probability = []
num_success = 0

def function(x):
    return math.cos( x ) * math.sin( 2*x ) + 1

#random x plot value is between 0 and 8
def boxX():
    return (8)*rnd.random()

# random y plot value is between 0 and 2
def boxY():
    return (2)*rnd.random()

# box area = 2*8
def trueArea():
    return 2*8

true_integral = 8.6687
print("True integral is ", true_integral)

for m in range(1, 1001):
    history.append(monteCarlo(function, boxX, boxY, trueArea, numberSamples=
    if abs(history[-1] - true_integral) <= delta:
        num_success += 1
    running_probability.append(num_success/m)
    average_running_probability.append(sum(running_probability)/len(running_

print(f"The probability that the Monte Carlo approximation was within {delta


plt.plot(running_probability)
plt.xlabel('Number of Samples')
plt.ylabel('Probability of Success')
plt.title('Probability of Success with Monte Carlo Integration')
plt.show()

```

True integral is 8.6687

The probability that the Monte Carlo approximation was within 0.18611 of the true integral after 1000 samples is 0.8936





## Q4, part4

```
In [36]: img = Image.open('q4_part4.png')
display(img)
```

$$4) P(|\hat{p} - p| \leq 0.25) > 0.9$$

$$1 - 2\Phi(z) > 0.9$$

$$z = \frac{\hat{p} - p}{\sigma} = \frac{n\hat{p} - np}{\sigma}$$

$$= \frac{\sqrt{n} \cdot \frac{1}{4}}{\sqrt{p_0 \cdot \text{boxSize}}}$$

$$\therefore \sqrt{n} > \frac{\text{boxSize} \sqrt{p_0} \Phi^{-1}(0.95)^2}{\frac{1}{4}}$$

$$\therefore n > \underline{\underline{2771}}$$

```
In [34]: img = Image.open('q4_part5.png')
display(img)
```

5)  $\int_0^8 \cos x \sin 2x + 1 \, dx \approx \underline{\underline{8.6687}}$

Very similar to the approximate answers I got

In [ ]: