

Bases

Brian D. Fitzpatrick

February 23, 2022



Overview

Bases

Definition

Examples

Fundamental Subspaces

Bases

Examples

Digraphs

Bases of $\text{Null}(A)$

Bases of $\text{Null}(A^T)$

Bases

Definition

Definition

A list of vectors $\beta = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_d\}$ in a vector space V is a *basis* if

Bases

Definition

Definition

A list of vectors $\beta = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_d\}$ in a vector space V is a *basis* if

Spanning Axiom

$$\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_d\} = V$$

Bases

Definition

Definition

A list of vectors $\beta = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_d\}$ in a vector space V is a *basis* if

Spanning Axiom

$$\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_d\} = V$$

Linear Independence Axiom

$$\{\mathbf{v}_1, \dots, \mathbf{v}_d\} \text{ linearly independent}$$

Bases

Definition

Definition

A list of vectors $\beta = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_d\}$ in a vector space V is a *basis* if

Spanning Axiom

$$\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_d\} = V$$

Linear Independence Axiom

$$\{\mathbf{v}_1, \dots, \mathbf{v}_d\} \text{ linearly independent}$$

We think of a basis as a *minimal spanning set*.

Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$

Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$



Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$




Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$




Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$




Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$



These vectors are all *dependent*.

Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$


These vectors are all *dependent*. Choosing any one gives a basis of V .


$$V = \text{Span}\{[1 \ 2 \ 3]^T\} \quad V = \text{Span}\{[2 \ 4 \ 6]^T\} \quad V = \text{Span}\{[-1 \ -2 \ -3]^T\}$$

Bases

Examples

Example

Consider the vector space $V \subset \mathbb{R}^3$ given by

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ -3 \end{bmatrix} \right\}$$


These vectors are all *dependent*. Choosing any one gives a basis of V .

$$V = \text{Span}\{[1 \ 2 \ 3]^T\} \quad V = \text{Span}\{[2 \ 4 \ 6]^T\} \quad V = \text{Span}\{[-1 \ -2 \ -3]^T\}$$

Vector spaces have infinitely many bases!

Bases

Examples

Example

To find vectors $\mathbf{v} \in \text{Null}(A)$, we must solve $A\mathbf{v} = \mathbf{0}$ for \mathbf{v} .

$$\text{rref} \begin{bmatrix} 2 & 1 & 5 & 25 \\ -1 & -12 & 9 & 68 \\ 3 & 5 & 4 & 13 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 3 & 16 \\ 0 & 1 & -1 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Bases

Examples

Example

To find vectors $\mathbf{v} \in \text{Null}(A)$, we must solve $A\mathbf{v} = \mathbf{0}$ for \mathbf{v} .

$$\text{rref} \begin{matrix} A \\ \begin{bmatrix} 2 & 1 & 5 & 25 \\ -1 & -12 & 9 & 68 \\ 3 & 5 & 4 & 13 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 3 & 16 \\ 0 & 1 & -1 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Bases

Examples

Example

To find vectors $\mathbf{v} \in \text{Null}(A)$, we must solve $A\mathbf{v} = \mathbf{0}$ for \mathbf{v} .

$$\text{rref} \begin{matrix} A \\ \begin{bmatrix} 2 & 1 & 5 & 25 \\ -1 & -12 & 9 & 68 \\ 3 & 5 & 4 & 13 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 3 & 16 \\ 0 & 1 & -1 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} -3c_1 - 16c_2 \\ c_1 + 7c_2 \\ c_1 \\ c_2 \end{bmatrix} = c_1 \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \\ 0 \end{bmatrix} + c_2 \cdot \begin{bmatrix} -16 \\ 7 \\ 0 \\ 1 \end{bmatrix}$$

Bases

Examples

Example

To find vectors $\mathbf{v} \in \text{Null}(A)$, we must solve $A\mathbf{v} = \mathbf{0}$ for \mathbf{v} .

$$\text{rref} \begin{matrix} A \\ \downarrow \\ \begin{bmatrix} 2 & -1 & 5 & 25 \\ -1 & -12 & 9 & 68 \\ 3 & 5 & 4 & 13 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 3 & 16 \\ 0 & 1 & -1 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} -3c_1 - 16c_2 \\ c_1 + 7c_2 \\ c_1 \\ c_2 \end{bmatrix} = c_1 \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \\ 0 \end{bmatrix} + c_2 \cdot \begin{bmatrix} -16 \\ 7 \\ 0 \\ 1 \end{bmatrix}$$

Here, we have shown that

$$\text{Null}(A) = \text{Span}\left\{ \begin{bmatrix} -3 & 1 & 1 & 0 \end{bmatrix}^T, \begin{bmatrix} -16 & 7 & 0 & 1 \end{bmatrix}^T \right\}$$

Bases

Examples

Example

To find vectors $\mathbf{v} \in \text{Null}(A)$, we must solve $A\mathbf{v} = \mathbf{0}$ for \mathbf{v} .

$$\text{rref} \begin{matrix} A \\ \downarrow \end{matrix} \begin{bmatrix} 2 & 1 & 5 & 25 \\ -1 & -12 & 9 & 68 \\ 3 & 5 & 4 & 13 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 3 & 16 \\ 0 & 1 & -1 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} -3c_1 - 16c_2 \\ c_1 + 7c_2 \\ c_1 \\ c_2 \end{bmatrix} = c_1 \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \\ 0 \end{bmatrix} + c_2 \cdot \begin{bmatrix} -16 \\ 7 \\ 0 \\ 1 \end{bmatrix}$$

Here, we have shown that

$$\text{Null}(A) = \text{Span}\left\{ \begin{bmatrix} -3 & 1 & 1 & 0 \end{bmatrix}^T, \begin{bmatrix} -16 & 7 & 0 & 1 \end{bmatrix}^T \right\}$$

Bases

Examples

Example

To find vectors $\mathbf{v} \in \text{Null}(A)$, we must solve $A\mathbf{v} = \mathbf{0}$ for \mathbf{v} .

$$\text{rref} \begin{matrix} A \\ \downarrow \end{matrix} \begin{bmatrix} 2 & 1 & 5 & 25 \\ -1 & -12 & 9 & 68 \\ 3 & 5 & 4 & 13 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 3 & 16 \\ 0 & 1 & -1 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} -3c_1 - 16c_2 \\ c_1 + 7c_2 \\ c_1 \\ c_2 \end{bmatrix} = c_1 \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \\ 0 \end{bmatrix} + c_2 \cdot \begin{bmatrix} -16 \\ 7 \\ 0 \\ 1 \end{bmatrix}$$

"pivot solutions" to $A\mathbf{v} = \mathbf{0}$ \uparrow \uparrow

Here, we have shown that

$$\text{Null}(A) = \text{Span}\left\{ \begin{bmatrix} -3 & 1 & 1 & 0 \end{bmatrix}^T, \begin{bmatrix} -16 & 7 & 0 & 1 \end{bmatrix}^T \right\}$$

Bases

Examples

Example

To find vectors $\mathbf{v} \in \text{Null}(A)$, we must solve $A\mathbf{v} = \mathbf{0}$ for \mathbf{v} .

$$\text{rref} \begin{matrix} A \\ \downarrow \end{matrix} \begin{bmatrix} 2 & 1 & 5 & 25 \\ -1 & -12 & 9 & 68 \\ 3 & 5 & 4 & 13 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 3 & 16 \\ 0 & 1 & -1 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} -3c_1 - 16c_2 \\ c_1 + 7c_2 \\ c_1 \\ c_2 \end{bmatrix} = c_1 \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \\ 0 \end{bmatrix} + c_2 \cdot \begin{bmatrix} -16 \\ 7 \\ 0 \\ 1 \end{bmatrix}$$

"pivot solutions" to $A\mathbf{v} = \mathbf{0}$ \uparrow \uparrow

Here, we have shown that

$$\text{Null}(A) = \text{Span}\left\{ \begin{bmatrix} -3 & 1 & 1 & 0 \end{bmatrix}^T, \begin{bmatrix} -16 & 7 & 0 & 1 \end{bmatrix}^T \right\}$$


This is called the *pivot basis* of $\text{Null}(A)$.

Fundamental Subspaces

Bases

Theorem

We may systematically construct bases of the four fundamental subspaces.




Fundamental Subspaces

Bases

Theorem


We may systematically construct bases of the four fundamental subspaces.

$$\begin{aligned}\text{Col}(A^T) &= \text{Span}\{\text{pivot columns of } A^T\} \\ &= \text{Span}\{\text{nonzero rows of } \text{rref}(A)\}\end{aligned}$$



$$\mathbb{R}^n \xrightarrow{A} \mathbb{R}^m$$

$$\begin{aligned}\text{Col}(A) &= \text{Span}\{\text{pivot columns of } A\} \\ &= \text{Span}\{\text{nonzero rows of } \text{rref}(A^T)\}\end{aligned}$$




Fundamental Subspaces

Bases

Theorem

We may systematically construct bases of the four fundamental subspaces.




Fundamental Subspaces

Bases

Theorem

We may systematically construct bases of the four fundamental subspaces.



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} 2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{bmatrix} 2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \xrightarrow{A} \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \xrightarrow{A^T} \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces

Examples


Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.

$\beta_1 =$



Fundamental Subspaces

Examples


Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} 2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.

$\beta_1 =$



Fundamental Subspaces

Examples

Example

Consider the calculations


$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} 2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.

$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} 2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$


This information gives bases of the four fundamental subspaces.

$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_2 =$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{matrix} \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} 2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{matrix} \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{matrix}$$


This information gives bases of the four fundamental subspaces.

$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_2 =$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{matrix} \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \end{matrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} 2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{matrix} \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{matrix}$$


This information gives bases of the four fundamental subspaces.

$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_2 = \{[1 \ 0 \ 2 \ 1]^T, [0 \ 1 \ 0 \ 1]^T\}$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.


$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_2 = \{[1 \ 0 \ 2 \ 1]^T, [0 \ 1 \ 0 \ 1]^T\}$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$

$\beta =$



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.


$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_2 = \{[1 \ 0 \ 2 \ 1]^T, [0 \ 1 \ 0 \ 1]^T\}$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$

$\beta =$



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.


$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_2 = \{[1 \ 0 \ 2 \ 1]^T, [0 \ 1 \ 0 \ 1]^T\}$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$

$$\beta = \{[-2 \ 0 \ 1 \ 0]^T, [-1 \ -1 \ 0 \ 1]^T\}$$



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \overset{A \rightarrow}{\begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix}} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \overset{A^T \rightarrow}{\begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix}} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.



Fundamental Subspaces

Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.


$$\beta_1 = \{[-2 \ -7 \ -4 \ -9]^T, [3 \ 10 \ 6 \ 13]^T\}$$

$$\beta_2 = \{[1 \ 0 \ 2 \ 1]^T, [0 \ 1 \ 0 \ 1]^T\}$$

$$\beta = \{[-2 \ 0 \ 1 \ 0]^T, [-1 \ -1 \ 0 \ 1]^T\}$$

$$\beta_1 = \{[-2 \ 3 \ -1]^T, [-7 \ 10 \ -2]^T\}$$

$$\beta_2 = \{[1 \ 0 \ -4]^T, [0 \ 1 \ -3]^T\}$$



Fundamental Subspaces


Examples

Example

Consider the calculations

$$\text{rref} \begin{matrix} A \rightarrow \\ \begin{bmatrix} -2 & -7 & -4 & -9 \\ 3 & 10 & 6 & 13 \\ -1 & -2 & -2 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{matrix} A^T \rightarrow \\ \begin{bmatrix} -2 & 3 & -1 \\ -7 & 10 & -2 \\ -4 & 6 & -2 \\ -9 & 13 & -3 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

This information gives bases of the four fundamental subspaces.




Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.




Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.




Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.




Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.




Let $\{c_1, \dots, c_k\}$ be cycles, each passing through **exactly one** purged vector.

Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.



Let $\{c_1, \dots, c_k\}$ be cycles, each passing through **exactly one** purged vector.


$$\mathbf{v}_{c_1} = \begin{bmatrix} a_1 & \textcolor{red}{a_2} & a_3 & a_4 & a_5 \\ -1 & \textcolor{red}{1} & 1 & 0 & 0 \end{bmatrix}$$

Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.



Let $\{c_1, \dots, c_k\}$ be cycles, each passing through **exactly one** purged vector.


$$\mathbf{v}_{c_1} = \begin{bmatrix} a_1 & \textcolor{red}{a_2} & a_3 & a_4 & a_5 \\ -1 & \textcolor{red}{1} & 1 & 0 & 0 \end{bmatrix}$$

Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.



Let $\{c_1, \dots, c_k\}$ be cycles, each passing through **exactly one** purged vector.


$$\mathbf{v}_{c_1} = \begin{bmatrix} a_1 & \textcolor{red}{a_2} & a_3 & a_4 & a_5 \\ -1 & \textcolor{red}{1} & 1 & 0 & 0 \end{bmatrix}$$

Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.



Let $\{c_1, \dots, c_k\}$ be cycles, each passing through *exactly one* purged vector.


$$\mathbf{v}_{c_1} = \begin{bmatrix} a_1 & \textcolor{red}{a_2} & a_3 & a_4 & a_5 \\ -1 & \textcolor{red}{1} & 1 & 0 & 0 \end{bmatrix} \quad \mathbf{v}_{c_2} = \begin{bmatrix} a_1 & a_2 & a_3 & \textcolor{red}{a_4} & a_5 \\ 1 & 0 & 0 & \textcolor{red}{1} & 0 \end{bmatrix}$$

Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.



Let $\{c_1, \dots, c_k\}$ be cycles, each passing through *exactly one* purged vector.


$$\mathbf{v}_{c_1} = \begin{bmatrix} a_1 & \textcolor{red}{a_2} & a_3 & a_4 & a_5 \\ -1 & \textcolor{red}{1} & 1 & 0 & 0 \end{bmatrix} \quad \mathbf{v}_{c_2} = \begin{bmatrix} a_1 & a_2 & a_3 & \textcolor{red}{a_4} & a_5 \\ 1 & 0 & 0 & \textcolor{red}{1} & 0 \end{bmatrix}$$

Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.



Let $\{c_1, \dots, c_k\}$ be cycles, each passing through *exactly one* purged vector.

$$\mathbf{v}_{c_1} = \begin{bmatrix} a_1 & \textcolor{red}{a_2} & a_3 & a_4 & a_5 \\ -1 & \textcolor{red}{1} & 1 & 0 & 0 \end{bmatrix} \quad \mathbf{v}_{c_2} = \begin{bmatrix} a_1 & a_2 & a_3 & \textcolor{red}{a_4} & a_5 \\ 1 & 0 & 0 & \textcolor{red}{1} & 0 \end{bmatrix}$$


Then $\{\mathbf{v}_{c_1}, \dots, \mathbf{v}_{c_k}\}$ is a basis of $\text{Null}(A)$.

Digraphs

Bases of $\text{Null}(A)$

Theorem

Purge the minimum number of arrows necessary to break all cycles.



Let $\{c_1, \dots, c_k\}$ be cycles, each passing through *exactly one* purged vector.

$$\mathbf{v}_{c_1} = \begin{bmatrix} a_1 & \textcolor{red}{a_2} & a_3 & a_4 & a_5 \\ -1 & \textcolor{red}{1} & 1 & 0 & 0 \end{bmatrix} \quad \mathbf{v}_{c_2} = \begin{bmatrix} a_1 & a_2 & a_3 & \textcolor{red}{a_4} & a_5 \\ 1 & 0 & 0 & \textcolor{red}{1} & 0 \end{bmatrix}$$


Then $\{\mathbf{v}_{c_1}, \dots, \mathbf{v}_{c_k}\}$ is a basis of $\text{Null}(A)$. Here, $\text{Null}(A) = \text{Span}\{\mathbf{v}_{c_1}, \mathbf{v}_{c_2}\}$.

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .




Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .




Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{\mathbf{v}_{G_1}, \dots, \mathbf{v}_{G_k}\}$.


$$\mathbf{v}_{G_1} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 1 & 1 & 1 & 0 & 0 \end{bmatrix}^T$$

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{\mathbf{v}_{G_1}, \dots, \mathbf{v}_{G_k}\}$.


$$\mathbf{v}_{G_1} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 1 & 1 & 1 & 0 & 0 \end{bmatrix}^T$$

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{v_{G_1}, \dots, v_{G_k}\}$.


$$v_{G_1} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 1 & 1 & 1 & 0 & 0 \end{bmatrix}^T$$

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{\mathbf{v}_{G_1}, \dots, \mathbf{v}_{G_k}\}$.


$$\mathbf{v}_{G_1} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 1 & 1 & 1 & 0 & 0 \end{bmatrix}^T \quad \mathbf{v}_{G_2} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}^T$$

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{\mathbf{v}_{G_1}, \dots, \mathbf{v}_{G_k}\}$.


$$\mathbf{v}_{G_1} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 1 & 1 & 1 & 0 & 0 \end{bmatrix}^T \quad \mathbf{v}_{G_2} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}^T$$

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{\mathbf{v}_{G_1}, \dots, \mathbf{v}_{G_k}\}$.


$$\mathbf{v}_{G_1} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 1 & 1 & 1 & 0 & 0 \end{bmatrix}^T \quad \mathbf{v}_{G_2} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}^T$$

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{\mathbf{v}_{G_1}, \dots, \mathbf{v}_{G_k}\}$.

$$\mathbf{v}_{G_1} = \begin{matrix} & v_1 & v_2 & v_3 & v_4 & v_5 \\ \begin{matrix} v_1 & v_2 & v_3 & v_4 & v_5 \end{matrix}^T & \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad \mathbf{v}_{G_2} = \begin{matrix} & v_1 & v_2 & v_3 & v_4 & v_5 \\ \begin{matrix} v_1 & v_2 & v_3 & v_4 & v_5 \end{matrix}^T & \begin{bmatrix} 0 & 0 & 0 & 1 & 1 \end{bmatrix} \end{matrix}$$


These vectors form a basis of $\text{Null}(A^T)$.

Digraphs

Bases of $\text{Null}(A^T)$

Theorem

Let $\{G_1, \dots, G_k\}$ be the connected components of a digraph G .



Consider the associated *classification vectors* $\{\mathbf{v}_{G_1}, \dots, \mathbf{v}_{G_k}\}$.

$$\mathbf{v}_{G_1} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 1 & 1 & 1 & 0 & 0 \end{bmatrix}^T \quad \mathbf{v}_{G_2} = \begin{bmatrix} v_1 & v_2 & v_3 & v_4 & v_5 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}^T$$

These vectors form a basis of $\text{Null}(A^T)$. Here, $\text{Null}(A^T) = \text{Span}\{\mathbf{v}_{G_1}, \mathbf{v}_{G_2}\}$.