

01:640:495 - Lecture 10

Pranav Tikkawar

March 6, 2025

1. Following are two figures from the article “A singularly Valuable Decomposition: The SVD of a Matrix” by Dan Kalman. Can you explain the idea the figures are communicating?

Solution: The first figure shows orthogonal projection of a basis of \mathbb{R}^3 onto a plane. Then it shows the operation of multiplying the projected basis to another basis. The second figure shows the null space of \mathbf{A} mapping to the range of \mathbf{A}^T and the null space of \mathbf{A}^T mapping to the range of \mathbf{A} .

2. Suppose $\mathbf{v}_1, \mathbf{v}_2$ are eigenvectors of $\mathbf{A}^T \mathbf{A}$ that are orthogonal $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$. Show their images under \mathbf{A} are orthogonal - that is $\mathbf{A}\mathbf{v}_1 \cdot \mathbf{A}\mathbf{v}_2 = 0$.

Solution: We can see that

$$\begin{aligned}\langle \mathbf{A}v_1, \mathbf{A}v_2 \rangle &= (\mathbf{A}v_1)^T (\mathbf{A}v_2) \\ &= v_1^T \mathbf{A}^T \mathbf{A} v_2 \\ &= v_1^T \lambda_1 v_2 \\ &= \lambda_1 v_1^T v_2 \\ &= 0\end{aligned}$$

3. Let $\mathbf{A} = \begin{bmatrix} 1 & -1 \\ -1 & 2 \\ 2 & -1 \end{bmatrix}$ For your convenience, you are told that $\mathbf{A}^T \mathbf{A} = \begin{bmatrix} 6 & -5 \\ -5 & 6 \end{bmatrix}$ has eigenvalues $\lambda = 11, 1$ with eigenvectors $\begin{bmatrix} -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ respectively a basis for the two eigenspaces (null space (kernel) of $\mathbf{A}^T \mathbf{A} - \lambda \mathbf{I}$). Also, $\mathbf{A} \mathbf{A}^T = \begin{bmatrix} 2 & -3 & 3 \\ -3 & 5 & -4 \\ 3 & -4 & 5 \end{bmatrix}$ has

eigenvalues $\lambda = 1, 0, 11$ with eigenvectors $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$, $\begin{bmatrix} -3 \\ -1 \\ 1 \end{bmatrix}$, $\begin{bmatrix} \frac{2}{3} \\ -1 \\ 1 \end{bmatrix}$ respectively a basis for the three eigenspaces (null space (kernel) of $\mathbf{A}\mathbf{A}^T - \lambda\mathbf{I}$).

- (a) Obtain a singular value decomposition of \mathbf{A} .

Solution: For our SVD we have

$$\Sigma = \begin{bmatrix} \sqrt{11} & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$V = \begin{bmatrix} \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$U = \begin{bmatrix} \frac{2}{\sqrt{22}} & 0 & \frac{-3}{\sqrt{10}} \\ \frac{-3}{\sqrt{22}} & \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{10}} \\ \frac{\sqrt{22}}{3} & \frac{\sqrt{2}}{1} & \frac{\sqrt{10}}{1} \\ \frac{\sqrt{22}}{3} & \frac{\sqrt{2}}{1} & \frac{\sqrt{10}}{1} \end{bmatrix}$$

- (b) Use it to write \mathbf{A} as sum of rank 1 projection matrices.

Solution: We can write out the SVD as

$$11 \begin{bmatrix} \frac{2}{\sqrt{22}} \\ \frac{-3}{\sqrt{22}} \\ \frac{\sqrt{22}}{3} \end{bmatrix} \begin{bmatrix} \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} + 1 \begin{bmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

- (c) What is the rank of \mathbf{A} ? Is \mathbf{A} a ‘full rank’ matrix?

Solution: The rank of \mathbf{A} is 2. It is not a full rank matrix.

- (d) Determine the spectral norm of \mathbf{A} . (Recall it is the maximum $\|\mathbf{A}\mathbf{X}\|$ among \mathbf{X} with $\|\mathbf{X}\| = 1$ with $\|\cdot\|$ denoting the usual Euclidean norm (length).)

Solution: The spectral norm of \mathbf{A} is $\sqrt{11}$. It is the value of the largest singular value.

4. Matrices \mathbf{A} , \mathbf{b} are given matrices of sizes $m \times n$, $m \times 1$ respectively. And, $\mathbf{x} \in \mathbb{R}^n$ variable. Consider the function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by $f(\mathbf{x}) = \|\mathbf{A}\mathbf{x} - \mathbf{b}\|_2$.

- (a) Express $f(\mathbf{x})$ using matrix operations.

Solution:

$$\begin{aligned}
 f(x) &= (Ax - b)^T(Ax - b) \\
 &= x^T A^T Ax + b^T b - b^T Ax - x^T A^T b \\
 &= \langle Ax, Ax \rangle + \langle b, b \rangle - \langle b, Ax \rangle - \langle Ax, b \rangle \\
 &= \langle Ax, Ax \rangle + \langle b, b \rangle - 2\langle b, Ax \rangle \\
 &= \|Ax\|^2 + \|b\|^2 - 2\langle b, Ax \rangle
 \end{aligned}$$

This is essentially splitting up the norm into its components.

- (b) Assuming the derivative exists, compute the derivative (gradient) linear map $\frac{df}{dx}(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$. You may use $\mathbf{h} \in \mathbb{R}^n$ as the argument of this linear map (so, for the “direction”). Examining the expression of this linear map, also determine the $1 \times n$ matrix of this linear map $\frac{df}{dx}(\mathbf{x})$. *Hint: Differentiate a curve passing through \mathbf{x} in the direction \mathbf{h} i.e., use $\frac{df}{dx}(\mathbf{x})(\mathbf{h}) = \frac{d}{dt} \bigg|_{t=0} f(\mathbf{x} + t\mathbf{h})$ and previous part.*

Solution: Using $x + th$ as x in the previous part, we get

$$\begin{aligned}
 f(x + th) &= \|A(x + th)\|^2 + \|b\|^2 - 2\langle b, A(x + th) \rangle \\
 &= x^T A^T Ax + tx^T A^T Ah + th^T A^T Ax + t^2 h^T A^T Ah - 2b^T Ax - 2tb^T Ah + b^T b
 \end{aligned}$$

Taking the derivative of this with respect to t and evaluating at $t = 0$ gives us the derivative.

$$\begin{aligned}
 \frac{d}{dt} \bigg|_{t=0} f(x + th) &= \frac{d}{dt} \bigg|_{t=0} (\|A(x + th)\|^2 + \|b\|^2 - 2\langle b, A(x + th) \rangle) \\
 &= \frac{d}{dt} \bigg|_{t=0} x^T A^T Ax + tx^T A^T Ah + th^T A^T Ax + t^2 h^T A^T Ah - 2b^T Ax - 2tb^T Ah + b^T b \\
 &= x^T A^T Ah + h^T A^T Ax - 2th^T A^T Ah - 2b^T Ah \bigg|_{t=0} \\
 &= 2x^T A^T Ah - 2b^T Ah \\
 &= 2(Ax - b)^T Ah
 \end{aligned}$$

Thus we get

$$\frac{d}{dx} f(x) = 2(Ax - b)^T A$$

- (c) Determine \mathbf{x} where the linear map $\frac{df}{dx}$ is zero. Do you recognize this scenario from earlier?

Solution: The linear map is zero then $(Ax - b)^T A = 0$. This is the normal equation for the least squares problem.

$$(Ax - b)^T A = 0$$

$$A^T Ax = A^T b$$

$$x = (A^T A)^{-1} A^T b$$