



PRACTICAL 9 LINEAR AND MULTIPLE LINEAR REGRESSION

Dr. Sudeep Tanwar

LINEAR REGRESSION

In regression, one variable is considered independent (=predictor) variable (X) and the other the dependent (=outcome) variable Y .

Widely applied:


- to predict the outputs,
- forecasting the data,
- analyzing the time series, and
- finding the causal effect dependencies between the variables

SIMPLE LINEAR REGRESSION

- Only one independent variable, X
- Relationship between X and Y is described by a linear function
- Changes in Y are assumed to be caused by changes in X

SIMPLE LINEAR REGRESSION

$$y = a + bx$$



SIMPLE LINEAR REGRESSION

a and b are given by the following formulas:

$$a \text{ (intercept)} = \frac{\sum y \sum x^2 - \sum x \sum xy}{(\sum x^2) - (\sum x)^2}$$

$$b \text{ (slope)} = \frac{n \sum xy - (\sum x)(\sum y)}{n \sum x^2 - (\sum x)^2}$$

Where,

x and y are two variables on the regression line.

b = Slope of the line.

a = y -intercept of the line.

x = Values of the first data set.

y = Values of the second data set.

Question: Find linear regression equation for the following two sets of data:

x	2	4	6	8
y	3	7	5	10

Solution:

Construct the following table:

x	y	x^2	xy
2	3	4	6
4	7	16	28
6	5	36	30
8	10	64	80
$\sum x = 20$	$\sum y = 25$	$\sum x^2 = 120$	$\sum xy = 144$

EXAMPLE

$$b = \frac{n \sum xy - (\sum x)(\sum y)}{n \sum x^2 - (\sum x)^2}$$

$$b = \frac{4 \times 144 - 20 \times 25}{4 \times 120 - 400}$$

$$b = 0.95$$

$$a = \frac{\sum y \sum x^2 - \sum x \sum xy}{n(\sum x^2) - (\sum x)^2}$$

$$a = \frac{25 \times 120 - 20 \times 144}{4(120) - 400}$$

$$a = 1.5$$


Linear regression is given by:

$$y = a + bx$$

$$y = 1.5 + 0.95 x$$

CODE

```
for i in range(10):
    x.append(random.randint(1,100));
    y.append(random.randint(1,100));
x.sort();
y.sort();
n=len(x);
E_y=sum(y);
E_x=sum(x);
E_xy=sum([i*j for i,j in zip(x,y)]);
E_x2=sum([i*i for i in x]);
E_y2=sum([i*i for i in y]);
intercept=(E_y*E_x2-E_x*E_xy)/(E_x2-E_x**2);
slope=(n*E_xy-E_x*E_y)/(n*E_x2-E_x**2);
```



```
x_val=list(range(80));
y_val=[intercept+slope*i for i in x_val];
print("\nintercept={ }, slope={ }\n".format(intercept,slope));
plt.plot(x,y);
plt.plot(x_val,y_val);
plt.xlabel("\nX-axis\n");
plt.ylabel("\nY-axis\n");
plt.title("\nLinear Regression\n");
plt.show();
```


MULTIPLE LIN REGRESSION

In linear regression, there is only one independent and dependent variable involved. But, in the case of multiple regression, there will be a set of independent variables that helps us to explain better or predict the dependent variable y .

The multiple regression equation is given by

$$Y = a + b_1X_1 + b_2X_2 + \dots + b_kX_k + e_i$$

MULTIPLE LIN REGRESSION

Multiple linear regression is used to estimate the relationship between **two or more independent variables** and **one dependent variable**.

$$b_1 = \frac{(\sum x_2^2)(\sum x_1 y) - (\sum x_1 x_2)(\sum x_2 y)}{(\sum x_1^2)(\sum x_2^2) - (\sum x_1 x_2)^2}$$

$$a = \bar{Y} - b_1 \bar{X}_1 - b_2 \bar{X}_2$$

and

$$b_2 = \frac{(\sum x_1^2)(\sum x_2 y) - (\sum x_1 x_2)(\sum x_1 y)}{(\sum x_1^2)(\sum x_2^2) - (\sum x_1 x_2)^2}$$