Module 15-1 Simple Harmonic Motion


- •1 An object undergoing simple harmonic motion takes 0.25 s to travel from one point of zero velocity to the next such point. The distance between those points is 36 cm. Calculate the (a) period, (b) frequency, and (c) amplitude of the motion.
- •2 A 0.12 kg body undergoes simple harmonic motion of amplitude 8.5 cm and period 0.20 s. (a) What is the magnitude of the maximum force acting on it? (b) If the oscillations are produced by a spring, what is the spring constant?
- •3 What is the maximum acceleration of a platform that oscillates at amplitude 2.20 cm and frequency 6.60 Hz?
- •4 An automobile can be considered to be mounted on four identical springs as far as vertical oscillations are concerned. The springs of a certain car are adjusted so that the oscillations have a frequency of 3.00 Hz. (a) What is the spring constant of each spring if the mass of the car is 1450 kg and the mass is evenly distributed over the springs? (b) What will be the oscillation frequency if five passengers, averaging 73.0 kg each, ride in the car with an even distribution of mass?

4. (a) Since the problem gives the frequency f = 3.00 Hz, we have $\omega = 2\pi f = 6\pi$ rad/s (understood to be valid to three significant figures). Each spring is considered to support 1815 kg, then Eq. 15-12 leads to

$$\omega = \sqrt{\frac{k}{m_{\text{car}}/4}} \implies k = \frac{1}{4} (1450 \text{kg}) (6\pi \text{ rad/s})^2 = 1.29 \times 10^5 \text{ N/m}.$$

$$\omega_{\text{new}} = \sqrt{\frac{k}{m_{\text{total}}/4}} \implies f_{\text{new}} = \frac{1}{2\pi} \sqrt{\frac{1.29 \times 10^5 \text{ N/m}}{(1815/4) \text{ kg}}} = 2.68 \text{ Hz}.$$

- •5 SSM In an electric shaver, the blade moves back and forth over a distance of 2.0 mm in simple harmonic motion, with frequency 120 Hz. Find (a) the amplitude, (b) the maximum blade speed, and (c) the magnitude of the maximum blade acceleration.
- •6 A particle with a mass of 1.00×10^{-20} kg is oscillating with simple harmonic motion with a period of 1.00×10^{-5} s and a maximum speed of 1.00×10^{3} m/s. Calculate (a) the angular frequency and (b) the maximum displacement of the particle.
- •7 SSM A loudspeaker produces a musical sound by means of the oscillation of a diaphragm whose amplitude is limited to 1.00 μ m. (a) At what frequency is the magnitude a of the diaphragm's acceleration equal to g? (b) For greater frequencies, is a greater than or less than g?



•9 The position function $x = (6.0 \text{ m}) \cos[(3\pi \text{ rad/s})t + \pi/3 \text{ rad}]$ gives the simple harmonic motion of a body. At t = 2.0 s, what are the (a) displacement, (b) velocity, (c)





Figure 15-30 Problem 8.

Hint: what is the position x at t=0 in the graph...

acceleration, and (d) phase of the motion? Also, what are the (e) frequency and (f) period of the motion?

- •10 An oscillating block-spring system takes 0.75 s to begin repeating its motion. Find (a) the period, (b) the frequency in hertz, and (c) the angular frequency in radians per second.
- •11 In Fig. 15-31, two identical springs of spring constant 7580 N/m



are attached to a block of mass 0.245 kg. What is the frequency of oscillation on the frictionless floor?

•12 What is the phase constant for the harmonic oscillator with the velocity function v(t) given in Fig. 15-32 if the position function x(t) has the form $x = x_m \cos(\omega t + \phi)$? The vertical axis scale is set by $v_s = 4.0 \text{ cm/s}$.




Figure 15-32 Problem 12.

•13 SSM An oscillator consists of a block of mass 0.500 kg connected to

a spring. When set into oscillation with amplitude 35.0 cm, the oscillator repeats its motion every 0.500 s. Find the (a) period, (b) frequency, (c) angular frequency, (d) spring constant, (e) maximum speed, and (f) magnitude of the maximum force on the block from the spring.

Hint: For springs in parallel, spring constants add linearly k1+k2. And you need to find f not ω

••14 A simple harmonic oscillator consists of a block of mass 2.00 kg attached to a spring of spring constant 100 N/m. When t = 1.00 s, the position and velocity of the block are x = 0.129 m and v = 3.415 m/s. (a) What is the amplitude of the oscillations? What were the (b) position and (c) velocity of the block at t = 0 s?

••15 SSM Two particles oscillate in simple harmonic motion along a common straight-line segment of length A. Each particle has a period of 1.5 s, but they differ in phase by $\pi/6$ rad. (a) How far apart are they (in terms of A) 0.50 s after the lagging particle leaves one end of the path? (b) Are they then moving in the same direction, toward each other, or away from each other?

••16 Two particles execute simple harmonic motion of the same amplitude and frequency along close parallel lines. They pass each other moving in opposite directions each time their displacement is half their amplitude. What is their phase difference?

16. They pass each other at time t, at $x_1 = x_2 = \frac{1}{2}x_m$ where

```
x_1 = x_m \cos(\omega t + \phi_1) and x_2 = x_m \cos(\omega t + \phi_2). v_1 = -x_m \omega \sin(\omega t + \phi_1) and v_2 = -x_m \omega \sin(\omega t + \phi_2).
```


From this, we conclude that $\cos(\omega t + \phi_1) = \cos(\omega t + \phi_2) = \frac{1}{2}$, and therefore that the phases (the arguments of the cosines) are either both equal to $\pi/3$ or one is $\pi/3$ while the other opposite sign. Thus, one phase is $\pi/3$ and the other phase is $-\pi/3$; the wt term cancels if we take the phase difference, which is seen to be $\pi/3 - (-\pi/3) = 2\pi/3$.

••17 ILW An oscillator consists of a block attached to a spring (k = 400 N/m). At some time t, the position (measured from the system's equilibrium location), velocity, and acceleration of the block are x = 0.100 m, v = -13.6 m/s, and $a = -123 \text{ m/s}^2$. Calculate (a) the frequency of oscillation, (b) the mass of the block, and (c) the amplitude of the motion.

••18 At a certain harbor, the tides cause the ocean surface to rise and fall a distance d (from highest level to lowest level) in simple harmonic motion, with a period of 12.5 h. How long does it take for the water to fall a distance 0.250d from its highest level?

••20 © Figure 15-33a is a partial graph of the position function x(t) for a simple harmonic oscillator with an angular frequency of

1.20 rad/s; Fig. 15-33b is a partial graph of the corresponding velocity function v(t). The vertical axis scales are set by $x_s = 5.0$ cm and $v_s = 5.0$ cm/s. What is the phase constant of the SHM if the position function x(t) is in the general form $x = x_m \cos(\omega t + \phi)$?



Hint: find the phase angle first. divide v(t) by x(t)