

Lecture Notes in Mathematics

An Introduction to Riemannian Geometry

Sigmundur Gudmundsson

(Lund University)

(version 1.0340 - 20th of June 2017)

The latest version of this document can be found at

<http://www.matematik.lu.se/matematiklu/personal/sigma/>

Preface

These lecture notes grew out of an M.Sc. course on differential geometry which I gave at the University of Leeds 1992. Their main purpose is to introduce the beautiful theory of Riemannian geometry a still very active area of mathematical research. This is a subject with no lack of interesting examples. They are indeed the key to a good understanding of it and will therefore play a major role throughout this work. Of special interest are the classical Lie groups allowing concrete calculations of many of the abstract notions on the menu.

The study of Riemannian geometry is rather meaningless without some basic knowledge on Gaussian geometry i.e. the geometry of curves and surfaces in 3-dimensional Euclidean space. For this I recommend the excellent textbook: M. P. do Carmo, *Differential geometry of curves and surfaces*, Prentice Hall (1976).

These lecture notes are written for students with a good understanding of linear algebra, real analysis of several variables, the classical theory of ordinary differential equations and some topology. The most important results stated in the text are also proven there. Others are left to the reader as exercises, which follow at the end of each chapter. This format is aimed at students willing to put **hard work** into the course.

For further reading I recommend the excellent standard text: M. P. do Carmo, *Riemannian Geometry*, Birkhäuser (1992).

I am very grateful to my enthusiastic students and many other readers who have, throughout the years, contributed to the text by giving numerous valuable comments on the presentation.

Norra Nöbbelöv the 23rd of February 2017

Sigmundur Gudmundsson

Contents

Chapter 1.	Introduction	5
Chapter 2.	Differentiable Manifolds	7
Chapter 3.	The Tangent Space	23
Chapter 4.	The Tangent Bundle	39
Chapter 5.	Riemannian Manifolds	55
Chapter 6.	The Levi-Civita Connection	71
Chapter 7.	Geodesics	83
Chapter 8.	The Riemann Curvature Tensor	101
Chapter 9.	Curvature and Local Geometry	115

CHAPTER 1

Introduction

On the 10th of June 1854 Georg Friedrich Bernhard Riemann (1826-1866) gave his famous "Habilitationsvortrag" in the Colloquium of the Philosophical Faculty at Göttingen. His talk "Über die Hypothesen, welche der Geometrie zu Grunde liegen" is often said to be the most important in the history of differential geometry. Johann Carl Friedrich Gauss (1777-1855) was in the audience, at the age of 77, and is said to have been very impressed by his former student.

Riemann's revolutionary ideas generalised the geometry of surfaces which had earlier been initiated by Gauss. Later this lead to an exact definition of the modern concept of an abstract Riemannian manifold.

The development of the 20th century has turned Riemannian geometry into one of the most important parts of modern mathematics. For an excellent survey of this vast field we recommend the following work written by one of the main actors: Marcel Berger, *A Panoramic View of Riemannian Geometry*, Springer (2003).

CHAPTER 2

Differentiable Manifolds

In this chapter we introduce the important notion of a differentiable manifold. This generalises curves and surfaces in \mathbb{R}^3 studied in classical differential geometry. Our manifolds are modelled on the standard differentiable structure on the vector spaces \mathbb{R}^m via compatible charts. We give many examples, study their submanifolds and differentiable maps between manifolds.

Let \mathbb{R}^m be the standard m -dimensional real vector space equipped with the topology induced by the Euclidean metric d on \mathbb{R}^m given by

$$d(x, y) = \sqrt{(x_1 - y_1)^2 + \dots + (x_m - y_m)^2}.$$

For a non-negative integer r and an open subset U of \mathbb{R}^m we will by $C^r(U, \mathbb{R}^n)$ denote the r -times continuously **differentiable** maps from U to \mathbb{R}^n . By **smooth** maps $U \rightarrow \mathbb{R}^n$ we mean the elements of

$$C^\infty(U, \mathbb{R}^n) = \bigcap_{r=0}^{\infty} C^r(U, \mathbb{R}^n).$$

The set of **real analytic** maps from U to \mathbb{R}^n will be denoted by $C^\omega(U, \mathbb{R}^n)$. For the theory of real analytic maps we recommend the book: S. G. Krantz and H. R. Parks, *A Primer of Real Analytic Functions*, Birkhäuser (1992).

Definition 2.1. Let (M, \mathcal{T}) be a topological Hausdorff space with a countable basis. Then M is called a **topological manifold** if there exists an $m \in \mathbb{Z}^+$ such that for each point $p \in M$ we have an open neighbourhood U of p , an open subset V of \mathbb{R}^m and a homeomorphism $x : U \rightarrow V$. The pair (U, x) is called a **chart** (or **local coordinates**) on M . The integer m is called the **dimension** of M . To denote that the dimension of M is m we write M^m .

According to Definition 2.1 a topological manifold M is locally homeomorphic to the standard \mathbb{R}^m for some positive integer m . We will now introduce a differentiable structure on M via its charts and turn M into a differentiable manifold.

Definition 2.2. Let M be a topological manifold of dimension m . Then a C^r -**atlas** on M is a collection

$$\mathcal{A} = \{(U_\alpha, x_\alpha) \mid \alpha \in \mathcal{I}\}$$

of charts on M such that \mathcal{A} covers the whole of M i.e.

$$M = \bigcup_{\alpha} U_\alpha$$

and for all $\alpha, \beta \in \mathcal{I}$ the corresponding **transition maps**

$$x_\beta \circ x_\alpha^{-1}|_{x_\alpha(U_\alpha \cap U_\beta)} : x_\alpha(U_\alpha \cap U_\beta) \subset \mathbb{R}^m \rightarrow \mathbb{R}^m$$

are r -times continuously differentiable i.e. of class C^r .

A chart (U, x) on M is said to be **compatible** with a C^r -atlas \mathcal{A} if the union $\mathcal{A} \cup \{(U, x)\}$ is a C^r -atlas. A C^r -atlas $\hat{\mathcal{A}}$ is said to be **maximal** if it contains all the charts that are compatible with it. A maximal atlas $\hat{\mathcal{A}}$ on M is also called a C^r -**structure** on M . The pair $(M, \hat{\mathcal{A}})$ is said to be a C^r -**manifold**, or a **differentiable manifold** of class C^r , if M is a topological manifold and $\hat{\mathcal{A}}$ is a C^r -structure on M . A differentiable manifold is said to be **smooth** if its transition maps are C^∞ and **real analytic** if they are C^ω .

Remark 2.3. It should be noted that a given C^r -atlas \mathcal{A} on a topological manifold M determines a unique C^r -structure $\hat{\mathcal{A}}$ on M containing \mathcal{A} . It simply consists of all charts compatible with \mathcal{A} .

Example 2.4. For the standard topological space $(\mathbb{R}^m, \mathcal{T}_m)$ we have the trivial C^ω -atlas

$$\mathcal{A} = \{(\mathbb{R}^m, x) \mid x : p \mapsto p\}$$

inducing the standard C^ω -structure $\hat{\mathcal{A}}$ on \mathbb{R}^m .

Example 2.5. Let S^m denote the unit sphere in \mathbb{R}^{m+1} i.e.

$$S^m = \{p \in \mathbb{R}^{m+1} \mid p_1^2 + \cdots + p_{m+1}^2 = 1\}$$

equipped with the subset topology \mathcal{T} induced by the standard \mathcal{T}_{m+1} on \mathbb{R}^{m+1} . Let N be the north pole $N = (1, 0) \in \mathbb{R} \times \mathbb{R}^m$ and S be the south pole $S = (-1, 0)$ on S^m , respectively. Put $U_N = S^m \setminus \{N\}$, $U_S = S^m \setminus \{S\}$ and define $x_N : U_N \rightarrow \mathbb{R}^m$, $x_S : U_S \rightarrow \mathbb{R}^m$ by

$$\begin{aligned} x_N : (p_1, \dots, p_{m+1}) &\mapsto \frac{1}{1 - p_1}(p_2, \dots, p_{m+1}), \\ x_S : (p_1, \dots, p_{m+1}) &\mapsto \frac{1}{1 + p_1}(p_2, \dots, p_{m+1}). \end{aligned}$$

Then the transition maps

$$x_S \circ x_N^{-1}, x_N \circ x_S^{-1} : \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}^m \setminus \{0\}$$

are both given by

$$x \mapsto \frac{x}{|x|^2},$$

so $\mathcal{A} = \{(U_N, x_N), (U_S, x_S)\}$ is a C^ω -atlas on S^m . The C^ω -manifold $(S^m, \hat{\mathcal{A}})$ is called the m -dimensional **standard sphere**.

Another interesting example of a differentiable manifold is the m -dimensional real projective space $\mathbb{R}P^m$.

Example 2.6. On the set $\mathbb{R}^{m+1} \setminus \{0\}$ we define the equivalence relation \equiv by

$$p \equiv q \text{ if and only if there exists a } \lambda \in \mathbb{R}^* \text{ such that } p = \lambda q.$$

Let $\mathbb{R}P^m$ be the quotient space $(\mathbb{R}^{m+1} \setminus \{0\}) / \equiv$ and

$$\pi : \mathbb{R}^{m+1} \setminus \{0\} \rightarrow \mathbb{R}P^m$$

be the natural projection mapping a point $p \in \mathbb{R}^{m+1} \setminus \{0\}$ onto the equivalence class $[p] \in \mathbb{R}P^m$ i.e. the line

$$[p] = \{\lambda p \in \mathbb{R}^{m+1} \mid \lambda \in \mathbb{R}^*\}$$

through the origin generated by p . Equip $\mathbb{R}P^m$ with the quotient topology \mathcal{T} induced by π and \mathcal{T}_{m+1} on \mathbb{R}^{m+1} . For $k \in \{1, \dots, m+1\}$ define the open subset

$$U_k = \{[p] \in \mathbb{R}P^m \mid p_k \neq 0\}$$

of $\mathbb{R}P^m$ and the charts $x_k : U_k \subset \mathbb{R}P^m \rightarrow \mathbb{R}^m$ by

$$x_k : [p] \mapsto \left(\frac{p_1}{p_k}, \dots, \frac{p_{k-1}}{p_k}, 1, \frac{p_{k+1}}{p_k}, \dots, \frac{p_{m+1}}{p_k} \right).$$

If $[p] = [q]$ then $p = \lambda q$ for some $\lambda \in \mathbb{R}^*$ so $p_l/p_k = q_l/q_k$ for all l . This means that the map x_k is well defined for all k . The corresponding transition maps

$$x_k \circ x_l^{-1}|_{x_l(U_l \cap U_k)} : x_l(U_l \cap U_k) \subset \mathbb{R}^m \rightarrow \mathbb{R}^m$$

are given by

$$\left(\frac{p_1}{p_l}, \dots, \frac{p_{l-1}}{p_l}, 1, \frac{p_{l+1}}{p_l}, \dots, \frac{p_{m+1}}{p_l} \right) \mapsto \left(\frac{p_1}{p_k}, \dots, \frac{p_{k-1}}{p_k}, 1, \frac{p_{k+1}}{p_k}, \dots, \frac{p_{m+1}}{p_k} \right)$$

so the collection

$$\mathcal{A} = \{(U_k, x_k) \mid k = 1, \dots, m+1\}$$

is a C^ω -atlas on $\mathbb{R}P^m$. The differentiable manifold $(\mathbb{R}P^m, \hat{\mathcal{A}})$ is called the m -dimensional **real projective space**.

Example 2.7. Let $\hat{\mathbb{C}}$ be the extended complex plane given by

$$\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$$

and put $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$, $U_0 = \mathbb{C}$ and $U_\infty = \hat{\mathbb{C}} \setminus \{0\}$. Then define the charts $x_0 : U_0 \rightarrow \mathbb{C}$ and $x_\infty : U_\infty \rightarrow \mathbb{C}$ on $\hat{\mathbb{C}}$ by $x_0 : z \mapsto z$ and $x_\infty : w \mapsto 1/w$, respectively. The corresponding transition maps

$$x_\infty \circ x_0^{-1}, x_0 \circ x_\infty^{-1} : \mathbb{C}^* \rightarrow \mathbb{C}^*$$

are both given by $z \mapsto 1/z$ so $\mathcal{A} = \{(U_0, x_0), (U_\infty, x_\infty)\}$ is a C^ω -atlas on $\hat{\mathbb{C}}$. The real analytic manifold $(\hat{\mathbb{C}}, \hat{\mathcal{A}})$ is called the **Riemann sphere**.

For the product of two differentiable manifolds we have the following important result.

Proposition 2.8. *Let $(M_1, \hat{\mathcal{A}}_1)$ and $(M_2, \hat{\mathcal{A}}_2)$ be two differentiable manifolds of class C^r . Let $M = M_1 \times M_2$ be the product space with the product topology. Then there exists an atlas \mathcal{A} on M turning $(M, \hat{\mathcal{A}})$ into a differentiable manifold of class C^r and the dimension of M satisfies*

$$\dim M = \dim M_1 + \dim M_2.$$

PROOF. See Exercise 2.1. □

The concept of a submanifold of a given differentiable manifold will play an important role as we go along and we will be especially interested in the connection between the geometry of a submanifold and that of its ambient space.

Definition 2.9. Let m, n be positive integers with $m \leq n$ and $(N^n, \hat{\mathcal{A}}_N)$ be a C^r -manifold. A subset M of N is said to be a **submanifold** of N if for each point $p \in M$ there exists a chart $(U_p, x_p) \in \hat{\mathcal{A}}_N$ such that $p \in U_p$ and $x_p : U_p \subset N \rightarrow \mathbb{R}^m \times \mathbb{R}^{n-m}$ satisfies

$$x_p(U_p \cap M) = x_p(U_p) \cap (\mathbb{R}^m \times \{0\}).$$

The natural number $(n - m)$ is called the **codimension** of M in N .

Proposition 2.10. *Let m, n be positive integers with $m \leq n$ and $(N^n, \hat{\mathcal{A}}_N)$ be a C^r -manifold. Let M be a submanifold of N equipped with the subset topology and $\pi : \mathbb{R}^m \times \mathbb{R}^{n-m} \rightarrow \mathbb{R}^m$ be the natural projection onto the first factor. Then*

$$\mathcal{A}_M = \{(U_p \cap M, (\pi \circ x_p)|_{U_p \cap M}) \mid p \in M\}$$

*is a C^r -atlas for M . Hence the pair $(M, \hat{\mathcal{A}}_M)$ is an m -dimensional C^r -manifold. The differentiable structure $\hat{\mathcal{A}}_M$ on M is called the **induced structure** by $\hat{\mathcal{A}}_N$.*

PROOF. See Exercise 2.2. \square

Remark 2.11. Our next aim is to prove Theorem 2.15 which is a useful tool for the construction of submanifolds of \mathbb{R}^m . For this we use the classical inverse mapping theorem stated below. Note that if

$$F : U \rightarrow \mathbb{R}^n$$

is a differentiable map defined on an open subset U of \mathbb{R}^m then its differential $dF_p : \mathbb{R}^m \rightarrow \mathbb{R}^n$ at the point $p \in U$ is a linear map given by the $n \times m$ matrix

$$dF_p = \begin{pmatrix} \partial F_1 / \partial x_1(p) & \dots & \partial F_1 / \partial x_m(p) \\ \vdots & & \vdots \\ \partial F_n / \partial x_1(p) & \dots & \partial F_n / \partial x_m(p) \end{pmatrix}.$$

If $\gamma : \mathbb{R} \rightarrow U$ is a curve in U such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v \in \mathbb{R}^m$ then the composition $F \circ \gamma : \mathbb{R} \rightarrow \mathbb{R}^n$ is a curve in \mathbb{R}^n and according to the chain rule we have

$$dF_p \cdot v = \frac{d}{ds}(F \circ \gamma(s))|_{s=0}.$$

This is the tangent vector of the curve $F \circ \gamma$ at $F(p) \in \mathbb{R}^n$.

The above shows that the differential dF_p can be seen as a linear map that maps tangent vectors at $p \in U$ to tangent vectors at the image point $F(p) \in \mathbb{R}^n$. This will later be generalised to the manifold setting.

We now state the classical inverse mapping theorem well known from multivariable analysis.

Fact 2.12. *Let U be an open subset of \mathbb{R}^m and $F : U \rightarrow \mathbb{R}^m$ be a C^r -map. If $p \in U$ and the differential*

$$dF_p : \mathbb{R}^m \rightarrow \mathbb{R}^m$$

of F at p is invertible then there exist open neighbourhoods U_p around p and U_q around $q = F(p)$ such that $\hat{F} = F|_{U_p} : U_p \rightarrow U_q$ is bijective and the inverse $(\hat{F})^{-1} : U_q \rightarrow U_p$ is a C^r -map. The differential $(d\hat{F}^{-1})_q$ of \hat{F}^{-1} at q satisfies

$$(d\hat{F}^{-1})_q = (dF_p)^{-1}$$

i.e. it is the inverse of the differential dF_p of F at p .

Before stating the classical implicit mapping theorem we remind the reader of the following well known notions.

Definition 2.13. Let m, n be positive natural numbers, U be an open subset of \mathbb{R}^m and $F : U \rightarrow \mathbb{R}^n$ be a C^r -map. A point $p \in U$ is said to be **regular** for F , if the differential

$$dF_p : \mathbb{R}^m \rightarrow \mathbb{R}^n$$

is of full rank, but **critical** otherwise. A point $q \in F(U)$ is said to be a **regular value** of F if every point in the pre-image $F^{-1}(\{q\})$ of q is regular.

Remark 2.14. Note that if m, n are positive integers with $m \geq n$ then $p \in U$ is a regular point for

$$F = (F_1, \dots, F_n) : U \rightarrow \mathbb{R}^n$$

if and only if the gradients $\text{grad}F_1, \dots, \text{grad}F_n$ of the coordinate functions $F_1, \dots, F_n : U \rightarrow \mathbb{R}$ are linearly independent at p , or equivalently, the differential dF_p of F at p satisfies the following condition

$$\det(dF_p \cdot (dF_p)^t) \neq 0.$$

Theorem 2.15 (The Implicit Mapping Theorem). *Let m, n be positive integers with $m > n$ and $F : U \rightarrow \mathbb{R}^n$ be a C^r -map from an open subset U of \mathbb{R}^m . If $q \in F(U)$ is a regular value of F then the pre-image $F^{-1}(\{q\})$ of q is an $(m - n)$ -dimensional submanifold of \mathbb{R}^m of class C^r .*

PROOF. Let p be an element of $F^{-1}(\{q\})$ and K_p be the kernel of the differential dF_p i.e. the $(m - n)$ -dimensional subspace of \mathbb{R}^m given by $K_p = \{v \in \mathbb{R}^m \mid dF_p \cdot v = 0\}$. Let $\pi_p : \mathbb{R}^m \rightarrow \mathbb{R}^{m-n}$ be a linear map such that $\pi_p|_{K_p} : K_p \rightarrow \mathbb{R}^{m-n}$ is bijective, $\pi_p|_{K_p^\perp} = 0$ and define the map $G_p : U \rightarrow \mathbb{R}^n \times \mathbb{R}^{m-n}$ by

$$G_p : x \mapsto (F(x), \pi_p(x)).$$

Then the differential $(dG_p)_p : \mathbb{R}^m \rightarrow \mathbb{R}^m$ of G_p , with respect to the decompositions $\mathbb{R}^m = K_p^\perp \oplus K_p$ and $\mathbb{R}^m = \mathbb{R}^n \oplus \mathbb{R}^{m-n}$, is given by

$$(dG_p)_p = \begin{pmatrix} dF_p|_{K_p^\perp} & 0 \\ 0 & \pi_p \end{pmatrix},$$

hence bijective. It now follows from the inverse function theorem that there exist open neighbourhoods V_p around p and W_p around $G_p(p)$ such that $\hat{G}_p = G_p|_{V_p} : V_p \rightarrow W_p$ is bijective, the inverse $\hat{G}_p^{-1} : W_p \rightarrow V_p$ is C^r , $d(\hat{G}_p^{-1})_{G_p(p)} = (dG_p)_p^{-1}$ and $d(\hat{G}_p^{-1})_y$ is bijective for all $y \in W_p$. Now put $\tilde{U}_p = F^{-1}(\{q\}) \cap V_p$ then

$$\tilde{U}_p = \hat{G}_p^{-1}((\{q\} \times \mathbb{R}^{m-n}) \cap W_p)$$

so if $\pi : \mathbb{R}^n \times \mathbb{R}^{m-n} \rightarrow \mathbb{R}^{m-n}$ is the natural projection onto the second factor, then the map

$$\tilde{x}_p = \pi \circ G_p|_{\tilde{U}_p} : \tilde{U}_p \rightarrow (\{q\} \times \mathbb{R}^{m-n}) \cap W_p \rightarrow \mathbb{R}^{m-n}$$

is a chart on the open neighbourhood \tilde{U}_p of p . The point $q \in F(U)$ is a regular value so the set

$$\mathcal{A} = \{(\tilde{U}_p, \tilde{x}_p) \mid p \in F^{-1}(\{q\})\}$$

is a C^r -atlas for $F^{-1}(\{q\})$. \square

Employing the implicit function theorem, we obtain the following interesting examples of the m -dimensional sphere S^m and its tangent bundle TS^m as differentiable submanifolds of \mathbb{R}^{m+1} and \mathbb{R}^{2m+2} , respectively.

Example 2.16. Let $F : \mathbb{R}^{m+1} \rightarrow \mathbb{R}$ be the C^ω -map given by

$$F : (p_1, \dots, p_{m+1}) \mapsto p_1^2 + \dots + p_{m+1}^2.$$

Then the differential dF_p of F at p is given by $dF_p = 2p$, so

$$dF_p \cdot (dF_p)^t = 4|p|^2 \in \mathbb{R}.$$

This means that $1 \in \mathbb{R}$ is a regular value of F so the fibre

$$S^m = \{p \in \mathbb{R}^{m+1} \mid |p|^2 = 1\} = F^{-1}(\{1\})$$

of F is an m -dimensional submanifold of \mathbb{R}^{m+1} . This is the standard m -dimensional sphere introduced in Example 2.5.

Example 2.17. Let $F : \mathbb{R}^{m+1} \times \mathbb{R}^{m+1} \rightarrow \mathbb{R}^2$ be the C^ω -map defined by $F : (p, v) \mapsto (|p|^2 - 1)/2, \langle p, v \rangle)$. Then the differential $dF_{(p,v)}$ of F at (p, v) satisfies

$$dF_{(p,v)} = \begin{pmatrix} p & 0 \\ v & p \end{pmatrix}.$$

A simple calculation shows that

$$\det(dF \cdot (dF)^t) = \det \begin{pmatrix} |p|^2 & \langle p, v \rangle \\ \langle p, v \rangle & |v|^2 + |p|^2 \end{pmatrix} = 1 + |v|^2 > 0$$

on $F^{-1}(\{0\})$. This means that

$$F^{-1}(\{0\}) = \{(p, v) \in \mathbb{R}^{m+1} \times \mathbb{R}^{m+1} \mid |p|^2 = 1 \text{ and } \langle p, v \rangle = 0\},$$

which we denote by TS^m , is a $2m$ -dimensional submanifold of \mathbb{R}^{2m+2} . We will later see that TS^m is what is called the **tangent bundle** of the m -dimensional sphere S^m .

We now apply the implicit function theorem to construct the important orthogonal group $\mathbf{O}(m)$ as a submanifold of the set of the real vector space of $m \times m$ matrices $\mathbb{R}^{m \times m}$.

Example 2.18. Let $\text{Sym}(\mathbb{R}^m)$ be the $m(m+1)/2$ dimensional linear subspace of $\mathbb{R}^{m \times m}$ consisting of all symmetric $m \times m$ matrices

$$\text{Sym}(\mathbb{R}^m) = \{y \in \mathbb{R}^{m \times m} \mid y^t = y\}.$$

Let $F : \mathbb{R}^{m \times m} \rightarrow \text{Sym}(\mathbb{R}^m)$ be the map defined by

$$F : x \mapsto x^t x.$$

If $\gamma : I \rightarrow \mathbb{R}^{m \times m}$ is a curve in $\mathbb{R}^{m \times m}$ then

$$\frac{d}{ds}(F \circ \gamma(s)) = \dot{\gamma}(s)^t \gamma(s) + \gamma(s)^t \dot{\gamma}(s),$$

so the differential dF_x of F at $x \in \mathbb{R}^{m \times m}$ satisfies

$$dF_x : X \mapsto X^t x + x^t X.$$

This means that for an arbitrary element p in

$$\mathbf{O}(m) = F^{-1}(\{e\}) = \{p \in \mathbb{R}^{m \times m} \mid p^t p = e\}$$

and $Y \in \text{Sym}(\mathbb{R}^m)$ we have $dF_p(pY/2) = Y$. Hence the differential dF_p is surjective, so the identity matrix $e \in \text{Sym}(\mathbb{R}^m)$ is a regular value of F . Following the implicit function theorem $\mathbf{O}(m)$ is a submanifold of $\mathbb{R}^{m \times m}$ of dimension $m(m-1)/2$. The set $\mathbf{O}(m)$ is the well known **orthogonal group**.

The concept of a differentiable map $U \rightarrow \mathbb{R}^n$, defined on an open subset of \mathbb{R}^m , can be generalised to mappings between manifolds. We will see that the most important properties of these objects in the classical case are also valid in the manifold setting.

Definition 2.19. Let $(M^m, \hat{\mathcal{A}}_M)$ and $(N^n, \hat{\mathcal{A}}_N)$ be C^r -manifolds. A map $\phi : M \rightarrow N$ is said to be **differentiable** of class C^r if for all charts $(U, x) \in \hat{\mathcal{A}}_M$ and $(V, y) \in \hat{\mathcal{A}}_N$ the map

$$y \circ \phi \circ x^{-1}|_{x(U \cap \phi^{-1}(V))} : x(U \cap \phi^{-1}(V)) \subset \mathbb{R}^m \rightarrow \mathbb{R}^n$$

is of class C^r . A differentiable map $\gamma : I \rightarrow M$ defined on an open interval of \mathbb{R} is called a differentiable **curve** in M . A differentiable map $f : M \rightarrow \mathbb{R}$ with values in \mathbb{R} is called a differentiable **function** on M . The set of smooth functions defined on M is denoted by $C^\infty(M)$.

It is an easy exercise, using Definition 2.19, to prove the following result concerning the composition of differentiable maps between manifolds.

Proposition 2.20. *Let $(M_1, \hat{\mathcal{A}}_1), (M_2, \hat{\mathcal{A}}_2), (M_3, \hat{\mathcal{A}}_3)$ be C^r -manifolds and $\phi : (M_1, \hat{\mathcal{A}}_1) \rightarrow (M_2, \hat{\mathcal{A}}_2)$, $\psi : (M_2, \hat{\mathcal{A}}_2) \rightarrow (M_3, \hat{\mathcal{A}}_3)$ be differentiable maps of class C^r . Then the composition $\psi \circ \phi : (M_1, \hat{\mathcal{A}}_1) \rightarrow (M_3, \hat{\mathcal{A}}_3)$ is a differentiable map of class C^r .*

PROOF. See Exercise 2.5. \square

Definition 2.21. Two manifolds $(M, \hat{\mathcal{A}}_M)$ and $(N, \hat{\mathcal{A}}_N)$ of class C^r are said to be **diffeomorphic** if there exists a bijective C^r -map $\phi : M \rightarrow N$ such that the inverse $\phi^{-1} : N \rightarrow M$ is of class C^r . In that case the map ϕ is called a **diffeomorphism** between $(M, \hat{\mathcal{A}}_M)$ and $(N, \hat{\mathcal{A}}_N)$.

Proposition 2.22. *Let $(M, \hat{\mathcal{A}})$ be an m -dimensional C^r -manifold and (U, x) be a chart on M . Then the bijective continuous map $x : U \rightarrow x(U) \subset \mathbb{R}^m$ is a diffeomorphism.*

PROOF. See Exercise 2.6. \square

It can be shown that the 2-dimensional sphere S^2 in \mathbb{R}^3 and the Riemann sphere $\hat{\mathbb{C}}$ are diffeomorphic, see Exercise 2.7.

Definition 2.23. For a differentiable manifold $(M, \hat{\mathcal{A}})$ we denote by $\mathcal{D}(M)$ the set of all its diffeomorphisms. If $\phi, \psi \in \mathcal{D}(M)$ then it is clear that the composition $\psi \circ \phi$ and the inverse ϕ^{-1} are also diffeomorphisms. The pair $(\mathcal{D}(M), \circ)$ is called the **diffeomorphism group** of $(M, \hat{\mathcal{A}})$. The operation is clearly associative and the identity map is its neutral element.

Definition 2.24. Two C^r -structures $\hat{\mathcal{A}}_1$ and $\hat{\mathcal{A}}_2$ on the same topological manifold M are said to be **different** if the identity map $\text{id}_M : (M, \hat{\mathcal{A}}_1) \rightarrow (M, \hat{\mathcal{A}}_2)$ is not a diffeomorphism.

It can be seen that even the real line \mathbb{R} carries infinitely many different differentiable structures, see Exercise 2.8.

Deep Result 2.25. *Let $(M, \hat{\mathcal{A}}_M), (N, \hat{\mathcal{A}}_N)$ be differentiable manifolds of class C^r of the same dimension m . If M and N are homeomorphic as topological spaces and $m \leq 3$ then $(M, \hat{\mathcal{A}}_M)$ and $(N, \hat{\mathcal{A}}_N)$ are diffeomorphic.*

The following remarkable result was proven by John Milnor in his famous paper: *Differentiable structures on spheres*, Amer. J. Math. **81** (1959), 962-972.

Deep Result 2.26. *The 7-dimensional sphere S^7 has exactly 28 different differentiable structures.*

The next very useful statement generalises a classical result from real analysis of several variables.

Proposition 2.27. *Let $(N_1, \hat{\mathcal{A}}_1)$ and $(N_2, \hat{\mathcal{A}}_2)$ be two differentiable manifolds of class C^r and M_1, M_2 be submanifolds of N_1 and N_2 , respectively. If $\phi : N_1 \rightarrow N_2$ is a differentiable map of class C^r such that $\phi(M_1)$ is contained in M_2 then the restriction $\phi|_{M_1} : M_1 \rightarrow M_2$ is differentiable of class C^r .*

PROOF. See Exercise 2.9. □

Example 2.28. The following maps are all smooth.

- (i) $\phi_1 : \mathbb{R}^1 \rightarrow S^1 \subset \mathbb{C}, \phi_1 : t \mapsto e^{it}$,
- (ii) $\phi_2 : \mathbb{R}^{m+1} \setminus \{0\} \rightarrow S^m \subset \mathbb{R}^{m+1}, \phi_2 : x \mapsto x/|x|$,
- (iii) $\phi_3 : S^2 \subset \mathbb{R}^3 \rightarrow S^3 \subset \mathbb{R}^4, \phi_3 : (x, y, z) \mapsto (x, y, z, 0)$,
- (iv) $\phi_4 : S^3 \subset \mathbb{C}^2 \rightarrow S^2 \subset \mathbb{C} \times \mathbb{R}, \phi_4 : (z_1, z_2) \mapsto (2z_1\bar{z}_2, |z_1|^2 - |z_2|^2)$,
- (v) $\phi_6 : S^m \rightarrow \mathbb{R}P^m, \phi_6 : x \mapsto [x]$.
- (vi) $\phi_5 : \mathbb{R}^{m+1} \setminus \{0\} \rightarrow \mathbb{R}P^m, \phi_5 : x \mapsto [x]$,

In differential geometry we are especially interested in manifolds carrying a group structure compatible with their differentiable structures. Such manifolds are named after the famous mathematician Sophus Lie (1842-1899) and will play an important role throughout this work.

Definition 2.29. A **Lie group** is a smooth manifold G with a group structure \cdot such that the map $\rho : G \times G \rightarrow G$ with

$$\rho : (p, q) \mapsto p \cdot q^{-1}$$

is smooth. For an element p in G the **left translation** by p is the map $L_p : G \rightarrow G$ defined by $L_p : q \mapsto p \cdot q$.

Example 2.30. Let $(\mathbb{R}^m, +, \cdot)$ be the m -dimensional vector space equipped with its standard differential structure. Then $(\mathbb{R}^m, +)$ with $\rho : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ given by

$$\rho : (p, q) \mapsto p - q$$

is a Lie group.

For Lie groups, we now introduce the left-translations. These will play a very important role in what follows.

Proposition 2.31. *Let G be a Lie group and p be an element of G . Then the left translation $L_p : G \rightarrow G$ is a smooth diffeomorphism.*

PROOF. See Exercise 2.11 □

Proposition 2.32. *Let (G, \cdot) be a Lie group and K be a submanifold of G which is a subgroup. Then (K, \cdot) is a Lie group.*

PROOF. The statement is a direct consequence of Definition 2.29 and Proposition 2.27. \square

The set of non-zero complex numbers (\mathbb{C}^*, \cdot) , together with the standard multiplication, form a Lie group. The unit circle (S^1, \cdot) is an interesting compact Lie subgroup of (\mathbb{C}^*, \cdot) . Another subgroup is the set of the non-zero real numbers (\mathbb{R}^*, \cdot) containing the positive real numbers (\mathbb{R}^+, \cdot) as a subgroup.

Definition 2.33. Let (G, \cdot) be a group and V be a vector space. Then a **linear representation** of G on V is a map $\rho : G \rightarrow \mathbf{Aut}(V)$ into the space of automorphisms of V i.e. the invertible linear endomorphisms such that for all $g, h \in G$ we have

$$\rho(g \cdot h) = \rho(g) * \rho(h).$$

Here $*$ denotes the matrix multiplication in $\mathbf{Aut}(V)$.

Example 2.34. The Lie group of non-zero complex numbers (\mathbb{C}^*, \cdot) has a well known linear representation $\rho : \mathbb{C}^* \rightarrow \mathbf{Aut}(\mathbb{R}^2)$ given by

$$\rho : a + ib \mapsto \begin{pmatrix} a & b \\ -b & a \end{pmatrix}.$$

This is obviously injective and it respects the standard multiplicative structures of \mathbb{C}^* and $\mathbb{R}^{2 \times 2}$ i.e.

$$\begin{aligned} \rho((a + ib) \cdot (x + iy)) &= \rho((ax - by) + i(ay + bx)) \\ &= \begin{pmatrix} ax - by & ay + bx \\ -(ay + bx) & ax - by \end{pmatrix} \\ &= \begin{pmatrix} a & b \\ -b & a \end{pmatrix} * \begin{pmatrix} x & y \\ -y & x \end{pmatrix} \\ &= \rho(a + ib) * \rho(x + iy). \end{aligned}$$

As an introduction to Example 2.36 we now play the same game in the complex case.

Example 2.35. Let $\rho : \mathbb{C}^2 \rightarrow \mathbb{C}^{2 \times 2}$ be the real linear map given by

$$\rho : (z, w) \mapsto \begin{pmatrix} z & w \\ -\bar{w} & \bar{z} \end{pmatrix}.$$

Then an easy calculation shows that the following is true

$$\rho(z_1, w_1) * \rho(z_2, w_2) = \begin{pmatrix} z_1 & w_1 \\ -\bar{w}_1 & \bar{z}_1 \end{pmatrix} * \begin{pmatrix} z_2 & w_2 \\ -\bar{w}_2 & \bar{z}_2 \end{pmatrix}$$

$$\begin{aligned}
&= \begin{pmatrix} z_1 z_2 - w_1 \bar{w}_2 & z_1 w_2 + w_1 \bar{z}_2 \\ -\bar{z}_1 \bar{w}_2 - \bar{w}_1 z_2 & \bar{z}_1 \bar{z}_2 - \bar{w}_1 w_2 \end{pmatrix} \\
&= \rho(z_1 z_2 - w_1 \bar{w}_2, z_1 w_2 + w_1 \bar{z}_2).
\end{aligned}$$

We now introduce the important quaternions and the three dimensional sphere which carries a natural group structure.

Example 2.36. Let \mathbb{H} be the set of **quaternions** given by

$$\mathbb{H} = \{z + wj \mid z, w \in \mathbb{C}\} \cong \mathbb{C}^2.$$

We equip \mathbb{H} with an addition, a multiplication and the conjugation satisfying

- (i) $\overline{(z + wj)} = \bar{z} - wj$,
- (ii) $(z_1 + w_1 j) + (z_2 + w_2 j) = (z_1 + z_2) + (w_1 + w_2)j$,
- (iii) $(z_1 + w_1 j) \cdot (z_2 + w_2 j) = (z_1 z_2 - w_1 \bar{w}_2) + (z_1 w_2 + w_1 \bar{z}_2)j$.

These extend the standard operations on \mathbb{C} as a subset of \mathbb{H} . It is easily seen that the non-zero quaternions (\mathbb{H}^*, \cdot) form a Lie group. On \mathbb{H} we define the quaternionic scalar product

$$\mathbb{H} \times \mathbb{H} \rightarrow \mathbb{H}, \quad (p, q) \mapsto p \cdot \bar{q}$$

and a real-valued norm given by $|p|^2 = p \cdot \bar{p}$. Then the 3-dimensional unit sphere S^3 in $\mathbb{H} \cong \mathbb{C}^2 \cong \mathbb{R}^4$, with the restricted multiplication, forms a compact Lie subgroup (S^3, \cdot) of (\mathbb{H}^*, \cdot) . They are both non-abelian.

We will now introduce some of the classical real and complex matrix Lie groups. As a reference on this topic we recommend the wonderful book: A. W. Knap, *Lie Groups Beyond an Introduction*, Birkhäuser (2002).

Example 2.37. Let Nil be the subset of $\mathbb{R}^{3 \times 3}$ given by

$$\text{Nil} = \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \in \mathbb{R}^{3 \times 3} \mid x, y, z \in \mathbb{R} \right\}.$$

Then Nil has a natural differentiable structure determined by the global coordinates $\phi : \text{Nil} \rightarrow \mathbb{R}^3$ with

$$\phi : \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mapsto (x, y, z).$$

It is easily seen that if $*$ is the standard matrix multiplication, then $(\text{Nil}, *)$ is a Lie group.

Example 2.38. Let Sol be the subset of $\mathbb{R}^{3 \times 3}$ given by

$$\text{Sol} = \left\{ \begin{pmatrix} e^z & 0 & x \\ 0 & e^{-z} & y \\ 0 & 0 & 1 \end{pmatrix} \in \mathbb{R}^{3 \times 3} \mid x, y, z \in \mathbb{R} \right\}.$$

Then Sol has a natural differentiable structure determined by the global coordinates $\phi : \text{Sol} \rightarrow \mathbb{R}^3$ with

$$\phi : \begin{pmatrix} e^z & 0 & x \\ 0 & e^{-z} & y \\ 0 & 0 & 1 \end{pmatrix} \mapsto (x, y, z).$$

It is easily seen that if $*$ is the standard matrix multiplication, then $(\text{Sol}, *)$ is a Lie group.

Example 2.39. The set of invertible real $m \times m$ matrices

$$\mathbf{GL}_m(\mathbb{R}) = \{A \in \mathbb{R}^{m \times m} \mid \det A \neq 0\},$$

equipped with the standard matrix multiplication, has the structure of a Lie group. It is called the **real general linear group** and its neutral element e is the identity matrix. The subset $\mathbf{GL}_m(\mathbb{R})$ of $\mathbb{R}^{m \times m}$ is open so $\dim \mathbf{GL}_m(\mathbb{R}) = m^2$.

As a subgroup of $\mathbf{GL}_m(\mathbb{R})$ we have the **real special linear group** $\mathbf{SL}_m(\mathbb{R})$ given by

$$\mathbf{SL}_m(\mathbb{R}) = \{A \in \mathbb{R}^{m \times m} \mid \det A = 1\}.$$

We will show in Example 3.10 that the dimension of the submanifold $\mathbf{SL}_m(\mathbb{R})$ of $\mathbb{R}^{m \times m}$ is $m^2 - 1$.

Another subgroup of $\mathbf{GL}_m(\mathbb{R})$ is the **orthogonal group**

$$\mathbf{O}(m) = \{A \in \mathbb{R}^{m \times m} \mid A^t A = e\}.$$

As we have already seen in Example 2.18 this is a submanifold of $\mathbb{R}^{m \times m}$ of dimension of $m(m-1)/2$.

As a subgroup of $\mathbf{O}(m)$ and even $\mathbf{SL}_m(\mathbb{R})$ we have the **special orthogonal group** $\mathbf{SO}(m)$ which is defined as

$$\mathbf{SO}(m) = \mathbf{O}(m) \cap \mathbf{SL}_m(\mathbb{R}).$$

It can be shown that $\mathbf{O}(m)$ is diffeomorphic to $\mathbf{SO}(m) \times \mathbf{O}(1)$, see Exercise 2.10. Note that $\mathbf{O}(1) = \{\pm 1\}$ so $\mathbf{O}(m)$ can be seen as double cover of $\mathbf{SO}(m)$. This means that

$$\dim \mathbf{SO}(m) = \dim \mathbf{O}(m) = m(m-1)/2.$$

To the above mentioned real Lie groups we have their following complex close relatives.

Example 2.40. The set of invertible complex $m \times m$ matrices

$$\mathbf{GL}_m(\mathbb{C}) = \{A \in \mathbb{C}^{m \times m} \mid \det A \neq 0\},$$

equipped with the standard matrix multiplication, has the structure of a Lie group. It is called the **complex general linear group** and its neutral element e is the identity matrix. The subset $\mathbf{GL}_m(\mathbb{C})$ of $\mathbb{C}^{m \times m}$ is open so $\dim(\mathbf{GL}_m(\mathbb{C})) = 2m^2$.

As a subgroup of $\mathbf{GL}_m(\mathbb{C})$ we have the **complex special linear group** $\mathbf{SL}_m(\mathbb{C})$ given by

$$\mathbf{SL}_m(\mathbb{C}) = \{A \in \mathbb{C}^{m \times m} \mid \det A = 1\}.$$

The dimension of the submanifold $\mathbf{SL}_m(\mathbb{C})$ of $\mathbb{C}^{m \times m}$ is $2(m^2 - 1)$.

Another subgroup of $\mathbf{GL}_m(\mathbb{C})$ is the **unitary group** $\mathbf{U}(m)$ given by

$$\mathbf{U}(m) = \{A \in \mathbb{C}^{m \times m} \mid \bar{A}^t A = e\}.$$

Calculations similar to those for the orthogonal group show that the dimension of $\mathbf{U}(m)$ is m^2 .

As a subgroup of $\mathbf{U}(m)$ and $\mathbf{SL}_m(\mathbb{C})$ we have the **special unitary group** $\mathbf{SU}(m)$ which is defined as

$$\mathbf{SU}(m) = \mathbf{U}(m) \cap \mathbf{SL}_m(\mathbb{C}).$$

It can be shown that $\mathbf{U}(1)$ is diffeomorphic to the circle S^1 and that $\mathbf{U}(m)$ is diffeomorphic to $\mathbf{SU}(m) \times \mathbf{U}(1)$, see Exercise 2.10. This means that $\dim \mathbf{SU}(m) = m^2 - 1$.

For the rest of this work we will assume, when not stating otherwise, that our manifolds and maps are smooth i.e. in the C^∞ -category.

Exercises

Exercise 2.1. Find a proof of Proposition 2.8.

Exercise 2.2. Find a proof of Proposition 2.10.

Exercise 2.3. Let S^1 be the unit circle in the complex plane \mathbb{C} given by $S^1 = \{z \in \mathbb{C} \mid |z|^2 = 1\}$. Use the maps $x : \mathbb{C} \setminus \{i\} \rightarrow \mathbb{C}$ and $y : \mathbb{C} \setminus \{-i\} \rightarrow \mathbb{C}$ with

$$x : z \mapsto \frac{i+z}{1+iz}, \quad y : z \mapsto \frac{1+iz}{i+z}$$

to show that S^1 is a 1-dimensional submanifold of $\mathbb{C} \cong \mathbb{R}^2$.

Exercise 2.4. Use the implicit function theorem to show that the m -dimensional **torus**

$$\begin{aligned} T^m &= \{(x, y) \in \mathbb{R}^m \times \mathbb{R}^m \mid x_1^2 + y_1^2 = \cdots = x_m^2 + y_m^2 = 1\} \\ &\cong \{z \in \mathbb{C}^m \mid |z_1|^2 = \cdots = |z_m|^2 = 1\} \end{aligned}$$

is a differentiable submanifold of $\mathbb{R}^{2m} \cong \mathbb{C}^m$.

Exercise 2.5. Find a proof of Proposition 2.20.

Exercise 2.6. Find a proof of Proposition 2.22.

Exercise 2.7. Prove that the 2-dimensional sphere S^2 as a differentiable submanifold of the standard \mathbb{R}^3 and the Riemann sphere $\hat{\mathbb{C}}$ are diffeomorphic.

Exercise 2.8. Equip the real line \mathbb{R} with the standard topology and for each odd integer $k \in \mathbb{Z}^+$ let $\hat{\mathcal{A}}_k$ be the C^ω -structure defined on \mathbb{R} by the atlas

$$\mathcal{A}_k = \{(\mathbb{R}, x_k) \mid x_k : p \mapsto p^k\}.$$

Show that the differentiable structures $\hat{\mathcal{A}}_k$ are all different but that the differentiable manifolds $(\mathbb{R}, \hat{\mathcal{A}}_k)$ are all diffeomorphic.

Exercise 2.9. Find a proof of Proposition 2.27.

Exercise 2.10. Let the spheres S^1 , S^3 and the Lie groups $\mathbf{SO}(n)$, $\mathbf{O}(n)$, $\mathbf{SU}(n)$, $\mathbf{U}(n)$ be equipped with their standard differentiable structures. Use Proposition 2.27 to prove the following diffeomorphisms

$$S^1 \cong \mathbf{SO}(2), \quad S^3 \cong \mathbf{SU}(2),$$

$$\mathbf{SO}(n) \times \mathbf{O}(1) \cong \mathbf{O}(n), \quad \mathbf{SU}(n) \times \mathbf{U}(1) \cong \mathbf{U}(n).$$

Exercise 2.11. Find a proof of Proposition 2.31.

Exercise 2.12. Let $(G, *)$ and (H, \cdot) be two Lie groups. Prove that the product manifold $G \times H$ has the structure of a Lie group.

The Tangent Space

In this chapter we introduce the notion of the tangent space $T_p M$ of a differentiable manifold M at a point $p \in M$. This is a vector space of the same dimension as M . We start by studying the standard \mathbb{R}^m and show how a tangent vector v at a point $p \in \mathbb{R}^m$ can be interpreted as a first order linear differential operator, annihilating constants, when acting on real-valued functions locally defined around p . Then we generalise to the manifold setting. To explain the notion of the tangent space we give several examples. Here the Lie classical Lie groups, introduced in Chapter 2, play an important role.

Let \mathbb{R}^m be the m -dimensional real vector space with the standard differentiable structure. If p is a point in \mathbb{R}^m and $\gamma : I \rightarrow \mathbb{R}^m$ is a C^1 -curve such that $\gamma(0) = p$ then the **tangent vector**

$$\dot{\gamma}(0) = \lim_{t \rightarrow 0} \frac{\gamma(t) - \gamma(0)}{t}$$

of γ at p is an element of \mathbb{R}^m . Conversely, for an arbitrary element v of \mathbb{R}^m we can easily find a curve $\gamma : I \rightarrow \mathbb{R}^m$ such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. One example is given by

$$\gamma : t \mapsto p + t \cdot v.$$

This shows that the **tangent space** i.e. the set of tangent vectors at the point $p \in \mathbb{R}^m$ can be identified with \mathbb{R}^m .

We will now describe how first order differential operators annihilating constants can be interpreted as tangent vectors. For a point p in \mathbb{R}^m we denote by $\varepsilon(p)$ the set of differentiable real-valued functions defined locally around p . Then it is well known from multivariable analysis that if $v \in \mathbb{R}^m$ and $f \in \varepsilon(p)$ then the **directional derivative** $\partial_v f$ of f at p in the direction of v is given by

$$\partial_v f = \lim_{t \rightarrow 0} \frac{f(p + tv) - f(p)}{t}.$$

Furthermore the operator ∂ has the following properties

$$\partial_v(\lambda \cdot f + \mu \cdot g) = \lambda \cdot \partial_v f + \mu \cdot \partial_v g,$$

$$\begin{aligned}\partial_v(f \cdot g) &= \partial_v f \cdot g(p) + f(p) \cdot \partial_v g, \\ \partial_{(\lambda \cdot v + \mu \cdot w)} f &= \lambda \cdot \partial_v f + \mu \cdot \partial_w f\end{aligned}$$

for all $\lambda, \mu \in \mathbb{R}$, $v, w \in \mathbb{R}^m$ and $f, g \in \varepsilon(p)$.

Motivated by the above, we make the following definition.

Definition 3.1. For a point p in \mathbb{R}^m let $T_p\mathbb{R}^m$ be the set of first order linear differential operators at p annihilating constants i.e. the set of mappings $\alpha : \varepsilon(p) \rightarrow \mathbb{R}$ such that

- (i) $\alpha(\lambda \cdot f + \mu \cdot g) = \lambda \cdot \alpha(f) + \mu \cdot \alpha(g)$,
- (ii) $\alpha(f \cdot g) = \alpha(f) \cdot g(p) + f(p) \cdot \alpha(g)$

for all $\lambda, \mu \in \mathbb{R}$ and $f, g \in \varepsilon(p)$.

The set of differential operators $T_p\mathbb{R}^m$ carries the structure of a real vector space. This is given by the addition $+$ and the multiplication \cdot by real numbers satisfying

$$\begin{aligned}(\alpha + \beta)(f) &= \alpha(f) + \beta(f), \\ (\lambda \cdot \alpha)(f) &= \lambda \cdot \alpha(f)\end{aligned}$$

for all $\alpha, \beta \in T_p\mathbb{R}^m$, $f \in \varepsilon(p)$ and $\lambda \in \mathbb{R}$.

The following result provides us with an important identification between \mathbb{R}^m and the tangent space $T_p\mathbb{R}^m$ as defined above.

Theorem 3.2. For a point p in \mathbb{R}^m the map $\Phi : \mathbb{R}^m \rightarrow T_p\mathbb{R}^m$ defined by $\Phi : v \mapsto \partial_v$ is a linear vector space isomorphism.

PROOF. The linearity of the map $\Phi : \mathbb{R}^m \rightarrow T_p\mathbb{R}^m$ follows directly from the fact that for all $\lambda, \mu \in \mathbb{R}$, $v, w \in \mathbb{R}^m$ and $f \in \varepsilon(p)$

$$\partial_{(\lambda \cdot v + \mu \cdot w)} f = \lambda \cdot \partial_v f + \mu \cdot \partial_w f.$$

Let $v, w \in \mathbb{R}^m$ be such that $v \neq w$. Choose an element $u \in \mathbb{R}^m$ such that $\langle u, v \rangle \neq \langle u, w \rangle$ and define $f : \mathbb{R}^m \rightarrow \mathbb{R}$ by $f(x) = \langle u, x \rangle$. Then $\partial_v f = \langle u, v \rangle \neq \langle u, w \rangle = \partial_w f$ so $\partial_v \neq \partial_w$. This proves that the linear map Φ is injective.

Let α be an arbitrary element of $T_p\mathbb{R}^m$. For $k = 1, \dots, m$ let the real-valued function $\hat{x}_k : \mathbb{R}^m \rightarrow \mathbb{R}$ be the natural projection onto the k -th component given by

$$\hat{x}_k : (x_1, \dots, x_m) \mapsto x_k$$

and put $v_k = \alpha(\hat{x}_k)$. For the constant function $1 : (x_1, \dots, x_m) \mapsto 1$ we have

$$\alpha(1) = \alpha(1 \cdot 1) = 1 \cdot \alpha(1) + 1 \cdot \alpha(1) = 2 \cdot \alpha(1),$$

so $\alpha(1) = 0$. By the linearity of α it follows that $\alpha(c) = 0$ for any constant $c \in \mathbb{R}$. Let $f \in \varepsilon(p)$ and following Lemma 3.3 locally write

$$f(x) = f(p) + \sum_{k=1}^m (\hat{x}_k(x) - p_k) \cdot \psi_k(x),$$

where $\psi_k \in \varepsilon(p)$ with

$$\psi_k(p) = \frac{\partial f}{\partial x_k}(p).$$

We can now apply the differential operator $\alpha \in T_p \mathbb{R}^m$ and yield

$$\begin{aligned} \alpha(f) &= \alpha\left(f(p) + \sum_{k=1}^m (\hat{x}_k - p_k) \cdot \psi_k\right) \\ &= \alpha(f(p)) + \sum_{k=1}^m \alpha(\hat{x}_k - p_k) \cdot \psi_k(p) + \sum_{k=1}^m (\hat{x}_k(p) - p_k) \cdot \alpha(\psi_k) \\ &= \sum_{k=1}^m v_k \frac{\partial f}{\partial x_k}(p) \\ &= \langle v, \text{grad} f_p \rangle \\ &= \partial_v f, \end{aligned}$$

where $v = (v_1, \dots, v_m) \in \mathbb{R}^m$. This means that $\Phi(v) = \partial_v = \alpha$ so the linear map $\Phi : \mathbb{R}^m \rightarrow T_p \mathbb{R}^m$ is surjective and hence a vector space isomorphism. \square

Lemma 3.3. *Let p be a point in \mathbb{R}^m and $f : U \rightarrow \mathbb{R}$ be a differentiable function defined on an open ball around p . Then for each $k = 1, 2, \dots, m$ there exist functions $\psi_k : U \rightarrow \mathbb{R}$ such that for all $x \in U$*

$$f(x) = f(p) + \sum_{k=1}^m (x_k - p_k) \cdot \psi_k(x) \quad \text{and} \quad \psi_k(p) = \frac{\partial f}{\partial x_k}(p).$$

PROOF. It follows from the fundamental theorem of calculus that

$$\begin{aligned} f(x) - f(p) &= \int_0^1 \frac{\partial}{\partial t} (f(p + t(x - p))) dt \\ &= \sum_{k=1}^m (x_k - p_k) \cdot \int_0^1 \frac{\partial f}{\partial x_k} (p + t(x - p)) dt. \end{aligned}$$

The statement then immediately follows by setting

$$\psi_k(x) = \int_0^1 \frac{\partial f}{\partial x_k} (p + t(x - p)) dt.$$

\square

Remark 3.4. Let p be a point in \mathbb{R}^m , $v \in T_p\mathbb{R}^m$ be a tangent vector at p and $f : U \rightarrow \mathbb{R}$ be a C^1 -function defined on an open subset U of \mathbb{R}^m containing p . Let $\gamma : I \rightarrow U$ be a curve such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. The identification given by Theorem 3.2 tells us that v acts on f by

$$v(f) = \partial_v(f) = \langle v, \text{grad} f_p \rangle = df_p(\dot{\gamma}(0)) = \frac{d}{dt}(f \circ \gamma(t))|_{t=0}.$$

This implies that the real number $v(f)$ is independent of the choice of the curve γ as long as $\gamma(0) = p$ and $\dot{\gamma}(0) = v$.

As a direct consequence of Theorem 3.2 we have the following important result.

Corollary 3.5. *Let p be a point in \mathbb{R}^m and $\{e_k \mid k = 1, \dots, m\}$ be a basis for \mathbb{R}^m . Then the set $\{\partial_{e_k} \mid k = 1, \dots, m\}$ is a basis for the tangent space $T_p\mathbb{R}^m$ at p .*

We will now employ the ideas presented above to generalise to the manifold setting. Let M be a differentiable manifold and for a point $p \in M$ let $\varepsilon(p)$ denote the set of differentiable real-valued functions defined on an open neighborhood of p .

Definition 3.6. Let M be a differentiable manifold and p be a point on M . A **tangent vector** X_p at p is a map $X_p : \varepsilon(p) \rightarrow \mathbb{R}$ such that

- (i) $X_p(\lambda \cdot f + \mu \cdot g) = \lambda \cdot X_p(f) + \mu \cdot X_p(g)$,
- (ii) $X_p(f \cdot g) = X_p(f) \cdot g(p) + f(p) \cdot X_p(g)$

for all $\lambda, \mu \in \mathbb{R}$ and $f, g \in \varepsilon(p)$. The set of tangent vectors at p is called the **tangent space** at p and denoted by T_pM .

The tangent space T_pM of M at p has the structure of a real vector space. The addition $+$ and the multiplication \cdot by real numbers are simply given by

$$\begin{aligned} (X_p + Y_p)(f) &= X_p(f) + Y_p(f), \\ (\lambda \cdot X_p)(f) &= \lambda \cdot X_p(f), \end{aligned}$$

for all $X_p, Y_p \in T_pM$, $f \in \varepsilon(p)$ and $\lambda \in \mathbb{R}$.

We have not yet defined the differential of a map between manifolds, see Definition 3.14, but still think that the following remark is appropriate at this point.

Remark 3.7. Let M be an m -dimensional manifold and (U, x) be a chart around $p \in M$. Then the differential $dx_p : T_pM \rightarrow T_{x(p)}\mathbb{R}^m$ is a bijective linear map such that for a given element $X_p \in T_pM$ there

exists a tangent vector v in $T_{x(p)}\mathbb{R}^m \cong \mathbb{R}^m$ such that $dx_p(X_p) = v$. The image $x(U)$ is an open subset of \mathbb{R}^m containing $x(p)$ so we can find a curve $c : I \rightarrow x(U)$ with $c(0) = x(p)$ and $\dot{c}(0) = v$. Then the composition $\gamma = x^{-1} \circ c : I \rightarrow U$ is a curve in M through p since $\gamma(0) = p$. The element $d(x^{-1})_{x(p)}(v)$ of the tangent space $T_p M$ denoted by $\dot{\gamma}(0)$ is called the **tangent** to the curve γ at p . It follows from the relation

$$\dot{\gamma}(0) = d(x^{-1})_{x(p)}(v) = X_p$$

that the tangent space $T_p M$ can be thought of as the set of all tangents to curves through the point p .

If $f : U \rightarrow \mathbb{R}$ is a C^1 -function defined locally on U then it follows from Definition 3.14 that

$$\begin{aligned} X_p(f) &= (dx_p(X_p))(f \circ x^{-1}) \\ &= \frac{d}{dt}(f \circ x^{-1} \circ c(t))|_{t=0} \\ &= \frac{d}{dt}(f \circ \gamma(t))|_{t=0} \end{aligned}$$

It should be noted that the real number $X_p(f)$ is independent of the choice of the chart (U, x) around p and the curve $c : I \rightarrow x(U)$ as long as $\gamma(0) = p$ and $\dot{\gamma}(0) = X_p$.

We can now determine the tangent spaces of some of the explicit differentiable manifolds that we introduced in Chapter 2.

Example 3.8. Let $\gamma : I \rightarrow S^m$ be a curve into the m -dimensional unit sphere in \mathbb{R}^{m+1} with $\gamma(0) = p$ and $\dot{\gamma}(0) = X$. The curve satisfies

$$\langle \gamma(t), \gamma(t) \rangle = 1$$

and differentiation yields

$$\langle \dot{\gamma}(t), \gamma(t) \rangle + \langle \gamma(t), \dot{\gamma}(t) \rangle = 0.$$

This means that $\langle p, X \rangle = 0$ so every tangent vector $X \in T_p S^m$ must be orthogonal to p . On the other hand if $X \neq 0$ satisfies $\langle p, X \rangle = 0$ then $\gamma : \mathbb{R} \rightarrow S^m$ with

$$\gamma : t \mapsto \cos(t|X|) \cdot p + \sin(t|X|) \cdot X/|X|$$

is a curve into S^m with $\gamma(0) = p$ and $\dot{\gamma}(0) = X$. This shows that the tangent space $T_p S^m$ is actually given by

$$T_p S^m = \{X \in \mathbb{R}^{m+1} \mid \langle p, X \rangle = 0\}.$$

The following is a well known result from matrix theory.

Proposition 3.9. *For complex matrices, the standard exponential map $\text{Exp} : \mathbb{C}^{m \times m} \rightarrow \mathbb{C}^{m \times m}$, is given by the convergent power series*

$$\text{Exp} : Z \mapsto \sum_{k=0}^{\infty} \frac{Z^k}{k!}.$$

Then for two elements $Z, W \in \mathbb{C}^{m \times m}$ we have the following

- (i) $\det(\text{Exp}(Z)) = \exp(\text{trace} Z)$,
- (ii) $\text{Exp}(Z^t) = \overline{\text{Exp}(Z)}^t$,
- (iii) $\text{Exp}(\bar{Z}) = \overline{\text{Exp}(Z)}$,
- (iv) if $ZW = WZ$ then $\text{Exp}(Z + W) = \text{Exp}(Z)\text{Exp}(W)$.

PROOF. See Exercise 3.2 □

The real general linear group $\mathbf{GL}_m(\mathbb{R})$ is an open subset of $\mathbb{R}^{m \times m}$ so its tangent space $T_p \mathbf{GL}_m(\mathbb{R})$ at any point p is simply $\mathbb{R}^{m \times m}$. The tangent space $T_e \mathbf{SL}_m(\mathbb{R})$ of the special linear group $\mathbf{SL}_m(\mathbb{R})$ at the neutral element e can be determined as follows.

Example 3.10. If X is a matrix in $\mathbb{R}^{m \times m}$ with $\text{trace } X = 0$ then we define a curve $A : \mathbb{R} \rightarrow \mathbb{R}^{m \times m}$ by

$$A : s \mapsto \text{Exp}(sX).$$

Then $A(0) = e$, $\dot{A}(0) = X$ and

$$\det(A(s)) = \det(\text{Exp}(sX)) = \exp(\text{trace}(sX)) = \exp(0) = 1.$$

This shows that A is a curve into the special linear group $\mathbf{SL}_m(\mathbb{R})$ and that X is an element of the tangent space $T_e \mathbf{SL}_m(\mathbb{R})$ of $\mathbf{SL}_m(\mathbb{R})$ at the neutral element e . Hence the linear space

$$\{X \in \mathbb{R}^{m \times m} \mid \text{trace } X = 0\}$$

of dimension $m^2 - 1$ is contained in the tangent space $T_e \mathbf{SL}_m(\mathbb{R})$.

On the other hand, the curve given by $s \mapsto \text{Exp}(se) = \exp(s)e$ is not contained in $\mathbf{SL}_m(\mathbb{R})$ so the dimension of $T_e \mathbf{SL}_m(\mathbb{R})$ is at most $m^2 - 1$. This shows that

$$T_e \mathbf{SL}_m(\mathbb{R}) = \{X \in \mathbb{R}^{m \times m} \mid \text{trace } X = 0\}.$$

Next we determine the tangent space of the orthogonal group at the neutral element.

Example 3.11. Let $\gamma : I \rightarrow \mathbf{O}(m)$ be a curve into the orthogonal group $\mathbf{O}(m)$ such that $\gamma(0) = e$. Then $\gamma(s)^t \cdot \gamma(s) = e$ for all $s \in I$ and differentiation gives

$$(\dot{\gamma}(s)^t \cdot \gamma(s) + \gamma(s)^t \cdot \dot{\gamma}(s))|_{s=0} = 0$$

or equivalently $\dot{\gamma}(0)^t + \dot{\gamma}(0) = 0$. This means that each tangent vector of $\mathbf{O}(m)$ at e is a skew-symmetric matrix.

On the other hand, for an arbitrary real skew-symmetric matrix X define the curve $A : \mathbb{R} \rightarrow \mathbb{R}^{m \times m}$ by $A : s \mapsto \text{Exp}(sX)$. Then

$$\begin{aligned} A(s)^t A(s) &= \text{Exp}(sX)^t \text{Exp}(sX) \\ &= \text{Exp}(sX^t) \text{Exp}(sX) \\ &= \text{Exp}(s(X^t + X)) \\ &= \text{Exp}(0) \\ &= e. \end{aligned}$$

This shows that A is a curve on the orthogonal group, $A(0) = e$ and $\dot{A}(0) = X$ so X is an element of $T_e \mathbf{O}(m)$. Hence

$$T_e \mathbf{O}(m) = \{X \in \mathbb{R}^{m \times m} \mid X^t + X = 0\}.$$

The dimension of $T_e \mathbf{O}(m)$ is therefore $m(m-1)/2$. The orthogonal group $\mathbf{O}(m)$ is diffeomorphic to $\mathbf{SO}(m) \times \{\pm 1\}$ so $\dim(\mathbf{SO}(m)) = \dim(\mathbf{O}(m))$ hence

$$T_e \mathbf{SO}(m) = T_e \mathbf{O}(m) = \{X \in \mathbb{R}^{m \times m} \mid X^t + X = 0\}.$$

To summarize, we have now proven the following result.

Proposition 3.12. *Let e be the neutral element of the classical real Lie groups $\mathbf{GL}_m(\mathbb{R})$, $\mathbf{SL}_m(\mathbb{R})$, $\mathbf{O}(m)$, $\mathbf{SO}(m)$. Then their tangent spaces at e are given by*

$$\begin{aligned} T_e \mathbf{GL}_m(\mathbb{R}) &= \mathbb{R}^{m \times m}, \\ T_e \mathbf{SL}_m(\mathbb{R}) &= \{X \in \mathbb{R}^{m \times m} \mid \text{trace} X = 0\}, \\ T_e \mathbf{O}(m) &= \{X \in \mathbb{R}^{m \times m} \mid X^t + X = 0\}, \\ T_e \mathbf{SO}(m) &= T_e \mathbf{O}(m) \cap T_e \mathbf{SL}_m(\mathbb{R}) = T_e \mathbf{O}(m). \end{aligned}$$

For the classical complex Lie groups similar methods can be used to prove the following.

Proposition 3.13. *Let e be the neutral element of the classical complex Lie groups $\mathbf{GL}_m(\mathbb{C})$, $\mathbf{SL}_m(\mathbb{C})$, $\mathbf{U}(m)$, and $\mathbf{SU}(m)$. Then their tangent spaces at e are given by*

$$\begin{aligned} T_e \mathbf{GL}_m(\mathbb{C}) &= \mathbb{C}^{m \times m}, \\ T_e \mathbf{SL}_m(\mathbb{C}) &= \{Z \in \mathbb{C}^{m \times m} \mid \text{trace} Z = 0\}, \\ T_e \mathbf{U}(m) &= \{Z \in \mathbb{C}^{m \times m} \mid \bar{Z}^t + Z = 0\}, \\ T_e \mathbf{SU}(m) &= T_e \mathbf{U}(m) \cap T_e \mathbf{SL}_m(\mathbb{C}). \end{aligned}$$

PROOF. See Exercise 3.4 □

We now introduce the important notion of the differential of a map between manifolds. This will play an important role in what follows.

Definition 3.14. Let $\phi : M \rightarrow N$ be a differentiable map between differentiable manifolds. Then the **differential** $d\phi_p$ of ϕ at a point p in M is the map $d\phi_p : T_p M \rightarrow T_{\phi(p)} N$ such that for all $X_p \in T_p M$ and $f \in \varepsilon(\phi(p))$ we have

$$(d\phi_p(X_p))(f) = X_p(f \circ \phi).$$

Remark 3.15. Let M and N be differentiable manifolds, $p \in M$ and $\phi : M \rightarrow N$ be a differentiable map. Further let $\gamma : I \rightarrow M$ be a curve on M such that $\gamma(0) = p$ and $\dot{\gamma}(0) = X_p$. Let $c : I \rightarrow N$ be the curve $c = \phi \circ \gamma$ in N with $c(0) = \phi(p)$ and put $Y_{\phi(p)} = \dot{c}(0)$. Then it is an immediate consequence of Definition 3.14 that for each function $f \in \varepsilon(\phi(p))$ defined locally around $\phi(p)$ we have

$$\begin{aligned} (d\phi_p(X_p))(f) &= X_p(f \circ \phi) \\ &= \frac{d}{dt}(f \circ \phi \circ \gamma(t))|_{t=0} \\ &= \frac{d}{dt}(f \circ c(t))|_{t=0} \\ &= Y_{\phi(p)}(f). \end{aligned}$$

Hence $d\phi_p(X_p) = Y_{\phi(p)}$ or equivalently $d\phi_p(\dot{\gamma}(0)) = \dot{c}(0)$. This result should be compared with Remark 2.11.

The following result describes the most important properties of the differential, in particular, the so called chain rule.

Proposition 3.16. Let $\phi : M_1 \rightarrow M_2$ and $\psi : M_2 \rightarrow M_3$ be differentiable maps between differentiable manifolds. Then for each point $p \in M_1$ we have

- (i) the map $d\phi_p : T_p M_1 \rightarrow T_{\phi(p)} M_2$ is linear,
- (ii) if $id_{M_1} : M_1 \rightarrow M_1$ is the identity map, then $d(id_{M_1})_p = id_{T_p M_1}$,
- (iii) $d(\psi \circ \phi)_p = d\psi_{\phi(p)} \circ d\phi_p$.

PROOF. The statement (i) follows immediately from the fact that for $\lambda, \mu \in \mathbb{R}$ and $X_p, Y_p \in T_p M$ we have

$$\begin{aligned} d\phi_p(\lambda \cdot X_p + \mu \cdot Y_p)(f) &= (\lambda \cdot X_p + \mu \cdot Y_p)(f \circ \phi) \\ &= \lambda \cdot X_p(f \circ \phi) + \mu \cdot Y_p(f \circ \phi) \\ &= \lambda \cdot d\phi_p(X_p)(f) + \mu \cdot d\phi_p(Y_p)(f). \end{aligned}$$

The statement (ii) is obvious. The statement (iii) is called the **chain rule**. If $X_p \in T_p M_1$ and $f \in \varepsilon(\psi \circ \phi(p))$, then

$$\begin{aligned} (d\psi_{\phi(p)} \circ d\phi_p)(X_p)(f) &= (d\psi_{\phi(p)}(d\phi_p(X_p)))(f) \\ &= (d\phi_p(X_p))(f \circ \psi) \\ &= X_p(f \circ \psi \circ \phi) \\ &= (d(\psi \circ \phi)_p(X_p))(f). \end{aligned}$$

□

As an immediate consequence of Proposition 3.16 we have the following interesting result.

Corollary 3.17. *Let $\phi : M \rightarrow N$ be a diffeomorphism with the inverse $\psi = \phi^{-1} : N \rightarrow M$. Then the differential $d\phi_p : T_p M \rightarrow T_{\phi(p)} N$ at p is bijective and $(d\phi_p)^{-1} = d\psi_{\phi(p)}$.*

PROOF. The statement follows directly from the relations

$$\begin{aligned} d\psi_{\phi(p)} \circ d\phi_p &= d(\psi \circ \phi)_p = d(\text{id}_M)_p = \text{id}_{T_p M}, \\ d\phi_p \circ d\psi_{\phi(p)} &= d(\phi \circ \psi)_{\phi(p)} = d(\text{id}_N)_{\phi(p)} = \text{id}_{T_{\phi(p)} N}. \end{aligned}$$

□

We are now ready to prove the following important result. This is of course a direct generalisation of the corresponding result in the classical theory for surfaces in \mathbb{R}^3 .

Theorem 3.18. *Let M^m be an m -dimensional differentiable manifold and p be a point in M . Then the tangent space $T_p M$ at p is an m -dimensional real vector space.*

PROOF. Let (U, x) be a chart on M . Then the linear map $dx_p : T_p M \rightarrow T_{x(p)} \mathbb{R}^m$ is a vector space isomorphism. The statement now follows from Theorem 3.2 and Corollary 3.17. □

Proposition 3.19. *Let M^m be a differentiable manifold, (U, x) be a chart on M and $\{e_k \mid k = 1, \dots, m\}$ be the canonical basis for \mathbb{R}^m . For an arbitrary point p in U we define the differential operator $(\frac{\partial}{\partial x_k})_p$ in $T_p M$ by*

$$\left(\frac{\partial}{\partial x_k}\right)_p : f \mapsto \frac{\partial f}{\partial x_k}(p) = \partial_{e_k}(f \circ x^{-1})(x(p)).$$

Then the set

$$\left\{\left(\frac{\partial}{\partial x_k}\right)_p \mid k = 1, 2, \dots, m\right\}$$

is a basis for the tangent space $T_p M$ of M at p .

PROOF. The chart $x : U \rightarrow x(U)$ is a diffeomorphism and the differential $(dx^{-1})_{x(p)} : T_{x(p)}\mathbb{R}^m \rightarrow T_pM$ of the inverse $x^{-1} : x(U) \rightarrow U$ satisfies

$$\begin{aligned} (dx^{-1})_{x(p)}(\partial_{e_k})(f) &= \partial_{e_k}(f \circ x^{-1})(x(p)) \\ &= \left(\frac{\partial}{\partial x_k}\right)_p(f) \end{aligned}$$

for all $f \in \varepsilon(p)$. The statement is then a direct consequence of Corollary 3.5. \square

The rest of this chapter is devoted to the introduction of special types of differentiable maps: immersions, embeddings and submersions.

Definition 3.20. For positive integers m, n with $m \leq n$, a differentiable map $\phi : M^m \rightarrow N^n$ between manifolds is said to be an **immersion** if for each $p \in M$ the differential $d\phi_p : T_pM \rightarrow T_{\phi(p)}N$ is injective. An **embedding** is an immersion $\phi : M \rightarrow N$ which is a homeomorphism onto its image $\phi(M)$.

Example 3.21. For positive integers m, n with $m < n$ we have the inclusion map $\phi : \mathbb{R}^{m+1} \rightarrow \mathbb{R}^{n+1}$ given by

$$\phi : (x_1, \dots, x_{m+1}) \mapsto (x_1, \dots, x_{m+1}, 0, \dots, 0).$$

The differential $d\phi_x$ at x is injective since $d\phi_x(v) = (v, 0)$. The map ϕ is obviously a homeomorphism onto its image $\phi(\mathbb{R}^{m+1})$ hence an embedding. It is easily seen that even the restriction $\phi|_{S^m} : S^m \rightarrow S^n$ of ϕ to the m -dimensional unit sphere S^m in \mathbb{R}^{m+1} is an embedding.

Definition 3.22. Let M be an m -dimensional differentiable manifold and U be an open subset of \mathbb{R}^m . An immersion $\phi : U \rightarrow M$ is called a **local parametrisation** of M . If the immersion ϕ is surjective it is said to be a **global parametrisation**.

Remark 3.23. If M is a differentiable manifold and (U, x) is a chart on M , then the inverse $x^{-1} : x(U) \rightarrow U$ of x is a global parametrisation of the open subset U of M .

Example 3.24. Let S^1 be the unit circle in the complex plane \mathbb{C} . For a non-zero integer $k \in \mathbb{Z}$ define $\phi_k : S^1 \rightarrow \mathbb{C}$ by $\phi_k : z \mapsto z^k$. For a point $w \in S^1$ let $\gamma_w : \mathbb{R} \rightarrow S^1$ be the curve with $\gamma_w : t \mapsto we^{it}$. Then $\gamma_w(0) = w$ and $\dot{\gamma}_w(0) = iw$. For the differential of ϕ_k we have

$$(d\phi_k)_w(\dot{\gamma}_w(0)) = \frac{d}{dt}(\phi_k \circ \gamma_w(t))|_{t=0} = \frac{d}{dt}(w^k e^{ikt})|_{t=0} = kiw^k \neq 0.$$

This shows that the differential $(d\phi_k)_w : T_w S^1 \cong \mathbb{R} \rightarrow T_w \mathbb{C} \cong \mathbb{R}^2$ is injective, so the map ϕ_k is an immersion. It is easily seen that ϕ_k is an embedding if and only if $k = \pm 1$.

Example 3.25. Let $q \in S^3$ be a quaternion of unit length and $\phi_q : S^1 \rightarrow S^3$ be the map defined by $\phi_q : z \mapsto qz$. For $w \in S^1$ let $\gamma_w : \mathbb{R} \rightarrow S^1$ be the curve given by $\gamma_w(t) = we^{it}$. Then $\gamma_w(0) = w$, $\dot{\gamma}_w(0) = iw$ and $\phi_q(\gamma_w(t)) = qwe^{it}$. By differentiating we yield

$$d\phi_q(\dot{\gamma}_w(0)) = \frac{d}{dt}(\phi_q(\gamma_w(t)))|_{t=0} = \frac{d}{dt}(qwe^{it})|_{t=0} = qiw.$$

Then $|d\phi_q(\dot{\gamma}_w(0))| = |qwi| = |q||w| = 1 \neq 0$ implies that the differential $d\phi_q$ is injective. It is easily checked that the immersion ϕ_q is an embedding.

In the next example we construct an interesting embedding of the real projective space $\mathbb{R}P^m$ into the vector space $\text{Sym}(\mathbb{R}^{m+1})$ of the real symmetric $(m+1) \times (m+1)$ matrices.

Example 3.26. Let S^m be the m -dimensional unit sphere in \mathbb{R}^{m+1} . For a point $p \in S^m$ let

$$l_p = \{\lambda p \in \mathbb{R}^{m+1} \mid \lambda \in \mathbb{R}\}$$

be the line through the origin generated by p and $\rho_p : \mathbb{R}^{m+1} \rightarrow \mathbb{R}^{m+1}$ be the reflection about the line l_p . Then ρ_p is an element of $\text{End}(\mathbb{R}^{m+1})$ i.e. the set of linear endomorphisms of \mathbb{R}^{m+1} which can be identified with $\mathbb{R}^{(m+1) \times (m+1)}$. It is easily checked that the reflection about the line l_p is given by

$$\rho_p : q \mapsto 2\langle q, p \rangle p - q.$$

It then follows from the equations

$$\rho_p(q) = 2\langle q, p \rangle p - q = 2p\langle p, q \rangle - q = (2pp^t - e)q$$

that the symmetric matrix in $\mathbb{R}^{(m+1) \times (m+1)}$ corresponding to ρ_p is just

$$(2pp^t - e).$$

We will now show that the map $\phi : S^m \rightarrow \text{Sym}(\mathbb{R}^{m+1})$ given by

$$\phi : p \mapsto \rho_p$$

is an immersion. Let p be an arbitrary point on S^m and $\alpha, \beta : I \rightarrow S^m$ be two curves meeting at p , that is $\alpha(0) = p = \beta(0)$, with $X = \dot{\alpha}(0)$ and $Y = \dot{\beta}(0)$. For $\gamma \in \{\alpha, \beta\}$ we have

$$\phi \circ \gamma : t \mapsto (q \mapsto 2\langle q, \gamma(t) \rangle \gamma(t) - q)$$

so

$$\begin{aligned}(d\phi)_p(\dot{\gamma}(0)) &= \frac{d}{dt}(\phi \circ \gamma(t))|_{t=0} \\ &= (q \mapsto 2\langle q, \dot{\gamma}(0) \rangle \gamma(0) + 2\langle q, \gamma(0) \rangle \dot{\gamma}(0)).\end{aligned}$$

This means that

$$d\phi_p(X) = (q \mapsto 2\langle q, X \rangle p + 2\langle q, p \rangle X)$$

and

$$d\phi_p(Y) = (q \mapsto 2\langle q, Y \rangle p + 2\langle q, p \rangle Y).$$

If $X \neq Y$ then $d\phi_p(X) \neq d\phi_p(Y)$ so the differential $d\phi_p$ is injective, hence the map $\phi : S^m \rightarrow \text{Sym}(\mathbb{R}^{m+1})$ is an immersion.

If two points $p, q \in S^m$ are linearly independent, then the lines l_p and l_q are different. But these are just the eigenspaces of ρ_p and ρ_q with the eigenvalue $+1$, respectively. This shows that the linear endomorphisms ρ_p, ρ_q of \mathbb{R}^{m+1} are different in this case.

On the other hand, if p and q are parallel then $p = \pm q$ hence $\rho_p = \rho_q$. This means that the image $\phi(S^m)$ can be identified with the quotient space S^m / \equiv where \equiv is the equivalence relation defined by

$$x \equiv y \text{ if and only if } x = \pm y.$$

This is of course the real projective space $\mathbb{R}P^m$ so the map ϕ induces an embedding $\Phi : \mathbb{R}P^m \rightarrow \text{Sym}(\mathbb{R}^{m+1})$ with

$$\Phi : [p] \rightarrow \rho_p.$$

For each $p \in S^m$ the reflection $\rho_p : \mathbb{R}^{m+1} \rightarrow \mathbb{R}^{m+1}$ about the line l_p satisfies

$$\rho_p^t \cdot \rho_p = e.$$

This shows that the image $\Phi(\mathbb{R}P^m) = \phi(S^m)$ is not only contained in the linear space $\text{Sym}(\mathbb{R}^{m+1})$ but also in the orthogonal group $\mathbf{O}(m+1)$ which we know is a submanifold of $\mathbb{R}^{(m+1) \times (m+1)}$

The following result was proven by Hassler Whitney in his famous paper, *Differentiable Manifolds*, Ann. of Math. **37** (1936), 645-680.

Deep Result 3.27. *For $1 \leq r \leq \infty$ let M be an m -dimensional C^r -manifold. Then there exists a C^r -embedding $\phi : M \rightarrow \mathbb{R}^{2m+1}$ of M into the $(2m+1)$ -dimensional real vector space \mathbb{R}^{2m+1} .*

The classical inverse function theorem generalises to the manifold setting as follows. The reader should compare this with Fact 2.12.

Theorem 3.28 (The Inverse Mapping Theorem). *Let $\phi : M \rightarrow N$ be a differentiable map between manifolds with $\dim M = \dim N$. If p is a point in M such that the differential $d\phi_p : T_p M \rightarrow T_{\phi(p)} N$ at p is bijective then there exist open neighborhoods U_p around p and U_q around $q = \phi(p)$ such that $\psi = \phi|_{U_p} : U_p \rightarrow U_q$ is bijective and the inverse $\psi^{-1} : U_q \rightarrow U_p$ is differentiable.*

PROOF. See Exercise 3.8 □

We will now generalise the classical implicit mapping theorem to manifolds. For this we need the following definition. Compare this with Definition 2.13.

Definition 3.29. Let m, n be positive integers and $\phi : M^m \rightarrow N^n$ be a differentiable map between manifolds. A point $p \in M$ is said to be **regular** for ϕ if the differential

$$d\phi_p : T_p M \rightarrow T_{\phi(p)} N$$

is of full rank, but **critical** otherwise. A point $q \in \phi(M)$ is said to be a **regular value** of ϕ if every point in the pre-image $\phi^{-1}(\{q\})$ of $\{q\}$ is regular.

The reader should compare the following result with Theorem 2.15.

Theorem 3.30 (The Implicit Mapping Theorem). *Let $\phi : M^m \rightarrow N^n$ be a differentiable map between manifolds such that $m > n$. If $q \in \phi(M)$ is a regular value, then the pre-image $\phi^{-1}(\{q\})$ of q is a submanifold of M^m of dimension $m - n$. The tangent space $T_p \phi^{-1}(\{q\})$ of $\phi^{-1}(\{q\})$ at p is the kernel of the differential $d\phi_p$ i.e.*

$$T_p \phi^{-1}(\{q\}) = \{X \in T_p M \mid d\phi_p(X) = 0\}.$$

PROOF. Let (V, y) be a chart on N with $q \in V$ and $y(q) = 0$. For a point $p \in \phi^{-1}(\{q\})$ we choose a chart (U, x) on M such that $p \in U$, $x(p) = 0$ and $\phi(U) \subset V$. Then the differential of the map

$$\psi = y \circ \phi \circ x^{-1}|_{x(U)} : x(U) \rightarrow \mathbb{R}^n$$

at the point 0 is given by

$$d\psi_0 = (dy)_q \circ d\phi_p \circ (dx^{-1})_0 : T_0 \mathbb{R}^m \rightarrow T_0 \mathbb{R}^n.$$

The pairs (U, x) and (V, y) are charts so the differentials $(dy)_q$ and $(dx^{-1})_0$ are bijective. This means that $d\psi_0$ is surjective since $d\phi_p$ is. It then follows from Theorem 2.15 that $x(\phi^{-1}(\{q\}) \cap U)$ is an $(m - n)$ -dimensional submanifold of $x(U)$. Hence $\phi^{-1}(\{q\}) \cap U$ is an $(m - n)$ -dimensional submanifold of U . This is true for each point $p \in \phi^{-1}(\{q\})$

so we have proven that $\phi^{-1}(\{q\})$ is a submanifold of M^m of dimension $(m - n)$.

Let $\gamma : I \rightarrow \phi^{-1}(\{q\})$ be a curve such that $\gamma(0) = p$. Then

$$(d\phi)_p(\dot{\gamma}(0)) = \frac{d}{dt}(\phi \circ \gamma(t))|_{t=0} = \frac{dq}{dt}|_{t=0} = 0.$$

This implies that $T_p\phi^{-1}(\{q\})$ is contained in and has the same dimension as the kernel of $d\phi_p$, so $T_p\phi^{-1}(\{q\}) = \text{Ker } d\phi_p$. \square

We conclude this chapter with a discussion on the important submersions between differentiable manifolds.

Definition 3.31. For positive integers m, n with $m \geq n$ a differentiable map $\phi : M^m \rightarrow N^n$ between two manifolds is said to be a **submersion** if for each $p \in M$ the differential $d\phi_p : T_pM \rightarrow T_{\phi(p)}N$ is surjective.

The reader should compare Definition 3.31 with Definition 3.20.

Example 3.32. If m, n are positive integers such that $m \geq n$ then we have the projection map $\pi : \mathbb{R}^m \rightarrow \mathbb{R}^n$ given by $\pi : (x_1, \dots, x_m) \mapsto (x_1, \dots, x_n)$. Its differential $d\pi_x$ at a point x is surjective since

$$d\pi_x(v_1, \dots, v_m) = (v_1, \dots, v_n).$$

This means that the projection is a submersion.

The following important example provides us with a submersion between spheres.

Example 3.33. Let S^3 and S^2 be the unit spheres in \mathbb{C}^2 and $\mathbb{C} \times \mathbb{R} \cong \mathbb{R}^3$, respectively. The **Hopf map** $\phi : S^3 \rightarrow S^2$ is given by

$$\phi : (z, w) \mapsto (2z\bar{w}, |z|^2 - |w|^2).$$

For $p = (z, w) \in S^3$ the **Hopf circle** C_p through p is given by

$$C_p = \{e^{i\theta}(z, w) \mid \theta \in \mathbb{R}\}.$$

The following shows that the Hopf map is constant along each Hopf circle

$$\begin{aligned} \phi(e^{i\theta}(z, w)) &= (2e^{i\theta}ze^{-i\theta}\bar{w}, |e^{i\theta}z|^2 - |e^{i\theta}w|^2) \\ &= (2z\bar{w}, |z|^2 - |w|^2) \\ &= \phi((z, w)). \end{aligned}$$

The map ϕ and its differential $d\phi_p : T_pS^3 \rightarrow T_{\phi(p)}S^2$ are surjective for each $p \in S^3$. This implies that each point $q \in S^2$ is a regular value and

the fibres of ϕ are 1-dimensional submanifolds of S^3 . They are actually the great circles given by

$$\phi^{-1}(\{(2z\bar{w}, |z|^2 - |w|^2)\}) = \{e^{i\theta}(z, w) \mid \theta \in \mathbb{R}\}.$$

This means that the 3-dimensional sphere S^3 is a disjoint union of great circles

$$S^3 = \bigcup_{q \in S^2} \phi^{-1}(\{q\}).$$

Exercises

Exercise 3.1. Let p be an arbitrary point of the unit sphere S^{2n+1} in $\mathbb{C}^{n+1} \cong \mathbb{R}^{2n+2}$. Determine the tangent space $T_p S^{2n+1}$ and show that this contains an n -dimensional complex vector subspace of \mathbb{C}^{n+1} .

Exercise 3.2. Use your local library to find a proof of Proposition 3.9.

Exercise 3.3. Prove that the matrices

$$X_1 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad X_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad X_3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

form a basis for the tangent space $T_e \mathbf{SL}_2(\mathbb{R})$ of the real special linear group $\mathbf{SL}_2(\mathbb{R})$ at the neutral element e . For each $k = 1, 2, 3$ find an explicit formula for the curve $\gamma_k : \mathbb{R} \rightarrow \mathbf{SL}_2(\mathbb{R})$ given by

$$\gamma_k : s \mapsto \text{Exp}(sX_k).$$

Exercise 3.4. Find a proof of Proposition 3.13.

Exercise 3.5. Prove that the matrices

$$Z_1 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad Z_2 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad Z_3 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix},$$

form a basis for the tangent space $T_e \mathbf{SU}(2)$ of the special unitary group $\mathbf{SU}(2)$ at the neutral element e . For each $k = 1, 2, 3$ find an explicit formula for the curve $\gamma_k : \mathbb{R} \rightarrow \mathbf{SU}(2)$ given by

$$\gamma_k : s \mapsto \text{Exp}(sZ_k).$$

Exercise 3.6. For each non-negative integer k define $\phi_k : \mathbb{C} \rightarrow \mathbb{C}$ and $\psi_k : \mathbb{C}^* \rightarrow \mathbb{C}$ by $\phi_k, \psi_k : z \mapsto z^k$. For which such k are ϕ_k, ψ_k immersions, embeddings or submersions?

Exercise 3.7. Prove that the differentiable map $\phi : \mathbb{R}^m \rightarrow \mathbb{C}^m$ given by

$$\phi : (x_1, \dots, x_m) \mapsto (e^{ix_1}, \dots, e^{ix_m})$$

is a parametrisation of the m -dimensional torus T^m in \mathbb{C}^m .

Exercise 3.8. Find a proof of Theorem 3.28.

Exercise 3.9. Prove that the Hopf-map $\phi : S^3 \rightarrow S^2$ with $\phi : (x, y) \mapsto (2x\bar{y}, |x|^2 - |y|^2)$ is a submersion.

The Tangent Bundle

In this chapter we introduce the tangent bundle TM of a differentiable manifold M . Intuitively, this is the object that we get by glueing at each point p on M the corresponding tangent space T_pM . The differentiable structure on M induces a natural differentiable structure on the tangent bundle TM turning it into a differentiable manifold. To explain the important notion of the tangent bundle we investigate several examples. The classical Lie groups will here play a particular important role.

We have already seen that for a point $p \in \mathbb{R}^m$ the **tangent space** $T_p\mathbb{R}^m$ can be identified with the m -dimensional vector space \mathbb{R}^m . This means that if we at each point $p \in \mathbb{R}^m$ glue the tangent space $T_p\mathbb{R}^m$ to \mathbb{R}^m we obtain the so called **tangent bundle** of \mathbb{R}^m

$$T\mathbb{R}^m = \{(p, v) \mid p \in \mathbb{R}^m \text{ and } v \in T_p\mathbb{R}^m\}.$$

For this we have the **natural projection** $\pi : T\mathbb{R}^m \rightarrow \mathbb{R}^m$ defined by

$$\pi : (p, v) \mapsto p$$

and for each point $p \in M$ the fibre $\pi^{-1}(\{p\})$ over p is precisely the tangent space $T_p\mathbb{R}^m$ at p .

Remark 4.1. Classically, a **vector field** X on \mathbb{R}^m is a differentiable map $X : \mathbb{R}^m \rightarrow \mathbb{R}^m$ but we would like to view it as a map $X : \mathbb{R}^m \rightarrow T\mathbb{R}^m$ into the tangent bundle and write

$$X : p \mapsto (p, X_p).$$

Following Proposition 3.19 two vector fields $X, Y : \mathbb{R}^m \rightarrow T\mathbb{R}^m$ can be written as

$$X = \sum_{k=1}^m a_k \frac{\partial}{\partial x_k} \quad \text{and} \quad Y = \sum_{k=1}^m b_k \frac{\partial}{\partial x_k},$$

where $a_k, b_k : \mathbb{R}^m \rightarrow \mathbb{R}$ are differentiable functions defined on \mathbb{R}^m . If $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is another such function the **commutator** $[X, Y]$ acts on f as follows.

$$[X, Y](f) = X(Y(f)) - Y(X(f))$$

$$\begin{aligned}
&= \sum_{k,l=1}^m \left(a_k \frac{\partial}{\partial x_k} \left(b_l \frac{\partial}{\partial x_l} \right) - b_k \frac{\partial}{\partial x_k} \left(a_l \frac{\partial}{\partial x_l} \right) \right) (f) \\
&= \sum_{k,l=1}^m \left(a_k \frac{\partial b_l}{\partial x_k} \frac{\partial}{\partial x_l} + a_k b_l \frac{\partial^2}{\partial x_k \partial x_l} \right. \\
&\quad \left. - b_k \frac{\partial a_l}{\partial x_k} \frac{\partial}{\partial x_l} - b_k a_l \frac{\partial^2}{\partial x_k \partial x_l} \right) (f) \\
&= \sum_{l=1}^m \left\{ \sum_{k=1}^m \left(a_k \frac{\partial b_l}{\partial x_k} - b_k \frac{\partial a_l}{\partial x_k} \right) \right\} \frac{\partial}{\partial x_l} (f).
\end{aligned}$$

This shows that the commutator $[X, Y]$ is a differentiable vector field on \mathbb{R}^m .

We will now generalise to the manifold setting. First we introduce the following notion of a topological vector bundle.

Definition 4.2. Let E and M be topological manifolds and $\pi : E \rightarrow M$ be a continuous surjective map. The triple (E, M, π) is called an n -dimensional **topological vector bundle** over M if

- (i) for each $p \in M$ the **fibre** $E_p = \pi^{-1}(\{p\})$ is an n -dimensional vector space,
- (ii) for each point $p \in M$ there exists a **bundle chart** $(\pi^{-1}(U), \psi)$ consisting of the pre-image $\pi^{-1}(U)$ of an open neighbourhood U of p and a homeomorphism $\psi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^n$ such that for all $q \in U$ the map $\psi_q = \psi|_{E_q} : E_q \rightarrow \{q\} \times \mathbb{R}^n$ is a vector space isomorphism.

A **bundle atlas** for (E, M, π) is a collection

$$\mathcal{B} = \{(\pi^{-1}(U_\alpha), \psi_\alpha) \mid \alpha \in \mathcal{I}\}$$

of bundle charts such that $M = \cup_\alpha U_\alpha$ and for all $\alpha, \beta \in \mathcal{I}$ there exists a map $A_{\alpha,\beta} : U_\alpha \cap U_\beta \rightarrow \mathbf{GL}_n(\mathbb{R})$ such that the corresponding continuous map

$$\psi_\beta \circ \psi_\alpha^{-1}|_{(U_\alpha \cap U_\beta) \times \mathbb{R}^n} : (U_\alpha \cap U_\beta) \times \mathbb{R}^n \rightarrow (U_\alpha \cap U_\beta) \times \mathbb{R}^n$$

is given by

$$(p, v) \mapsto (p, (A_{\alpha,\beta}(p))(v)).$$

The elements of $\{A_{\alpha,\beta} \mid \alpha, \beta \in \mathcal{I}\}$ are called the **transition maps** of the bundle atlas \mathcal{B} .

Definition 4.3. Let (E, M, π) be an n -dimensional topological vector bundle over M . A continuous map $\sigma : M \rightarrow E$ is called a **section** of the bundle (E, M, π) if $\pi \circ \sigma(p) = p$ for each $p \in M$.

Definition 4.4. A topological vector bundle (E, M, π) over M , of dimension n , is said to be **trivial** if there exists a global bundle chart $\psi : E \rightarrow M \times \mathbb{R}^n$.

Example 4.5. Let M be the one dimensional circle S^1 , E be the two dimensional cylinder $E = S^1 \times \mathbb{R}^1$ and $\pi : E \rightarrow M$ be the projection map given by $\pi : (z, t) \mapsto z$. Then (E, M, π) is a trivial line bundle i.e. a trivial one dimensional vector bundle over the circle. This because the identity map $\psi : S^1 \times \mathbb{R}^1 \rightarrow S^1 \times \mathbb{R}^1$ with $\psi : (z, t) \rightarrow (z, t)$ is a global bundle chart.

Example 4.6. For a positive integer n and a topological manifold M we have the n -dimensional **trivial vector bundle** $(M \times \mathbb{R}^n, M, \pi)$ over M , where $\pi : M \times \mathbb{R}^n \rightarrow M$ is the projection map with $\pi : (p, v) \mapsto p$. The bundle is trivial since the identity map $\psi : M \times \mathbb{R}^n \rightarrow M \times \mathbb{R}^n$ is a global bundle chart.

Example 4.7. Let M be the circle S^1 in \mathbb{R}^4 parametrised by $\gamma : \mathbb{R} \rightarrow \mathbb{R}^4$ with

$$\gamma : s \mapsto (\cos s, \sin s, 0, 0).$$

Further let E be the well known **Möbius band** in \mathbb{R}^4 parametrised by $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^4$ with

$$\phi : (s, t) \mapsto (\cos s, \sin s, 0, 0) + t \cdot (0, 0, \sin(s/2), \cos(s/2)).$$

Then E is a regular surface and the natural projection $\pi : E \rightarrow M$ given by $\pi : (x, y, z, w) \mapsto (x, y)$ is continuous and surjective. The triple (E, M, π) is a line bundle over the circle S^1 . The Möbius band is not orientable and hence not homeomorphic to the product $S^1 \times \mathbb{R}$. This shows that the bundle (E, M, π) is not trivial.

Definition 4.8. Let E and M be **differentiable** manifolds and $\pi : E \rightarrow M$ be a **differentiable** map such that (E, M, π) is an n -dimensional topological vector bundle. A bundle atlas \mathcal{B} for (E, M, π) is said to be differentiable if the corresponding transition maps are differentiable. A **differentiable vector bundle** is a topological vector bundle together with a maximal differentiable bundle atlas. By $C^\infty(E)$ we denote the set of all smooth sections of (E, M, π) .

From now on we will assume, when not stating otherwise, that all our vector bundles are smooth.

Definition 4.9. Let (E, M, π) be a vector bundle over a manifold M . Then we define the operations $+$ and \cdot on the set $C^\infty(E)$ of smooth sections of (E, M, π) by

- (i) $(v + w)_p = v_p + w_p$,
- (ii) $(f \cdot v)_p = f(p) \cdot v_p$

for all $p \in M$, $v, w \in C^\infty(E)$ and $f \in C^\infty(M)$. If U is an open subset of M then a set $\{v_1, \dots, v_n\}$ of smooth sections $v_1, \dots, v_n : U \rightarrow E$ on U is called a **local frame** for E if for each $p \in U$ the set $\{(v_1)_p, \dots, (v_n)_p\}$ is a basis for the vector space E_p .

Remark 4.10. According to Definition 2.19, the set of smooth real-valued functions on M is denoted by $C^\infty(M)$. With the above defined operations on $C^\infty(E)$ it becomes a **module** over $C^\infty(M)$ and in particular a vector space over the real numbers as the constant functions in $C^\infty(M)$.

The following example is the central part of this chapter. Here we construct the tangent bundle of a differentiable manifold.

Example 4.11. Let M^m be a differentiable manifold with maximal atlas $\hat{\mathcal{A}}$. Define the **set** TM by

$$TM = \{(p, v) \mid p \in M \text{ and } v \in T_p M\}$$

and let $\pi : TM \rightarrow M$ be the **projection map** satisfying

$$\pi : (p, v) \mapsto p.$$

Then the **fibre** $\pi^{-1}(\{p\})$ is the m -dimensional **tangent space** $T_p M$. The triple (TM, M, π) is called the **tangent bundle** of M . We will now equip this with the structure of a differentiable vector bundle.

For every chart $x : U \rightarrow \mathbb{R}^m$ from the maximal atlas $\hat{\mathcal{A}}$ of M we define the chart

$$x^* : \pi^{-1}(U) \rightarrow \mathbb{R}^m \times \mathbb{R}^m$$

on the tangent bundle TM by the formula

$$x^* : (p, \sum_{k=1}^m v_k(p) \left(\frac{\partial}{\partial x_k} \right)_p) \mapsto (x(p), (v_1(p), \dots, v_m(p))).$$

Proposition 3.19 shows that the map x^* is well defined. The collection

$$\{(x^*)^{-1}(W) \subset TM \mid (U, x) \in \hat{\mathcal{A}} \text{ and } W \subset x(U) \times \mathbb{R}^m \text{ open}\}$$

is a basis for a topology \mathcal{T}_{TM} on TM and $(\pi^{-1}(U), x^*)$ is a chart on the **topological manifold** (TM, \mathcal{T}_{TM}) of dimension $2m$.

If (U, x) and (V, y) are two charts in $\hat{\mathcal{A}}$ such that $p \in U \cap V$ then the transition map

$$(y^*) \circ (x^*)^{-1} : x^*(\pi^{-1}(U \cap V)) \rightarrow \mathbb{R}^m \times \mathbb{R}^m$$

is given by

$$(a, b) \mapsto (y \circ x^{-1}(a), \sum_{k=1}^m \frac{\partial y_1}{\partial x_k}(x^{-1}(a))b_k, \dots, \sum_{k=1}^m \frac{\partial y_m}{\partial x_k}(x^{-1}(a))b_k),$$

see Exercise 4.1. Since we are assuming that $y \circ x^{-1}$ is differentiable it follows that $(y^*) \circ (x^*)^{-1}$ is also differentiable. This means that

$$\mathcal{A}^* = \{(\pi^{-1}(U), x^*) \mid (U, x) \in \hat{\mathcal{A}}\}$$

is a C^r -atlas on TM so $(TM, \widehat{\mathcal{A}^*})$ is a **differentiable manifold**. The surjective projection map $\pi : TM \rightarrow M$ is clearly differentiable.

For each point $p \in M$ the fibre $\pi^{-1}(\{p\})$ is the tangent space $T_p M$ and hence an m -dimensional vector space. For a chart $x : U \rightarrow \mathbb{R}^m$ in the maximal atlas $\hat{\mathcal{A}}$ of M we define $\bar{x} : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^m$ by

$$\bar{x} : (p, \sum_{k=1}^m v_k(p) \left(\frac{\partial}{\partial x_k} \right)_p) \mapsto (p, (v_1(p), \dots, v_m(p))).$$

The restriction $\bar{x}_p = \bar{x}|_{T_p M} : T_p M \rightarrow \{p\} \times \mathbb{R}^m$ to the tangent space $T_p M$ is given by

$$\bar{x}_p : \sum_{k=1}^m v_k(p) \left(\frac{\partial}{\partial x_k} \right)_p \mapsto (v_1(p), \dots, v_m(p)),$$

so it is clearly a vector space isomorphism. This implies that the map

$$\bar{x} : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^m$$

is a bundle chart. If (U, x) and (V, y) are two charts in $\hat{\mathcal{A}}$ such that $p \in U \cap V$ then the transition map

$$(\bar{y}) \circ (\bar{x})^{-1} : (U \cap V) \times \mathbb{R}^m \rightarrow (U \cap V) \times \mathbb{R}^m$$

is given by

$$(p, b) \mapsto (p, \sum_{k=1}^m \frac{\partial y_1}{\partial x_k}(p) \cdot b_k, \dots, \sum_{k=1}^m \frac{\partial y_m}{\partial x_k}(p) \cdot b_k).$$

It is clear that the matrix

$$\begin{pmatrix} \partial y_1 / \partial x_1(p) & \dots & \partial y_1 / \partial x_m(p) \\ \vdots & \ddots & \vdots \\ \partial y_m / \partial x_1(p) & \dots & \partial y_m / \partial x_m(p) \end{pmatrix}$$

is of full rank so the corresponding linear map $A(p) : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a vector space isomorphism for all $p \in U \cap V$. This shows that

$$\mathcal{B} = \{(\pi^{-1}(U), \bar{x}) \mid (U, x) \in \hat{\mathcal{A}}\}$$

is a bundle atlas turning (TM, M, π) into a topological vector bundle of dimension m . It immediately follows from above that (TM, M, π) together with the maximal bundle atlas $\hat{\mathcal{B}}$ defined by \mathcal{B} is a **differentiable vector bundle**.

Definition 4.12. Let M be a differentiable manifold, then a section $X : M \rightarrow TM$ of the tangent bundle is called a **vector field**. The set of smooth vector fields $X : M \rightarrow TM$ is denoted by $C^\infty(TM)$.

Example 4.13. We have seen earlier that the 3-dimensional sphere S^3 in $\mathbb{H} \cong \mathbb{C}^2 \cong \mathbb{R}^4$ carries a group structure \cdot given by

$$(z, w) \cdot (\alpha, \beta) = (z\alpha - w\bar{\beta}, z\beta + w\bar{\alpha}).$$

This turns (S^3, \cdot) into a Lie group with neutral element $e = (1, 0)$. Put $v_1 = (i, 0)$, $v_2 = (0, 1)$ and $v_3 = (0, i)$ and for $k = 1, 2, 3$ define the curves $\gamma_k : \mathbb{R} \rightarrow S^3$ with

$$\gamma_k : t \mapsto \cos t \cdot (1, 0) + \sin t \cdot v_k.$$

Then $\gamma_k(0) = e$ and $\dot{\gamma}_k(0) = v_k$ so v_1, v_2, v_3 are elements of the tangent space $T_e S^3$ of S^3 at the neutral element e . They are linearly independent and hence generate $T_e S^3$. The group structure on S^3 can be used to extend vectors in $T_e S^3$ to vector fields on S^3 as follows. For $p \in S^3$ let $L_p : S^3 \rightarrow S^3$ be the left translation on S^3 by p given by $L_p : q \mapsto p \cdot q$. Then define the vector fields $X_1, X_2, X_3 \in C^\infty(TS^3)$ by

$$(X_k)_p = (dL_p)_e(v_k) = \frac{d}{dt}(L_p(\gamma_k(t)))|_{t=0}.$$

It is left as an exercise for the reader to show that at an arbitrary point $p = (z, w) \in S^3$ the values of X_k at p are given by

$$\begin{aligned} (X_1)_p &= (z, w) \cdot (i, 0) = (iz, -iw), \\ (X_2)_p &= (z, w) \cdot (0, 1) = (-w, z), \\ (X_3)_p &= (z, w) \cdot (0, i) = (iw, iz). \end{aligned}$$

Our next goal is to introduce the Lie bracket on the set of vector fields $C^\infty(TM)$ on M .

Definition 4.14. Let M be a smooth manifold. For two vector fields $X, Y \in C^\infty(TM)$ we define the **Lie bracket** $[X, Y]_p : C^\infty(M) \rightarrow \mathbb{R}$ of X and Y at $p \in M$ by

$$[X, Y]_p(f) = X_p(Y(f)) - Y_p(X(f)).$$

Remark 4.15. The reader should note that if M is a smooth manifold, $X \in C^\infty(TM)$ and $f \in C^\infty(M)$ then the derivative $X(f)$ is the smooth real-valued function on M given by $X(f) : q \mapsto X_q(f)$ for all $q \in M$.

The next result shows that the Lie bracket $[X, Y]_p$ actually is an element of the tangent space $T_p M$. The reader should compare this with Definition 3.6 and Remark 4.1.

Proposition 4.16. *Let M be a smooth manifold, $X, Y \in C^\infty(TM)$ be vector fields on M , $f, g \in C^\infty(M)$ and $\lambda, \mu \in \mathbb{R}$. Then*

- (i) $[X, Y]_p(\lambda \cdot f + \mu \cdot g) = \lambda \cdot [X, Y]_p(f) + \mu \cdot [X, Y]_p(g)$,
- (ii) $[X, Y]_p(f \cdot g) = [X, Y]_p(f) \cdot g(p) + f(p) \cdot [X, Y]_p(g)$.

PROOF.

$$\begin{aligned}
 & [X, Y]_p(\lambda f + \mu g) \\
 &= X_p(Y(\lambda f + \mu g)) - Y_p(X(\lambda f + \mu g)) \\
 &= \lambda X_p(Y(f)) + \mu X_p(Y(g)) - \lambda Y_p(X(f)) - \mu Y_p(X(g)) \\
 &= \lambda [X, Y]_p(f) + \mu [X, Y]_p(g). \\
 \\
 & [X, Y]_p(f \cdot g) \\
 &= X_p(Y(f \cdot g)) - Y_p(X(f \cdot g)) \\
 &= X_p(f \cdot Y(g) + g \cdot Y(f)) - Y_p(f \cdot X(g) + g \cdot X(f)) \\
 &= X_p(f)Y_p(g) + f(p)X_p(Y(g)) + X_p(g)Y_p(f) + g(p)X_p(Y(f)) \\
 &\quad - Y_p(f)X_p(g) - f(p)Y_p(X(g)) - Y_p(g)X_p(f) - g(p)Y_p(X(f)) \\
 &= f(p)\{X_p(Y(g)) - Y_p(X(g))\} + g(p)\{X_p(Y(f)) - Y_p(X(f))\} \\
 &= f(p)[X, Y]_p(g) + g(p)[X, Y]_p(f).
 \end{aligned}$$

□

Proposition 4.16 implies that if X, Y are smooth vector fields on M then the map $[X, Y] : M \rightarrow TM$ given by $[X, Y] : p \mapsto [X, Y]_p$ is a section of the tangent bundle. In Proposition 4.18 we will prove that this section is smooth. For this we need the following technical lemma.

Lemma 4.17. *Let M^m be a smooth manifold and $X : M \rightarrow TM$ be a section of TM . Then the following conditions are equivalent*

- (i) *the section X is smooth,*
- (ii) *if (U, x) is a chart on M then the functions $a_1, \dots, a_m : U \rightarrow \mathbb{R}$ given by*

$$X|_U = \sum_{k=1}^m a_k \frac{\partial}{\partial x_k},$$

are smooth,

- (iii) *if $f : V \rightarrow \mathbb{R}$ defined on an open subset V of M is smooth, then the function $X(f) : V \rightarrow \mathbb{R}$ with $X(f)(p) = X_p(f)$ is smooth.*

PROOF. This proof is divided into three parts. First we show that (i) implies (ii): The functions

$$a_k = \pi_{m+k} \circ x^* \circ X|_U : U \rightarrow \pi^{-1}(U) \rightarrow x(U) \times \mathbb{R}^m \rightarrow \mathbb{R}$$

are compositions of smooth maps so therefore smooth.

Secondly, we now show that (ii) gives (iii): Let (U, x) be a chart on M such that U is contained in V . By assumption the map

$$X(f|_U) = \sum_{i=1}^m a_i \frac{\partial f}{\partial x_i}$$

is smooth. This is true for each such chart (U, x) so the function $X(f)$ is smooth on V .

Finally we show that (iii) leads to (i): Note that the smoothness of the section X is equivalent to $x^* \circ X|_U : U \rightarrow \mathbb{R}^{2m}$ being smooth for all charts (U, x) on M . On the other hand, this is equivalent to

$$x_k^* = \pi_k \circ x^* \circ X|_U : U \rightarrow \mathbb{R}$$

being smooth for all $k = 1, 2, \dots, 2m$ and all charts (U, x) on M . It is trivial that the coordinate functions $x_k^* = x_k$ for $k = 1, \dots, m$ are smooth. But $x_{m+k}^* = a_k = X(x_k)$ for $k = 1, \dots, m$ hence also smooth by assumption. \square

Proposition 4.18. *Let M be a manifold and $X, Y \in C^\infty(TM)$ be vector fields on M . Then the section $[X, Y] : M \rightarrow TM$ of the tangent bundle given by $[X, Y] : p \mapsto [X, Y]_p$ is smooth.*

PROOF. Let $f : M \rightarrow \mathbb{R}$ be an arbitrary smooth function on M then $[X, Y](f) = X(Y(f)) - Y(X(f))$ is smooth so it follows from Lemma 4.17 that the section $[X, Y]$ is smooth. \square

For later use we prove the following important result.

Lemma 4.19. *Let M be a smooth manifold and $[\cdot, \cdot]$ be the Lie bracket on the tangent bundle TM . Then*

$$(i) \quad [X, f \cdot Y] = X(f) \cdot Y + f \cdot [X, Y],$$

$$(ii) \quad [f \cdot X, Y] = f \cdot [X, Y] - Y(f) \cdot X$$

for all $X, Y \in C^\infty(TM)$ and $f \in C^\infty(M)$.

PROOF. If $g \in C^\infty(M)$, then

$$\begin{aligned} [X, f \cdot Y](g) &= X(f \cdot Y(g)) - f \cdot Y(X(g)) \\ &= X(f) \cdot Y(g) + f \cdot X(Y(g)) - f \cdot Y(X(g)) \\ &= (X(f) \cdot Y + f \cdot [X, Y])(g). \end{aligned}$$

This proves the first statement and the second follows from the skew-symmetry of the Lie bracket. \square

We now define the notion of a Lie algebra. This is a fundamental concept in differential geometry.

Definition 4.20. A real vector space $(V, +, \cdot)$ equipped with an operation $[\cdot] : V \times V \rightarrow V$ is said to be a **Lie algebra** if the following relations hold

- (i) $[\lambda X + \mu Y, Z] = \lambda[X, Z] + \mu[Y, Z]$,
- (ii) $[X, Y] = -[Y, X]$,
- (iii) $[X, [Y, Z]] + [Z, [X, Y]] + [Y, [Z, X]] = 0$

for all $X, Y, Z \in V$ and $\lambda, \mu \in \mathbb{R}$. The equation (iii) is called the **Jacobi identity**.

Example 4.21. Let \mathbb{R}^3 be the standard 3-dimensional real vector space generated by $X = (1, 0, 0)$, $Y = (0, 1, 0)$ and $Z = (0, 0, 1)$. Let \times be the standard cross product on \mathbb{R}^3 and define the skew-symmetric bilinear operation $[\cdot] : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by

$$[X, Y] = X \times Y = Z,$$

$$[Z, X] = Z \times X = Y,$$

$$[Y, Z] = Y \times Z = X.$$

This turns \mathbb{R}^3 into a Lie algebra. Compare this with Exercise 4.7.

Theorem 4.22. Let M be a smooth manifold. The vector space $C^\infty(TM)$ of smooth vector fields on M equipped with the Lie bracket $[\cdot] : C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(TM)$ is a Lie algebra.

PROOF. See Exercise 4.4. □

Definition 4.23. If $\phi : M \rightarrow N$ is a **surjective** map between differentiable manifolds, then two vector fields $X \in C^\infty(TM)$ and $\bar{X} \in C^\infty(TN)$ are said to be **ϕ -related** if $d\phi_p(X_p) = \bar{X}_{\phi(p)}$ for all $p \in M$. In this situation we write $d\phi(X) = \bar{X}$.

Example 4.24. Let S^1 be the unit circle in the complex plane and $\phi : S^1 \rightarrow S^1$ be the map given by $\phi(z) = z^2$. Note that this is surjective but not bijective. Further let X be the vector field on S^1 satisfying $X(z) = iz$. Then

$$d\phi_z(X_z) = \frac{d}{d\theta}(\phi(ze^{i\theta}))|_{\theta=0} = \frac{d}{d\theta}((ze^{i\theta})^2)|_{\theta=0} = 2iz^2 = 2X_{\phi(z)}.$$

This shows that the vector field X is ϕ -related to $\bar{X} = 2X$.

Example 4.25. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a surjective C^1 -function and $x, y \in \mathbb{R}$ such that $x \neq y$, $f(x) = f(y)$ and $f'(x) \neq f'(y)$. Further

let $\gamma : \mathbb{R} \rightarrow \mathbb{R}$ be the curve with $\gamma(t) = t$ and define the vector field $X \in C^1(\mathbb{R})$ by $X_t = \dot{\gamma}(t)$. Then for each $t \in \mathbb{R}$ we have

$$df_t(X_t) = (f \circ \gamma(t))' = f'(t).$$

If $\bar{X} \in C^1(\mathbb{R})$ is a vector field which is f -related to X then

$$\bar{X}_{f(x)} = df_x(X_x) = f'(x) \neq f'(y) = df_y(X_y) = \bar{X}_{f(y)}.$$

This contradicts the existence of such a vector field \bar{X} .

Proposition 4.26. *Let $\phi : M \rightarrow N$ be a surjective map between differentiable manifolds and $X, Y \in C^\infty(TM)$, $\bar{X}, \bar{Y} \in C^\infty(TN)$ such that $d\phi(X) = \bar{X}$ and $d\phi(Y) = \bar{Y}$. Then*

$$d\phi([X, Y]) = [\bar{X}, \bar{Y}].$$

PROOF. Let $p \in M$ and $f : N \rightarrow \mathbb{R}$ be a smooth function, then

$$\begin{aligned} d\phi_p([X, Y]_p)(f) &= [X, Y]_p(f \circ \phi) \\ &= X_p(Y(f \circ \phi)) - Y_p(X(f \circ \phi)) \\ &= X_p(d\phi(Y)(f) \circ \phi) - Y_p(d\phi(X)(f) \circ \phi) \\ &= d\phi(X)_{\phi(p)}(d\phi(Y)(f)) - d\phi(Y)_{\phi(p)}(d\phi(X)(f)) \\ &= [\bar{X}, \bar{Y}]_{\phi(p)}(f). \end{aligned}$$

□

Proposition 4.27. *Let M and N be differentiable manifolds and $\phi : M \rightarrow N$ be a diffeomorphism. If $X, Y \in C^\infty(TM)$ are vector fields on M , then $d\phi(X)$ is a vector field on N and the tangent map $d\phi : C^\infty(TM) \rightarrow C^\infty(TN)$ is a Lie algebra homomorphism i.e.*

$$d\phi([X, Y]) = [d\phi(X), d\phi(Y)].$$

PROOF. The fact that ϕ is bijective implies that $d\phi(X)$ is a section of the tangent bundle. That $d\phi(X)$ is smooth follows directly from the fact that

$$d\phi(X)(f)(\phi(p)) = X(f \circ \phi)(p).$$

The last statement is a direct consequence of Proposition 4.26. □

Definition 4.28. Let M be a differentiable manifold. Two vector fields $X, Y \in C^\infty(TM)$ are said to **commute** if their Lie bracket vanishes i.e. $[X, Y] = 0$.

Proposition 4.29. *Let M be a differentiable manifold, (U, x) be a chart on M and*

$$\left\{ \frac{\partial}{\partial x_k} \mid k = 1, 2, \dots, m \right\}$$

be the induced local frame for the tangent bundle TM . Then the local frame fields commute i.e.

$$[\frac{\partial}{\partial x_k}, \frac{\partial}{\partial x_l}] = 0 \text{ for all } k, l = 1, \dots, m.$$

PROOF. The map $x : U \rightarrow x(U)$ is bijective and differentiable. The vector field $\partial/\partial x_k \in C^\infty(TU)$ is x -related to the coordinate vector field $\partial_{e_k} \in C^\infty(Tx(U))$. Then Proposition 4.27 implies that

$$dx([\frac{\partial}{\partial x_k}, \frac{\partial}{\partial x_l}]) = [\partial_{e_k}, \partial_{e_l}] = 0.$$

The last equation is an immediate consequence of the following well known fact

$$[\partial_{e_k}, \partial_{e_l}](f) = \partial_{e_k}(\partial_{e_l}(f)) - \partial_{e_l}(\partial_{e_k}(f)) = 0$$

for all $f \in C^2(x(U))$. The result now follows from the fact that the linear map $dx_p : T_p M \rightarrow T_{x(p)} \mathbb{R}^m$ is bijective for all $p \in U$. \square

We will now introduce the important notion of a left-invariant vector field on a Lie group. This will play an important role in what follows.

Definition 4.30. Let G be a Lie group. Then a vector field $X \in C^\infty(TG)$ on G is said to be **left-invariant** if it is L_p -related to itself for all $p \in G$ i.e.

$$(dL_p)_q(X_q) = X_{pq} \text{ for all } p, q \in G.$$

The set of left-invariant vector fields on G is called the **Lie algebra** of G and denoted by \mathfrak{g} .

Remark 4.31. It should be noted that if e is the neutral element of the Lie group G and $X \in \mathfrak{g}$ is a left-invariant vector field then

$$X_p = (dL_p)_e(X_e).$$

This shows that the value X_p of X at p is completely determined by the value X_e at e . Hence the map $\Phi : T_e G \rightarrow \mathfrak{g}$ given by

$$\Phi : X_e \mapsto (X : p \mapsto (dL_p)_e(X_e))$$

is a vector space isomorphism. As a direct consequence we see that the Lie algebra \mathfrak{g} is a finite dimensional subspace of $C^\infty(TG)$ of the same dimension as G .

Proposition 4.32. If G is a Lie group then its Lie algebra \mathfrak{g} is a **Lie subalgebra** of $C^\infty(TG)$ i.e. if $X, Y \in \mathfrak{g}$ then $[X, Y] \in \mathfrak{g}$.

PROOF. If $p \in G$ then the left translation $L_p : G \rightarrow G$ is a diffeomorphism so it follows from Proposition 4.27 that

$$dL_p([X, Y]) = [dL_p(X), dL_p(Y)] = [X, Y]$$

for all $X, Y \in \mathfrak{g}$. This proves that the Lie bracket $[X, Y]$ of two left-invariant vector fields X, Y is also left-invariant. \square

The linear isomorphism $\Phi : T_e G \rightarrow \mathfrak{g}$ given by

$$\Phi : X_e \mapsto (X : p \mapsto (dL_p)_e(X_e))$$

induces a natural Lie bracket $[\cdot, \cdot] : T_e G \times T_e G \rightarrow T_e G$ on the tangent space $T_e G$ via

$$[X_e, Y_e] = [X, Y]_e.$$

Notation 4.33. For the classical matrix Lie groups introduced in Chapter 3, we denote their Lie algebras by $\mathfrak{gl}_m(\mathbb{R})$, $\mathfrak{sl}_m(\mathbb{R})$, $\mathfrak{o}(m)$, $\mathfrak{so}(m)$, $\mathfrak{gl}_m(\mathbb{C})$, $\mathfrak{sl}_m(\mathbb{C})$, $\mathfrak{u}(m)$ and $\mathfrak{su}(m)$, respectively.

The following result is a useful tool for handling the Lie brackets for the classical matrix Lie groups. They can simply be calculated by means of the usual matrix multiplication.

Proposition 4.34. *Let G be one of the classical matrix Lie groups and $T_e G$ be the tangent space of G at the neutral element e . Then the Lie bracket $[\cdot, \cdot] : T_e G \times T_e G \rightarrow T_e G$ is given by*

$$[X_e, Y_e] = X_e \cdot Y_e - Y_e \cdot X_e$$

where \cdot is the standard matrix multiplication.

PROOF. We prove the result for the general linear group $\mathbf{GL}_m(\mathbb{R})$. For the other real groups the result follows from the fact that they are all subgroups of $\mathbf{GL}_m(\mathbb{R})$. The same proof can be used in the complex cases.

Let $X, Y \in \mathfrak{gl}_m(\mathbb{R})$ be left-invariant vector fields, $f : U \rightarrow \mathbb{R}$ be a function defined locally around the identity element e and p be an arbitrary point of U . Then the first order derivative $X_p(f)$ of f at p is given by

$$X_p(f) = \frac{d}{dt}(f(p \cdot \text{Exp}(tX_e)))|_{t=0} = df_p(p \cdot X_e) = df_p(X_p).$$

The general linear group $\mathbf{GL}_m(\mathbb{R})$ is an open subset of $\mathbb{R}^{m \times m}$ so we can apply standard arguments from multivariable calculus. The second order derivative $Y_e(X(f))$ satisfies

$$Y_e(X(f)) = \frac{d}{dt}(X_{\text{Exp}(tY_e)}(f))|_{t=0}$$

$$\begin{aligned}
&= \frac{d}{dt}(df_{\text{Exp}(tY_e)}(\text{Exp}(tY_e) \cdot X_e))|_{t=0} \\
&= d^2f_e(Y_e, X_e) + df_e(Y_e \cdot X_e).
\end{aligned}$$

Here d^2f_e is the symmetric Hessian of the function f . As an immediate consequence we obtain

$$\begin{aligned}
[X, Y]_e(f) &= X_e(Y(f)) - Y_e(X(f)) \\
&= d^2f_e(X_e, Y_e) + df_e(X_e \cdot Y_e) \\
&\quad - d^2f_e(Y_e, X_e) - df_e(Y_e \cdot X_e) \\
&= df_e(X_e \cdot Y_e - Y_e \cdot X_e).
\end{aligned}$$

This last calculation implies the statement. \square

The following result shows that the tangent bundle of any Lie group is trivial.

Theorem 4.35. *The tangent bundle TG of a Lie group G is trivial.*

PROOF. Let $\{(X_1)_e, \dots, (X_m)_e\}$ be a basis for T_eG and extend each $(X_k)_e \in T_eG$ to the left-invariant vector field $X_k \in \mathfrak{g}$ with

$$(X_k)_p = (dL_p)_e((X_k)_e).$$

For each $p \in G$ the left translation $L_p : G \rightarrow G$ is a diffeomorphism so the set $\{(X_1)_p, \dots, (X_m)_p\}$ is a basis for the tangent space T_pG . This means that the map $\psi : TG \rightarrow G \times \mathbb{R}^m$ given by

$$\psi : (p, \sum_{k=1}^m v_k \cdot (X_k)_p) \mapsto (p, (v_1, \dots, v_m))$$

is well defined. This is a global bundle chart so the tangent bundle TG is therefore trivial. \square

Exercises

Exercise 4.1. Let $(M, \hat{\mathcal{A}})$ be a smooth manifold, (U, x) , (V, y) be charts such that $U \cap V$ is non-empty and

$$f = y \circ x^{-1} : x(U \cap V) \rightarrow \mathbb{R}^m$$

be the corresponding transition map. Show that the local frames

$$\left\{ \frac{\partial}{\partial x_i} \mid i = 1, \dots, m \right\} \quad \text{and} \quad \left\{ \frac{\partial}{\partial y_j} \mid j = 1, \dots, m \right\}$$

for TM on $U \cap V$ are related as follows

$$\frac{\partial}{\partial x_i} = \sum_{j=1}^m \frac{\partial(f_j \circ x)}{\partial x_i} \cdot \frac{\partial}{\partial y_j}.$$

Exercise 4.2. Let $\mathbf{SO}(m)$ be the special orthogonal group.

- (i) Find a basis for the tangent space $T_e \mathbf{SO}(m)$,
- (ii) construct a non-vanishing vector field $Z \in C^\infty(T\mathbf{SO}(m))$,
- (iii) determine all smooth vector fields on $\mathbf{SO}(2)$.

The Hairy Ball Theorem. There does not exist a continuous non-vanishing vector field $X \in C^0(TS^{2m})$ on the even dimensional sphere S^{2m} .

Exercise 4.3. Employ the Hairy Ball Theorem to show that the tangent bundle TS^{2m} is not trivial. Then construct a non-vanishing vector field $X \in C^\infty(TS^{2m+1})$ on the odd-dimensional sphere S^{2m+1} .

Exercise 4.4. Find a proof of Theorem 4.22.

Exercise 4.5. The Lie algebra $\mathfrak{sl}_2(\mathbb{R})$ of the special linear group $\mathbf{SL}_2(\mathbb{R})$ is generated by

$$X = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad Z = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Show that the Lie brackets of $\mathfrak{sl}_2(\mathbb{R})$ satisfy

$$[X, Y] = 2Z, \quad [Z, X] = 2Y, \quad [Y, Z] = -2X.$$

Exercise 4.6. The Lie algebra $\mathfrak{su}(2)$ of the special unitary group $\mathbf{SU}(2)$ is generated by

$$X = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad Z = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

Show that the corresponding Lie bracket relations are given by

$$[X, Y] = 2Z, \quad [Z, X] = 2Y, \quad [Y, Z] = 2X.$$

Exercise 4.7. The Lie algebra $\mathfrak{so}(3)$ of the special orthogonal group $\mathbf{SO}(3)$ is generated by

$$X = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad Z = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Show that the corresponding Lie bracket relations are given by

$$[X, Y] = Z, \quad [Z, X] = Y, \quad [Y, Z] = X.$$

Compare this result with Example 4.21.

Riemannian Manifolds

In this chapter we introduce the notion of a Riemannian manifold (M, g) . The metric g provides us with a scalar product on each tangent space and can be used to measure angles and the lengths of curves in the manifold. This defines a distance function and turns the manifold into a metric space in a natural way. The Riemannian metric on a differentiable manifold is the most important example of what is called a tensor field.

Let M be a smooth manifold, $C^\infty(M)$ denote the commutative ring of smooth functions on M and $C^\infty(TM)$ be the set of smooth vector fields on M forming a **module** over $C^\infty(M)$. Put

$$C_0^\infty(TM) = C^\infty(M)$$

and for each positive integer r let

$$C_r^\infty(TM) = C^\infty(TM) \otimes \cdots \otimes C^\infty(TM)$$

be the r -fold tensor product of $C^\infty(TM)$ over the commutative ring $C^\infty(M)$.

Definition 5.1. Let M be a differentiable manifold. A smooth **tensor field** A on M of type (r, s) is a map

$$A : C_r^\infty(TM) \rightarrow C_s^\infty(TM)$$

which is multilinear over the commutative ring $C^\infty(M)$ i.e. satisfying

$$\begin{aligned} & A(X_1 \otimes \cdots \otimes X_{k-1} \otimes (f \cdot Y + g \cdot Z) \otimes X_{k+1} \otimes \cdots \otimes X_r) \\ &= f \cdot A(X_1 \otimes \cdots \otimes X_{k-1} \otimes Y \otimes X_{k+1} \otimes \cdots \otimes X_r) \\ & \quad + g \cdot A(X_1 \otimes \cdots \otimes X_{k-1} \otimes Z \otimes X_{k+1} \otimes \cdots \otimes X_r) \end{aligned}$$

for all $X_1, \dots, X_r, Y, Z \in C^\infty(TM)$, $f, g \in C^\infty(M)$ and $k = 1, \dots, r$.

Notation 5.2. For the rest of this work we will for $A(X_1 \otimes \cdots \otimes X_r)$ use the notation

$$A(X_1, \dots, X_r).$$

The next fundamental result provides us with the most important property of a tensor field. It shows that the value $A(X_1, \dots, X_r)(p)$ of $A(X_1, \dots, X_r)$ at a point $p \in M$ only depends on the values of the vector fields X_1, \dots, X_r at p and is independent of their values away from p .

Proposition 5.3. *Let $A : C_r^\infty(TM) \rightarrow C_s^\infty(TM)$ be a tensor field of type (r, s) and $p \in M$. Let X_1, \dots, X_r and Y_1, \dots, Y_r be smooth vector fields on M such that $(X_k)_p = (Y_k)_p$ for each $k = 1, \dots, r$. Then*

$$A(X_1, \dots, X_r)(p) = A(Y_1, \dots, Y_r)(p).$$

PROOF. We will prove the statement for $r = 1$, the rest follows by induction. Put $X = X_1$ and $Y = Y_1$ and let (U, x) be a chart on M . Choose a function $f \in C^\infty(M)$ such that $f(p) = 1$,

$$\text{support}(f) = \overline{\{p \in M \mid f(p) \neq 0\}}$$

is contained in U and define the vector fields $v_1, \dots, v_m \in C^\infty(TM)$ on M by

$$(v_k)_q = \begin{cases} f(q) \cdot (\frac{\partial}{\partial x_k})_q & \text{if } q \in U \\ 0 & \text{if } q \notin U \end{cases}$$

Then there exist functions $\rho_k, \sigma_k \in C^\infty(M)$ such that

$$f \cdot X = \sum_{k=1}^m \rho_k \cdot v_k \quad \text{and} \quad f \cdot Y = \sum_{k=1}^m \sigma_k \cdot v_k.$$

This implies that

$$\begin{aligned} A(X)(p) &= f(p)A(X)(p) \\ &= (f \cdot A(X))(p) \\ &= A(f \cdot X)(p) \\ &= A\left(\sum_{k=1}^m \rho_k \cdot v_k\right)(p) \\ &= \sum_{k=1}^m (\rho_k \cdot A(v_k))(p) \\ &= \sum_{k=1}^m \rho_k(p)A(v_k)(p) \end{aligned}$$

and similarly

$$A(Y)(p) = \sum_{k=1}^m \sigma_k(p)A(v_k)(p).$$

The fact that $X_p = Y_p$ shows that $\rho_k(p) = \sigma_k(p)$ for all k . As a direct consequence we see that

$$A(X)(p) = A(Y)(p).$$

□

Following the result of Proposition 5.3 we now introduce the following useful notation.

Notation 5.4. For a tensor field $A : C_r^\infty(TM) \rightarrow C_s^\infty(TM)$ of type (r, s) we will by A_p denote the real multilinear restriction of A to the r -fold tensor product $T_p M \otimes \cdots \otimes T_p M$ of the real vector space $T_p M$ given by

$$A_p : ((X_1)_p, \dots, (X_r)_p) \mapsto A(X_1, \dots, X_r)(p).$$

Next we introduce the notion of a Riemannian metric. This is the most important example of a tensor field in Riemannian geometry.

Definition 5.5. Let M be a smooth manifold. A **Riemannian metric** g on M is a tensor field $g : C_2^\infty(TM) \rightarrow C_0^\infty(TM)$ such that for each $p \in M$ the restriction g_p of g to the tensor product $T_p M \otimes T_p M$ with

$$g_p : (X_p, Y_p) \mapsto g(X, Y)(p)$$

is a scalar product on the tangent space $T_p M$. The pair (M, g) is said to be a **Riemannian manifold**. The study of Riemannian manifolds is called Riemannian Geometry. The geometric properties of (M, g) which only depend on the metric g are said to be **intrinsic** or **metric** properties.

The classical m -dimensional Euclidean space E^m is a Riemannian manifold as follows.

Example 5.6. The standard Euclidean scalar product on the vector space \mathbb{R}^m given by

$$\langle X, Y \rangle_{\mathbb{R}^m} = X^t \cdot Y = \sum_{k=1}^m X_k Y_k$$

defines a Riemannian metric on \mathbb{R}^m . The Riemannian manifold

$$E^m = (\mathbb{R}^m, \langle \cdot, \cdot \rangle_{\mathbb{R}^m})$$

is called the m -dimensional **Euclidean space**.

Definition 5.7. Let (M, g) be a Riemannian manifold and $\gamma : I \rightarrow M$ be a C^1 -curve in M . Then the **length** $L(\gamma)$ of γ is defined by

$$L(\gamma) = \int_I \sqrt{g(\dot{\gamma}(t), \dot{\gamma}(t))} dt.$$

The standard punctured round sphere has the following description as a Riemannian manifold.

Example 5.8. Equip the vector space \mathbb{R}^m with the Riemannian metric g given by

$$g_p(X, Y) = \frac{4}{(1 + |p|_{\mathbb{R}^m}^2)^2} \langle X, Y \rangle_{\mathbb{R}^m}.$$

The Riemannian manifold $\Sigma^m = (\mathbb{R}^m, g)$ is called the m -dimensional **punctured round sphere**. Let $\gamma : \mathbb{R}^+ \rightarrow \Sigma^m$ be the curve with

$$\gamma : t \mapsto (t, 0, \dots, 0).$$

Then the length $L(\gamma)$ of γ can be determined as follows.

$$L(\gamma) = 2 \int_0^\infty \frac{\sqrt{\langle \dot{\gamma}, \dot{\gamma} \rangle}}{1 + |\gamma|^2} dt = 2 \int_0^\infty \frac{dt}{1 + t^2} = 2[\arctan(t)]_0^\infty = \pi.$$

The important hyperbolic space H^m can be modelled in different ways. In the following Example 5.9 we present it as the open unit ball. For the upper half space model see Exercise 8.8.

Example 5.9. Let $B_1^m(0)$ be the open unit ball in \mathbb{R}^m given by

$$B_1^m(0) = \{p \in \mathbb{R}^m \mid |p|_{\mathbb{R}^m} < 1\}.$$

By the m -dimensional **hyperbolic space** we mean $B_1^m(0)$ equipped with the Riemannian metric

$$g_p(X, Y) = \frac{4}{(1 - |p|_{\mathbb{R}^m}^2)^2} \langle X, Y \rangle_{\mathbb{R}^m}.$$

Let $\gamma : (0, 1) \rightarrow B_1^m(0)$ be the curve given by

$$\gamma : t \mapsto (t, 0, \dots, 0).$$

Then the length $L(\gamma)$ of γ can be determined as follows.

$$L(\gamma) = 2 \int_0^1 \frac{\sqrt{\langle \dot{\gamma}, \dot{\gamma} \rangle}}{1 - |\gamma|^2} dt = 2 \int_0^1 \frac{dt}{1 - t^2} = [\log(\frac{1+t}{1-t})]_0^1 = \infty.$$

The following result tells us that a Riemannian manifold (M, g) has the structure of a metric space (M, d) in a natural way.

Proposition 5.10. *Let (M, g) be a Riemannian manifold which is path-connected. For two points $p, q \in M$ let C_{pq} denote the set of C^1 -curves $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = p$ and $\gamma(1) = q$ and define the function $d : M \times M \rightarrow \mathbb{R}_0^+$ by*

$$d(p, q) = \inf\{L(\gamma) \mid \gamma \in C_{pq}\}.$$

*Then (M, d) is a **metric space** i.e. for all $p, q, r \in M$ we have*

- (i) $d(p, q) \geq 0$,
- (ii) $d(p, q) = 0$ if and only if $p = q$,
- (iii) $d(p, q) = d(q, p)$,
- (iv) $d(p, q) \leq d(p, r) + d(r, q)$.

The topology on M induced by the metric d is identical to the one M carries as a topological manifold (M, \mathcal{T}) , see Definition 2.1.

PROOF. See for example: Peter Petersen, *Riemannian Geometry*, Graduate Texts in Mathematics **171**, Springer (1998). \square

A Riemannian metric on a differentiable manifold induces a Riemannian metric on its submanifolds as follows.

Definition 5.11. Let (N, h) be a Riemannian manifold and M be a submanifold. Then the smooth tensor field $g : C_2^\infty(TM) \rightarrow C_0^\infty(M)$ given by

$$g(X, Y) : p \mapsto h_p(X_p, Y_p)$$

is a Riemannian metric on M . It is called the **induced metric** on M in (N, h) .

We can now equip some of the manifolds which we introduced in Chapter 2 with a Riemannian metric.

Example 5.12. The standard Euclidean metric $\langle \cdot, \cdot \rangle_{\mathbb{R}^n}$ on \mathbb{R}^n induces Riemannian metrics on the following submanifolds.

- (i) the sphere $S^m \subset \mathbb{R}^n$, with $n = m + 1$,
- (ii) the tangent bundle $TS^m \subset \mathbb{R}^n$, where $n = 2(m + 1)$,
- (iii) the torus $T^m \subset \mathbb{R}^n$, with $n = 2m$,

Example 5.13. The vector space $\mathbb{C}^{m \times m}$ of complex $m \times m$ matrices carries a natural Riemannian metric g given by

$$g(Z, W) = \operatorname{Re}(\operatorname{trace}(\bar{Z}^t W))$$

for all $Z, W \in \mathbb{C}^{m \times m}$. This induces metrics on the submanifolds of $\mathbb{C}^{m \times m}$ such as $\mathbb{R}^{m \times m}$ and the classical Lie groups $\mathbf{GL}_m(\mathbb{R})$, $\mathbf{SL}_m(\mathbb{R})$, $\mathbf{O}(m)$, $\mathbf{SO}(m)$, $\mathbf{GL}_m(\mathbb{C})$, $\mathbf{SL}_m(\mathbb{C})$, $\mathbf{U}(m)$, $\mathbf{SU}(m)$.

Our next aim is to prove that every differentiable manifold M can be equipped with a Riemannian metric g . For this we need Fact 5.15.

Definition 5.14. Let M be a differentiable manifold. Then a partition of unity consists of a family $\{f_\alpha : M \rightarrow \mathbb{R} \mid \alpha \in \mathcal{I}\}$ of differentiable real-valued functions such that

- (i) $0 \leq f_\alpha \leq 1$ for all $\alpha \in \mathcal{I}$,

- (ii) every point $p \in M$ has a neighbourhood which intersects only finitely many of the sets $\text{support}(f_\alpha) = \overline{\{p \in M \mid f_\alpha(p) \neq 0\}}$, and
- (iii)

$$\sum_{\alpha \in \mathcal{I}} f_\alpha = 1.$$

Note that the sum in (iii) is finite at each point $p \in M$.

For the proof of the following interesting result, it is important that M is a Hausdorff space with a countable basis, see Definition 2.1.

Fact 5.15. *Let M be a differentiable manifold and $(U_\alpha)_{\alpha \in \mathcal{I}}$ be an open covering of M such that for each $\alpha \in \mathcal{I}$ the pair (U_α, ϕ_α) is a chart on M . Then there exist*

- (i) *a locally finite open cover $(W_\beta)_{\beta \in \mathcal{J}}$ such that each W_β is contained in U_α for some $\alpha \in \mathcal{I}$. Furthermore, for each $\beta \in \mathcal{J}$, W_β is an open neighbourhood for a chart (W_β, x_β) , and*
- (ii) *a partition of unity $(f_\beta)_{\beta \in \mathcal{J}}$ such that the support (f_β) is contained in the open subset W_β .*

We are now ready to prove the following important statement.

Theorem 5.16. *Let $(M^m, \hat{\mathcal{A}})$ be a differentiable manifold. Then there exists a Riemannian metric g on M .*

PROOF. For each point $p \in M$, let $(U_p, \phi_p) \in \hat{\mathcal{A}}$ be a chart such that $p \in U_p$. Then $(U_p)_{p \in M}$ is an open covering and let (W_β, x^β) be charts on M as in Fact 5.15. Let $(f_\beta)_{\beta \in \mathcal{J}}$ be a partition of unity such that the support (f_β) is contained in W_β . Further, let $\langle \cdot, \cdot \rangle_{\mathbb{R}^m}$ be the standard Euclidean metric on \mathbb{R}^m . Then for each $\beta \in \mathcal{J}$ we define

$$g_\beta : C_2^\infty(TM) \rightarrow C_0^\infty(TM)$$

by

$$g_\beta\left(\frac{\partial}{\partial x_k^\beta}, \frac{\partial}{\partial x_l^\beta}\right)(p) = \begin{cases} f_\beta(p) \cdot \langle e_k, e_l \rangle_{\mathbb{R}^m} & \text{if } p \in W_\beta \\ 0 & \text{if } p \notin W_\beta \end{cases}$$

Note that at each point only finitely many of g_β are non-zero. This means that the well defined tensor $g : C_2^\infty(TM) \rightarrow C_0^\infty(TM)$ given by

$$g = \sum_{\beta \in \mathcal{J}} g_\beta$$

is a Riemannian metric on M . □

Definition 5.17. Let (M, g) and (N, h) be Riemannian manifolds. A map $\phi : (M, g) \rightarrow (N, h)$ is said to be **conformal** if there exists a function $\lambda : M \rightarrow \mathbb{R}$ such that

$$e^{\lambda(p)} \cdot g_p(X_p, Y_p) = h_{\phi(p)}(d\phi_p(X_p), d\phi_p(Y_p)),$$

for all $X, Y \in C^\infty(TM)$ and $p \in M$. The positive real-valued function e^λ is called the **conformal factor** of ϕ . A conformal map with $\lambda \equiv 0$ i.e. $e^\lambda \equiv 1$ is said to be **isometric**. An isometric diffeomorphism is called an **isometry**.

Definition 5.18. For a Riemannian manifold (M, g) we denote by $\mathcal{I}(M)$ the set of its isometries. If $\phi, \psi \in \mathcal{I}(M)$ then it is clear that the composition $\psi \circ \phi$ and the inverse ϕ^{-1} are also isometries. The operation is clearly associative and the identity map is its neutral element. The pair $(\mathcal{I}(M), \circ)$ is called the **isometry group** of (M, g) .

It can be shown that the isometry group of a differentiable manifold has the structure of a Lie group. For this see: R. S. Palais, *On the differentiability of isometries*, Proc. Amer. Math. Soc. **8** (1957), 805-807.

We next introduce the important notion of a Riemannian homogeneous spaces. The classical reference is: S. Kobayashi, K. Nomizu, *Foundations of differential geometry*, Vol. II, John Wiley & Sons (1969).

Definition 5.19. The isometry group $\mathcal{I}(M)$ of a Riemannian manifold (M, g) is said to be **transitive** if for all $p, q \in M$ there exists an isometry $\phi_{pq} : M \rightarrow M$ such that $\phi_{pq}(p) = q$. In that case (M, g) is called a **Riemannian homogeneous space**.

An important subclass of Riemannian homogeneous spaces is that of symmetric spaces introduced in Definition 7.30.

Example 5.20. On the standard unit sphere S^m we have the action $\alpha : \mathbf{O}(m+1) \times S^m \rightarrow S^m$ of the orthogonal group $\mathbf{O}(m+1)$ given by

$$\alpha : (p, x) \mapsto p \cdot x$$

where \cdot is the standard matrix multiplication. The following shows that this action on S^m is isometric

$$\langle pX, pY \rangle = X^t p^t p Y = X^t Y = \langle X, Y \rangle.$$

This means that the orthogonal group $\mathbf{O}(m+1)$ is a subgroup of the isometry group $\mathcal{I}(S^m)$. It is easily seen that $\mathbf{O}(m+1)$ acts transitively on the sphere S^m so this is a homogeneous space.

Example 5.21. The standard Euclidean scalar product on the real vector space $\mathbb{R}^{m \times m}$ induces a Riemannian metric on the orthogonal group $\mathbf{O}(m)$ given by

$$g(X, Y) = \text{trace}(X^t \cdot Y).$$

Applying the left translation $L_p : \mathbf{O}(m) \rightarrow \mathbf{O}(m)$, with $L_p : q \mapsto pq$, we see that the tangent space $T_p \mathbf{O}(m)$ of $\mathbf{O}(m)$ at p is

$$T_p \mathbf{O}(m) = \{pX \mid X^t + X = 0\}.$$

The differential $(dL_p)_q : T_q \mathbf{O}(m) \rightarrow T_{pq} \mathbf{O}(m)$ of L_p at $q \in \mathbf{O}(m)$ satisfies

$$(dL_p)_q : qX \mapsto pqX.$$

We then have

$$\begin{aligned} g_{pq}((dL_p)_q(qX), (dL_p)_q(qY)) &= \text{trace}((pqX)^t pqY) \\ &= \text{trace}(X^t q^t p^t pqY) \\ &= \text{trace}(qX)^t (qY). \\ &= g_q(qX, qY). \end{aligned}$$

This shows that the left translation $L_p : \mathbf{O}(m) \rightarrow \mathbf{O}(m)$ is an isometry for all $p \in \mathbf{O}(m)$.

Definition 5.22. A Riemannian metric g on a Lie group G is said to be **left-invariant** if for each $p \in G$ the left translation $L_p : G \rightarrow G$ is an isometry. A Lie group (G, g) with a left-invariant metric is called a **Riemannian Lie group**.

Remark 5.23. It should be noted that if (G, g) is a Riemannian Lie group and $X, Y \in \mathfrak{g}$ are left-invariant vector fields then

$$g_p(X_p, Y_p) = g_p((dL_p)_e(X_e), (dL_p)_e(Y_e)) = g_e(X_e, Y_e).$$

This tells us that a left-invariant metric g on G is completely determined by the scalar product $g_e : T_e G \times T_e G \rightarrow \mathbb{R}$ on the tangent space at the neutral element $e \in G$.

Theorem 5.24. *A Riemannian Lie group (G, g) is a Riemannian homogeneous space.*

PROOF. For arbitrary elements $p, q \in G$ the left-translation $\phi_{pq} = L_{qp^{-1}}$ by $qp^{-1} \in G$ is an isometry satisfying $\phi_{pq}(p) = q$. This shows that the isometry group $\mathcal{I}(G)$ is transitive. \square

In Example 2.6 we introduced the real projective space $\mathbb{R}P^m$ as an abstract differentiable manifold. We will now equip this with a natural Riemannian metric.

Example 5.25. Let S^m be the unit sphere in \mathbb{R}^{m+1} and $\text{Sym}(\mathbb{R}^{m+1})$ be the vector space of real symmetric $(m+1) \times (m+1)$ matrices equipped with the Riemannian metric g given by

$$g(X, Y) = \frac{1}{8} \text{trace}(X^t \cdot Y).$$

As in Example 3.26, we define the immersion $\phi : S^m \rightarrow \text{Sym}(\mathbb{R}^{m+1})$ by

$$\phi : p \mapsto (\rho_p : q \mapsto 2\langle q, p \rangle p - q).$$

This maps a point $p \in S^m$ to the reflection $\rho_p : \mathbb{R}^{m+1} \rightarrow \mathbb{R}^{m+1}$ about the real line generated by p . This is clearly a symmetric bijective linear map.

Let $\alpha, \beta : \mathbb{R} \rightarrow S^m$ be two curves such that $\alpha(0) = p = \beta(0)$ and put $X = \dot{\alpha}(0)$, $Y = \dot{\beta}(0)$. Then for $\gamma \in \{\alpha, \beta\}$ we have

$$d\phi_p(\dot{\gamma}(0)) = (q \mapsto 2\langle q, \dot{\gamma}(0) \rangle p + 2\langle q, p \rangle \dot{\gamma}(0)).$$

If \mathcal{B} is an orthonormal basis for \mathbb{R}^{m+1} , then

$$\begin{aligned} g(d\phi_p(X), d\phi_p(Y)) &= \frac{1}{8} \text{trace}(d\phi_p(X)^t \cdot d\phi_p(Y)) \\ &= \frac{1}{8} \sum_{q \in \mathcal{B}} \langle q, d\phi_p(X)^t \cdot d\phi_p(Y) q \rangle \\ &= \frac{1}{8} \sum_{q \in \mathcal{B}} \langle d\phi_p(X) q, d\phi_p(Y) q \rangle \\ &= \frac{1}{2} \sum_{q \in \mathcal{B}} \langle \langle q, X \rangle p + \langle q, p \rangle X, \langle q, Y \rangle p + \langle q, p \rangle Y \rangle \\ &= \frac{1}{2} \sum_{q \in \mathcal{B}} \{ \langle p, p \rangle \langle X, q \rangle \langle q, Y \rangle + \langle X, Y \rangle \langle p, q \rangle \langle p, q \rangle \} \\ &= \frac{1}{2} \{ \langle X, Y \rangle + \langle X, Y \rangle \} \\ &= \langle X, Y \rangle. \end{aligned}$$

This proves that the immersion $\phi : S^m \rightarrow \text{Sym}(\mathbb{R}^{m+1})$ is isometric. In Example 3.26 we have seen that the image $\phi(S^m)$ can be identified with the real projective space $\mathbb{R}P^m$. This inherits the induced metric from $\text{Sym}(\mathbb{R}^{m+1})$. The map $\phi : S^m \rightarrow \mathbb{R}P^m$ is what is called an isometric double cover of $\mathbb{R}P^m$.

Proposition 5.26. *Let $\mathbb{R}P^2$ be the two dimensional real projective plane equipped with the Riemannian metric introduced in Example 5.25. Then the surface area of $\mathbb{R}P^2$ is 2π .*

PROOF. Example 5.25 shows that if m is a positive integer then the map $\phi : S^m \rightarrow \mathbb{R}P^m$ is an isometric double cover. Hence this is locally volume preserving. This implies that the m -dimensional volume satisfies

$$\text{vol}(S^m) = 2 \cdot \text{vol}(\mathbb{R}P^m).$$

In particular,

$$\text{vol}(\mathbb{R}P^2) = \frac{1}{2} \cdot \text{vol}(S^2) = 2\pi.$$

□

Long before John Nash became famous in Hollywood he proved the next remarkable result in his paper: J. Nash, *The imbedding problem for Riemannian manifolds*, Ann. Math. **63** (1956), 20-63. It implies that every Riemannian manifold can be realised as a submanifold of a Euclidean space. The original proof by Nash was later simplified, see for example: Matthias Günther, *On the perturbation problem associated to isometric embeddings of Riemannian manifolds*, Ann. Global Anal. Geom. **7** (1989), 69-77.

Deep Result 5.27. For $3 \leq r \leq \infty$ let (M, g) be a Riemannian C^r -manifold. Then there exists an isometric C^r -embedding of (M, g) into a Euclidean space \mathbb{R}^n .

Remark 5.28. The reader should note that in Example 5.25 we have embedded the Riemannian manifold $\mathbb{R}P^m$ in the Euclidean space $\text{Sym}(\mathbb{R}^{m+1})$.

Remark 5.29. We will now see that local parametrizations are very useful tools for studying the intrinsic geometry of a Riemannian manifold (M, g) . Let p be a point of M and $\hat{\psi} : U \rightarrow M$ be a local parametrization of M with $q \in U$ and $\hat{\psi}(q) = p$. The differential $d\hat{\psi}_q : T_q\mathbb{R}^m \rightarrow T_pM$ is bijective so, following the inverse mapping theorem, there exist neighbourhoods U_q of q and U_p of p such that the restriction $\psi = \hat{\psi}|_{U_q} : U_q \rightarrow U_p$ is a diffeomorphism. On U_q we have the canonical frame $\{e_1, \dots, e_m\}$ for TU_q so $\{d\psi(e_1), \dots, d\psi(e_m)\}$ is a local frame for TM over U_p . We then define the pull-back metric $\tilde{g} = \psi^*g$ on U_q by

$$\tilde{g}(e_k, e_l) = g(d\psi(e_k), d\psi(e_l)).$$

Then $\psi : (U_q, \tilde{g}) \rightarrow (U_p, g)$ is an isometry so the intrinsic geometry of (U_q, \tilde{g}) and that of (U_p, g) are exactly the same.

Example 5.30. Let G be one of the classical Lie groups and e be the neutral element of G . Let $\{X_1, \dots, X_m\}$ be a basis for the Lie algebra \mathfrak{g} of G . For $p \in G$ define $\psi_p : \mathbb{R}^m \rightarrow G$ by

$$\psi_p : (t_1, \dots, t_m) \mapsto L_p\left(\prod_{k=1}^m \text{Exp}(t_k X_k(e))\right)$$

where $L_p : G \rightarrow G$ is the left translation given by $L_p(q) = pq$. Then

$$(d\psi_p)_0(e_k) = X_k(p)$$

for all k . This means that the differential $(d\psi_p)_0 : T_0\mathbb{R}^m \rightarrow T_pG$ is an isomorphism so there exist open neighbourhoods U_0 of 0 and U_p of p such that the restriction of ψ to U_0 is bijective onto its image U_p and hence a local parametrization of G around p .

We will now study the normal bundle of a submanifold of a given Riemannian manifold. This is an important example of the notion of a vector bundle over a manifold.

Definition 5.31. Let (N, h) be a Riemannian manifold and M be a submanifold. For a point $p \in M$ we define the **normal space** N_pM of M at p by

$$N_pM = \{X \in T_pN \mid h_p(X, Y) = 0 \text{ for all } Y \in T_pM\}.$$

For all $p \in M$ we have the orthogonal decomposition

$$T_pN = T_pM \oplus N_pM.$$

The **normal bundle** of M in N is defined by

$$NM = \{(p, X) \mid p \in M, X \in N_pM\}.$$

Theorem 5.32. Let (N^n, h) be a Riemannian manifold and M^m be a smooth submanifold. Then the normal bundle (NM, M, π) is a smooth vector bundle over M of dimension $(n - m)$.

PROOF. See Exercise 5.6. □

Example 5.33. Let S^m be the unit sphere in \mathbb{R}^{m+1} equipped with its standard Euclidean metric $\langle \cdot, \cdot \rangle$. If $p \in S^m$ then the tangent space T_pS^m of S^m at p is

$$T_pS^m = \{X \in \mathbb{R}^{m+1} \mid \langle p, X \rangle = 0\}$$

so the normal space N_pS^m of S^m at p satisfies

$$N_pS^m = \{\lambda p \in \mathbb{R}^{m+1} \mid \lambda \in \mathbb{R}\}.$$

This shows that the normal bundle NS^m of S^m in \mathbb{R}^{m+1} is given by

$$NS^m = \{(p, \lambda p) \in \mathbb{R}^{2m+2} \mid p \in S^m, \lambda \in \mathbb{R}\}.$$

We will now determine the normal bundle $N\mathbf{O}(m)$ of the orthogonal group $\mathbf{O}(m)$ as a submanifold of $\mathbb{R}^{m \times m}$.

Example 5.34. The standard Euclidean scalar product, on the real vector space $\mathbb{R}^{m \times m}$, induces a left-invariant Riemannian metric g on the subset $\mathbf{O}(m)$. This satisfies

$$g(X, Y) = \text{trace}(X^t \cdot Y).$$

As we have already seen in Example 3.11 the tangent space $T_e\mathbf{O}(m)$ of $\mathbf{O}(m)$ at the neutral element e satisfies

$$T_e\mathbf{O}(m) = \{X \in \mathbb{R}^{m \times m} \mid X^t + X = 0\}.$$

This means that the tangent bundle $T\mathbf{O}(m)$ of $\mathbf{O}(m)$ is given by

$$T\mathbf{O}(m) = \{(p, pX) \mid p \in \mathbf{O}(m), X \in T_e\mathbf{O}(m)\}.$$

The real vector space $\mathbb{R}^{m \times m}$ has a natural linear decomposition

$$\mathbb{R}^{m \times m} = \text{Sym}(\mathbb{R}^m) \oplus T_e\mathbf{O}(m),$$

where every element $X \in \mathbb{R}^{m \times m}$ can be decomposed $X = X^\top + X^\perp$ in its symmetric and skew-symmetric parts given by

$$X^\top = \frac{1}{2}(X + X^t) \quad \text{and} \quad X^\perp = \frac{1}{2}(X - X^t).$$

If $X \in T_e\mathbf{O}(m)$ and $Y \in \text{Sym}(\mathbb{R}^m)$ then

$$\begin{aligned} g(X, Y) &= \text{trace}(X^t Y) \\ &= \text{trace}(Y^t X) \\ &= \text{trace}(XY^t) \\ &= \text{trace}(-X^t Y) \\ &= -g(X, Y). \end{aligned}$$

This shows that $g(X, Y) = 0$ so the normal space $N_e\mathbf{O}(m)$ at the neutral element e of $\mathbf{O}(m)$ satisfies

$$N_e\mathbf{O}(m) = \text{Sym}(\mathbb{R}^m).$$

This means that in this situation the normal bundle $N\mathbf{O}(m)$ of $\mathbf{O}(m)$ is given by

$$N\mathbf{O}(m) = \{(p, pY) \mid p \in \mathbf{O}(m), Y \in \text{Sym}(\mathbb{R}^m)\}.$$

A Riemannian metric g on a differentiable manifold M can be used to construct families of natural metrics on the tangent bundle TM of M . The best known such examples are the Sasaki and Cheeger-Gromoll metrics. For a detailed survey on the geometry of tangent bundles

equipped with these metrics we recommend the paper: S. Gudmundson, E. Kappos, *On the geometry of tangent bundles*, Expo. Math. **20** (2002), 1-41.

Exercises

Exercise 5.1. Let \mathbb{R}^m and \mathbb{C}^m be equipped with their standard Euclidean metrics given by

$$g(z, w) = \operatorname{Re} \sum_{k=1}^m z_k \bar{w}_k$$

and let

$$T^m = \{z \in \mathbb{C}^m \mid |z_1| = \dots = |z_m| = 1\}$$

be the m -dimensional torus in \mathbb{C}^m with the induced metric. Let $\phi : \mathbb{R}^m \rightarrow T^m$ be the standard parametrisation of the m -dimensional torus in \mathbb{C}^m satisfying $\phi : (x_1, \dots, x_m) \mapsto (e^{ix_1}, \dots, e^{ix_m})$. Show that ϕ is isometric.

Exercise 5.2. The **stereographic projection** from the north pole of the m -dimensional sphere

$$\phi : (S^m - \{(1, 0, \dots, 0)\}, \langle, \rangle_{\mathbb{R}^{m+1}}) \rightarrow (\mathbb{R}^m, \frac{4}{(1 + |x|^2)^2} \langle, \rangle_{\mathbb{R}^m})$$

is given by

$$\phi : (x_0, \dots, x_m) \mapsto \frac{1}{1 - x_0} (x_1, \dots, x_m).$$

Show that ϕ is an isometry.

Exercise 5.3. Let $B_1^2(0)$ be the open unit disk in the complex plane equipped with the hyperbolic metric

$$g(X, Y) = \frac{4}{(1 - |z|^2)^2} \langle X, Y \rangle_{\mathbb{R}^2}.$$

Equip the upper half plane $\{z \in \mathbb{C} \mid \operatorname{Im}(z) > 0\}$ with the Riemannian metric

$$g(X, Y) = \frac{1}{\operatorname{Im}(z)^2} \langle X, Y \rangle_{\mathbb{R}^2}.$$

Prove that the holomorphic function $f : B_1^2(0) \rightarrow \{z \in \mathbb{C} \mid \operatorname{Im}(z) > 0\}$ given by

$$f : z \mapsto \frac{i + z}{1 + iz}$$

is an isometry.

Exercise 5.4. Equip the unitary group $\mathbf{U}(m)$ with the Riemannian metric g given by

$$g(Z, W) = \operatorname{Re}(\operatorname{trace}(\bar{Z}^t \cdot W)).$$

Show that for each $p \in \mathbf{U}(m)$ the left translation $L_p : \mathbf{U}(m) \rightarrow \mathbf{U}(m)$ is an isometry.

Exercise 5.5. For the general linear group $\mathbf{GL}_m(\mathbb{R})$ we have two Riemannian metrics g and h satisfying

$$g_p(pZ, pW) = \text{trace}((pZ)^t \cdot pW), \quad h_p(pZ, pW) = \text{trace}(Z^t \cdot W).$$

Further let \hat{g}, \hat{h} be their induced metrics on the special linear group $\mathbf{SL}_m(\mathbb{R})$ as a subset of $\mathbf{GL}_m(\mathbb{R})$.

- (i) Which of the metrics g, h, \hat{g}, \hat{h} are left-invariant?
- (ii) Determine the normal space $N_e \mathbf{SL}_m(\mathbb{R})$ of $\mathbf{SL}_m(\mathbb{R})$ in $\mathbf{GL}_m(\mathbb{R})$ with respect to g
- (iii) Determine the normal bundle $N\mathbf{SL}_m(\mathbb{R})$ of $\mathbf{SL}_m(\mathbb{R})$ in $\mathbf{GL}_m(\mathbb{R})$ with respect to h .

Exercise 5.6. Find a proof of Theorem 5.32.

The Levi-Civita Connection

In this chapter we introduce the Levi-Civita connection ∇ on the tangent bundle TM of a Riemannian manifold (M, g) . This is the most important example of the general notion of a connection on a smooth vector bundle. We deduce the general explicit Koszul formula for the Levi-Civita connection. Then we show how this simplifies in the important case of a Lie group equipped with a left-invariant metric. We also give an example of a metric connection on the normal bundle of a submanifold of a Riemannian manifold and study its properties.

On the m -dimensional real vector space \mathbb{R}^m we have the well known differential operator

$$\partial : C^\infty(T\mathbb{R}^m) \times C^\infty(T\mathbb{R}^m) \rightarrow C^\infty(T\mathbb{R}^m)$$

mapping a pair of vector fields X, Y on \mathbb{R}^m to the classical **directional derivative** $\partial_X Y$ of Y in the direction of X given by

$$(\partial_X Y)(x) = \lim_{t \rightarrow 0} \frac{Y(x + tX(x)) - Y(x)}{t}.$$

The most fundamental properties of the operator ∂ are expressed by the following. If $\lambda, \mu \in \mathbb{R}$, $f, g \in C^\infty(\mathbb{R}^m)$ and $X, Y, Z \in C^\infty(T\mathbb{R}^m)$ then

- (i) $\partial_X(\lambda \cdot Y + \mu \cdot Z) = \lambda \cdot \partial_X Y + \mu \cdot \partial_X Z$,
- (ii) $\partial_X(f \cdot Y) = X(f) \cdot Y + f \cdot \partial_X Y$,
- (iii) $\partial_{(f \cdot X + g \cdot Y)} Z = f \cdot \partial_X Z + g \cdot \partial_Y Z$.

The next result shows that the differential operator ∂ is compatible with both the standard differentiable structure on \mathbb{R}^m and its Euclidean metric.

Proposition 6.1. *Let the real vector space \mathbb{R}^m be equipped with the standard Euclidean metric $\langle \cdot, \cdot \rangle$ and $X, Y, Z \in C^\infty(T\mathbb{R}^m)$ be smooth vector fields on \mathbb{R}^m . Then*

- (iv) $\partial_X Y - \partial_Y X = [X, Y]$,
- (v) $X(\langle Y, Z \rangle) = \langle \partial_X Y, Z \rangle + \langle Y, \partial_X Z \rangle$.

We will now generalise the differential operator ∂ on the Euclidean space \mathbb{R}^m to the so called Levi-Civita connection ∇ on a Riemannian manifold (M, g) . First we introduce the general concept of a connection on a smooth vector bundle.

Definition 6.2. Let M be a smooth manifold. A **connection** $\hat{\nabla}$, on a vector bundle (E, M, π) over M , is an operator

$$\hat{\nabla} : C^\infty(TM) \times C^\infty(E) \rightarrow C^\infty(E)$$

such that

- (i) $\hat{\nabla}_X(\lambda \cdot v + \mu \cdot w) = \lambda \cdot \hat{\nabla}_X v + \mu \cdot \hat{\nabla}_X w$,
- (ii) $\hat{\nabla}_X(f \cdot v) = X(f) \cdot v + f \cdot \hat{\nabla}_X v$,
- (iii) $\hat{\nabla}(f \cdot X + g \cdot Y)v = f \cdot \hat{\nabla}_X v + g \cdot \hat{\nabla}_Y v$

for all $\lambda, \mu \in \mathbb{R}$, $X, Y \in C^\infty(TM)$, $v, w \in C^\infty(E)$ and $f, g \in C^\infty(M)$. A section $v \in C^\infty(E)$ of the vector bundle E is said to be **parallel** with respect to the connection $\hat{\nabla}$ if

$$\hat{\nabla}_X v = 0,$$

for all vector fields $X \in C^\infty(TM)$.

Definition 6.3. Let M be a smooth manifold and $\hat{\nabla}$ be a connection on the tangent bundle (TM, M, π) . Then we define its **torsion**

$$T : C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(TM)$$

by

$$T(X, Y) = \hat{\nabla}_X Y - \hat{\nabla}_Y X - [X, Y],$$

where $[,]$ is the Lie bracket on $C^\infty(TM)$. The connection $\hat{\nabla}$ is said to be **torsion-free** if its torsion T vanishes i.e. if

$$[X, Y] = \hat{\nabla}_X Y - \hat{\nabla}_Y X$$

for all $X, Y \in C^\infty(TM)$.

Definition 6.4. Let (M, g) be a Riemannian manifold. Then a connection $\hat{\nabla}$ on the tangent bundle (TM, M, π) is said to be **metric**, or compatible with the Riemannian metric g , if

$$X(g(Y, Z)) = g(\hat{\nabla}_X Y, Z) + g(Y, \hat{\nabla}_X Z)$$

for all $X, Y, Z \in C^\infty(TM)$.

Let (M, g) be a Riemannian manifold and ∇ be a metric and torsion-free connection on its tangent bundle (TM, M, π) . Then it is easily seen that the following equations hold

$$g(\nabla_X Y, Z) = X(g(Y, Z)) - g(Y, \nabla_X Z),$$

$$\begin{aligned} g(\nabla_X Y, Z) &= g([X, Y], Z) + g(\nabla_Y X, Z) \\ &= g([X, Y], Z) + Y(g(X, Z)) - g(X, \nabla_Y Z), \end{aligned}$$

$$\begin{aligned} 0 &= -Z(g(X, Y)) + g(\nabla_Z X, Y) + g(X, \nabla_Z Y) \\ &= -Z(g(X, Y)) + g(\nabla_X Z + [Z, X], Y) + g(X, \nabla_Y Z - [Y, Z]). \end{aligned}$$

By adding these relations we yield the so called **Koszul formula**

$$\begin{aligned} 2 \cdot g(\nabla_X Y, Z) &= \{X(g(Y, Z)) + Y(g(Z, X)) - Z(g(X, Y)) \\ &\quad + g(Z, [X, Y]) + g(Y, [Z, X]) - g(X, [Y, Z])\}. \end{aligned}$$

If $\{E_1, \dots, E_m\}$ is a local orthonormal frame for the tangent bundle then

$$\nabla_X Y = \sum_{i=1}^m g(\nabla_X Y, E_i) E_i.$$

Hence it is a direct consequence the Koszul formula that there exists **at most** one metric and torsion-free connection on the tangent bundle TM of (M, g) .

Definition 6.5. Let (M, g) be a Riemannian manifold then the operator

$$\nabla : C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(TM)$$

given by

$$\begin{aligned} g(\nabla_X Y, Z) &= \frac{1}{2} \{X(g(Y, Z)) + Y(g(X, Z)) - Z(g(X, Y)) \\ &\quad + g([Z, X], Y) + g([Z, Y], X) + g(Z, [X, Y])\} \end{aligned}$$

is called the **Levi-Civita connection** on M .

Remark 6.6. It is very important to note that the Levi-Civita connection is an intrinsic object on (M, g) i.e. only depending on the differentiable structure of the manifold and its Riemannian metric.

Proposition 6.7. Let (M, g) be a Riemannian manifold. Then the Levi-Civita connection ∇ is a connection on the tangent bundle TM of M .

PROOF. It follows from Definition 3.6, Theorem 4.22 and the fact that g is a tensor field that

$$g(\nabla_X (\lambda \cdot Y_1 + \mu \cdot Y_2), Z) = \lambda \cdot g(\nabla_X Y_1, Z) + \mu \cdot g(\nabla_X Y_2, Z)$$

and

$$g(\nabla_{Y_1 + Y_2} X, Z) = g(\nabla_{Y_1} X, Z) + g(\nabla_{Y_2} X, Z)$$

for all $\lambda, \mu \in \mathbb{R}$ and $X, Y_1, Y_2, Z \in C^\infty(TM)$. Furthermore we have for all $f \in C^\infty(M)$

$$\begin{aligned}
& 2 \cdot g(\nabla_X fY, Z) \\
&= \{X(f \cdot g(Y, Z)) + f \cdot Y(g(X, Z)) - Z(f \cdot g(X, Y)) \\
&\quad + f \cdot g([Z, X], Y) + g([Z, f \cdot Y], X) + g(Z, [X, f \cdot Y])\} \\
&= \{X(f) \cdot g(Y, Z) + f \cdot X(g(Y, Z)) + f \cdot Y(g(X, Z)) \\
&\quad - Z(f) \cdot g(X, Y) - f \cdot Z(g(X, Y)) + f \cdot g([Z, X], Y) \\
&\quad + g(Z(f) \cdot Y + f \cdot [Z, Y], X) + g(Z, X(f) \cdot Y + f \cdot [X, Y])\} \\
&= 2 \cdot \{X(f) \cdot g(Y, Z) + f \cdot g(\nabla_X Y, Z)\} \\
&= 2 \cdot g(X(f) \cdot Y + f \cdot \nabla_X Y, Z)
\end{aligned}$$

and

$$\begin{aligned}
& 2 \cdot g(\nabla_f \cdot_X Y, Z) \\
&= \{f \cdot X(g(Y, Z)) + Y(f \cdot g(X, Z)) - Z(f \cdot g(X, Y)) \\
&\quad + g([Z, f \cdot X], Y) + f \cdot g([Z, Y], X) + g(Z, [f \cdot X, Y])\} \\
&= \{f \cdot X(g(Y, Z)) + Y(f) \cdot g(X, Z) + f \cdot Y(g(X, Z)) \\
&\quad - Z(f) \cdot g(X, Y) - f \cdot Z(g(X, Y)) \\
&\quad + g(Z(f) \cdot X, Y) + f \cdot g([Z, X], Y) \\
&\quad + f \cdot g([Z, Y], X) + f \cdot g(Z, [X, Y]) - g(Z, Y(f) \cdot X)\} \\
&= 2 \cdot f \cdot g(\nabla_X Y, Z).
\end{aligned}$$

This proves that ∇ is a connection on the tangent bundle (TM, M, π) . \square

The next result is called the **fundamental theorem** of Riemannian geometry.

Theorem 6.8. *Let (M, g) be a Riemannian manifold. Then the Levi-Civita connection is the unique metric and torsion-free connection on the tangent bundle (TM, M, π) .*

PROOF. The difference $g(\nabla_X Y, Z) - g(\nabla_Y X, Z)$ equals twice the skew-symmetric part (w.r.t the pair (X, Y)) of the right hand side of the equation in Definition 6.5. This implies that

$$\begin{aligned}
g(\nabla_X Y, Z) - g(\nabla_Y X, Z) &= \frac{1}{2} \{g(Z, [X, Y]) - g(Z, [Y, X])\} \\
&= g([X, Y], Z).
\end{aligned}$$

This proves that the Levi-Civita connection is torsion-free.

The sum $g(\nabla_X Y, Z) + g(\nabla_X Z, Y)$ equals twice the symmetric part (w.r.t the pair (Y, Z)) on the right hand side of Definition 6.5. This yields

$$\begin{aligned} g(\nabla_X Y, Z) + g(Y, \nabla_X Z) &= \frac{1}{2} \{X(g(Y, Z)) + X(g(Z, Y))\} \\ &= X(g(Y, Z)). \end{aligned}$$

This shows that the Levi-Civita connection is compatible with the Riemannian metric g on M . The stated result now follows from Proposition 6.7. \square

For later use we introduce the following useful notion.

Definition 6.9. Let $X \in C^\infty(TM)$ be a vector field on (M, g) . Then the **first order covariant derivative**

$$\nabla_X : C^\infty(TM) \rightarrow C^\infty(TM)$$

in the direction of X is given by

$$\nabla_X : Y \mapsto \nabla_X Y.$$

In Lie theory, the following linear operator is called the adjoint representation of the Lie algebra \mathfrak{g} of G .

Definition 6.10. Let G be a Lie group. For a left-invariant vector field $Z \in \mathfrak{g}$ we define the linear operator $\text{ad}_Z : \mathfrak{g} \rightarrow \mathfrak{g}$ by

$$\text{ad}_Z : X \mapsto [Z, X].$$

For the classical compact groups, introduced in Chapter 2, we have the following interesting result.

Proposition 6.11. *Let G be one of the classical compact Lie groups, $\mathbf{O}(m)$, $\mathbf{SO}(m)$, $\mathbf{U}(m)$ or $\mathbf{SU}(m)$, equipped with its left-invariant metric given by*

$$g(X, Y) = \text{Re}(\text{trace}(\bar{X}^t \cdot Y)).$$

If $Z \in \mathfrak{g}$ is a left-invariant vector field on G then the linear operator $\text{ad}_Z : \mathfrak{g} \rightarrow \mathfrak{g}$ is skew symmetric.

PROOF. See Exercise 6.2. \square

The following result shows that the Koszul formula simplifies considerably in the important case when the manifold is a Riemannian Lie group.

Proposition 6.12. *Let (G, g) be a Lie group equipped with a left-invariant metric. Then its Levi-Civita connection ∇ satisfies*

$$g(\nabla_X Y, Z) = \frac{1}{2} \{g([X, Y], Z) + g(\text{ad}_Z(X), Y) + g(X, \text{ad}_Z(Y))\}$$

for all $X, Y, Z \in \mathfrak{g}$. In particular, if for all $Z \in \mathfrak{g}$ the map ad_Z is skew symmetric with respect to the Riemannian metric g then

$$\nabla_X Y = \frac{1}{2} [X, Y].$$

PROOF. See Exercise 6.3. \square

The next example shows how the Levi-Civita connection can be presented by means of local coordinates. Hopefully this will convince the reader that those should be avoided whenever possible.

Example 6.13. Let (M, g) be a Riemannian manifold with Levi-Civita connection ∇ . Further let (U, x) be a chart on M and put $X_i = \partial/\partial x_i \in C^\infty(TU)$. Then $\{X_1, \dots, X_m\}$ is a local frame of TM on U . For (U, x) we define the **Christoffel symbols** $\Gamma_{ij}^k : U \rightarrow \mathbb{R}$ of the connection ∇ with respect to (U, x) by

$$\sum_{k=1}^m \Gamma_{ij}^k \cdot X_k = \nabla_{X_i} X_j.$$

On the subset $x(U)$ of \mathbb{R}^m we define the Riemannian metric \tilde{g} by

$$\tilde{g}(e_i, e_j) = g_{ij} = g(X_i, X_j).$$

The differential dx is bijective so Proposition 4.27 implies that

$$dx([X_i, X_j]) = [dx(X_i), dx(X_j)] = [\partial_{e_i}, \partial_{e_j}] = 0$$

and hence $[X_i, X_j] = 0$. It now follows from the definition of the Christoffel symbols and the Koszul formula that for each $l = 1, 2, \dots, m$ we have

$$\begin{aligned} \sum_{k=1}^m g_{kl} \cdot \Gamma_{ij}^k &= \sum_{k=1}^m \langle X_k, X_l \rangle \cdot \Gamma_{ij}^k \\ &= \left\langle \sum_{k=1}^m \Gamma_{ij}^k \cdot X_k, X_l \right\rangle \\ &= \langle \nabla_{X_i} X_j, X_l \rangle \\ &= \frac{1}{2} \{X_i \langle X_j, X_l \rangle + X_j \langle X_l, X_i \rangle - X_l \langle X_i, X_j \rangle\} \\ &= \frac{1}{2} \left\{ \frac{\partial g_{jl}}{\partial x_i} + \frac{\partial g_{li}}{\partial x_j} - \frac{\partial g_{ij}}{\partial x_l} \right\}. \end{aligned}$$

Hence for each pair (i, j) we have a system of m linear equations in the m variables Γ_{ij}^k where $k = 1, 2, \dots, m$. Because the metric g is definite we can solve this as follows: Let $g^{kl} = (g^{-1})_{kl}$ be the components of the the inverse g^{-1} of g then the Christoffel symbols Γ_{ij}^k satisfy

$$\Gamma_{ij}^k = \frac{1}{2} \sum_{l=1}^m g^{kl} \left\{ \frac{\partial g_{jl}}{\partial x_i} + \frac{\partial g_{li}}{\partial x_j} - \frac{\partial g_{ij}}{\partial x_l} \right\}.$$

We will now study the relation between the Levi-Civita connection of a Riemannian manifold and that of its submanifolds, see Theorem 6.20. For this we need the following.

Definition 6.14. Let (N, h) be a Riemannian manifold and M be a submanifold equipped with the induced metric. Further let $\tilde{X} \in C^\infty(TM)$ be a vector field on M and $\tilde{Y} \in C^\infty(NM)$ be a section of its normal bundle. Let U be an open subset of N such that $U \cap M \neq \emptyset$. Two vector fields $X, Y \in C^\infty(TU)$ are said to be **local extensions** of \tilde{X} and \tilde{Y} to U if $\tilde{X}_p = X_p$ and $\tilde{Y}_p = Y_p$ for all $p \in U \cap M$. If $U = N$ then X, Y are said to be **global extension**.

Fact 6.15. Let (N, h) be a Riemannian manifold and M be a submanifold equipped with the induced metric, $\tilde{X} \in C^\infty(TM)$, $\tilde{Y} \in C^\infty(NM)$ and $p \in M$. Then there exists an open neighbourhood U of N containing p and $X, Y \in C^\infty(TU)$ extending \tilde{X} and \tilde{Y} on U , respectively.

Remark 6.16. Let (N, h) be a Riemannian manifold and M be a submanifold equipped with the induced metric. Let $Z \in C^\infty(TN)$ be a vector field on N and $\tilde{Z} = Z|_M : M \rightarrow TN$ be the restriction of Z to M . Note that \tilde{Z} is not necessarily an element of $C^\infty(TM)$ i.e. a vector field on the submanifold M . For each $p \in M$ the tangent vector $\tilde{Z}_p \in T_p N$ can be decomposed

$$\tilde{Z}_p = \tilde{Z}_p^\top + \tilde{Z}_p^\perp$$

in a unique way into its tangential part $(\tilde{Z}_p)^\top \in T_p M$ and its normal part $(\tilde{Z}_p)^\perp \in N_p M$. For this we write $\tilde{Z} = \tilde{Z}^\top + \tilde{Z}^\perp$.

Proposition 6.17. Let (N, h) be a Riemannian manifold and M be a submanifold equipped with the induced metric. If $Z \in C^\infty(TN)$ is a vector field on N then the sections \tilde{Z}^\top of the tangent bundle TM and \tilde{Z}^\perp of the normal bundle NM are smooth.

PROOF. See Exercise 6.7. □

Remark 6.18. Let $\tilde{X}, \tilde{Y} \in C^\infty(TM)$ be vector fields on M and $p \in M$. Let $X, Y \in C^\infty(TU)$ extend \tilde{X}, \tilde{Y} on an open neighbourhood U of p in N . It will be shown in Remark 7.3 that $(\nabla_X Y)_p$ only depends on the value $X_p = \tilde{X}_p$ and the value of Y along some curve $\gamma : (-\epsilon, \epsilon) \rightarrow N$ such that $\gamma(0) = p$ and $\dot{\gamma}(0) = X_p = \tilde{X}_p$. Since $X_p \in T_p M$ we may choose the curve γ such that the image $\gamma((-\epsilon, \epsilon))$ is contained in M . Then $\tilde{Y}_{\gamma(t)} = Y_{\gamma(t)}$ for $t \in (-\epsilon, \epsilon)$. This means that $(\nabla_X Y)_p$ only depends on \tilde{X}_p and the value of \tilde{Y} along γ , hence independent of how the vector fields \tilde{X} and \tilde{Y} are extended.

Remark 6.18 shows that the following important operators $\tilde{\nabla}$ and B are well defined.

Definition 6.19. Let (N, h) be a Riemannian manifold and M be a submanifold with the induced metric. Then we define the operators

$$\tilde{\nabla} : C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(TM)$$

and

$$B : C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(NM)$$

by

$$\tilde{\nabla}_{\tilde{X}} \tilde{Y} = (\nabla_X Y)^\top \quad \text{and} \quad B(\tilde{X}, \tilde{Y}) = (\nabla_X Y)^\perp.$$

Here X and Y are some local extensions of $\tilde{X}, \tilde{Y} \in C^\infty(TM)$. The operator B is called the **second fundamental form** of M in (N, h) .

The next result provides us with the relationship between the Levi-Civita connection of a Riemannian manifold and that of its submanifolds.

Theorem 6.20. *Let (N, h) be a Riemannian manifold and M be a submanifold with the induced metric g . Then the operator*

$$\tilde{\nabla} : C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(TM),$$

given by

$$\tilde{\nabla}_{\tilde{X}} \tilde{Y} = (\nabla_X Y)^\top,$$

is the Levi-Civita connection of the submanifold (M, g) .

PROOF. See Exercise 6.8. □

The second fundamental form of a submanifold of a Riemannian manifold has the following important properties.

Proposition 6.21. *Let (N, h) be a Riemannian manifold and M be a submanifold with the induced metric. Then the second fundamental form B of M in (N, h) is symmetric and tensorial in both its arguments.*

PROOF. See Exercise 6.9. \square

We now introduce the important notion of minimal submanifolds of a Riemannian manifold.

Definition 6.22. Let (N, h) be a Riemannian manifold and M be a submanifold with the induced metric. Then M is said to be **minimal** if its second fundamental form

$$B : C^\infty(TM) \otimes C^\infty(TM) \rightarrow C^\infty(NM)$$

is traceless i.e.

$$\text{trace } B = \sum_{k=1}^m B(\tilde{X}_k, \tilde{X}_k) = 0.$$

Here $\{\tilde{X}_1, \tilde{X}_2, \dots, \tilde{X}_m\}$ is any local orthonormal frame for the tangent bundle TM .

Example 6.23. Let us now consider the classical situation of a regular surface Σ as a submanifold of the three dimensional Euclidean space \mathbb{R}^3 . Let $\{\tilde{X}, \tilde{Y}\}$ be a local orthonormal frame for the tangent bundle $T\Sigma$ of Σ around a point $p \in \Sigma$ and \tilde{N} be the local Gauss map with $\tilde{N} = \tilde{X} \times \tilde{Y}$. If X, Y, N are local extensions of $\tilde{X}, \tilde{Y}, \tilde{N}$, forming a local orthonormal frame for $T\mathbb{R}^3$, then the second fundamental form B of Σ in \mathbb{R}^3 satisfies

$$\begin{aligned} B(\tilde{X}, \tilde{Y}) &= (\partial_X Y)^\perp \\ &= \langle \partial_X Y, N \rangle N \\ &= -\langle Y, \partial_X N \rangle N \\ &= -\langle Y, dN(X) \rangle N \\ &= \langle \tilde{Y}, S_p(\tilde{X}) \rangle \tilde{N}, \end{aligned}$$

where $S_p : T_p\Sigma \rightarrow T_p\Sigma$ is the shape operator at p . Then the trace of B satisfies

$$\begin{aligned} \text{trace } B &= (\langle S_p(\tilde{X}), \tilde{X} \rangle + \langle S_p(\tilde{Y}), \tilde{Y} \rangle) \tilde{N} \\ &= (\text{trace } S_p) \tilde{N} \\ &= (k_1 + k_2) \tilde{N}. \end{aligned}$$

Here k_1 and k_2 are the eigenvalues of the symmetric shape operator S_p i.e. the principal curvatures at p . This shows that the surface Σ is a minimal submanifold of \mathbb{R}^3 if and only if the classical **mean curvature** $H = (k_1 + k_2)/2$ vanishes.

We conclude this chapter by observing that the Levi-Civita connection of a Riemannian manifold (N, h) induces a metric connection $\bar{\nabla}$ on the normal bundle NM of its submanifold M as follows.

Proposition 6.24. *Let (N, h) be a Riemannian manifold and M be a submanifold with the induced metric. Then the operator*

$$\bar{\nabla} : C^\infty(TM) \times C^\infty(NM) \rightarrow C^\infty(NM)$$

given by

$$\bar{\nabla}_{\tilde{X}} \tilde{Y} = (\nabla_X Y)^\perp$$

*is a well defined connection on the normal bundle NM . Here X and Y are some local extensions of $\tilde{X} \in C^\infty(TM)$ and $\tilde{Y} \in C^\infty(NM)$, respectively. Furthermore, the connection $\bar{\nabla}$ is **metric** i.e. it satisfies*

$$\tilde{X}(h(\tilde{Y}, \tilde{Z})) = h(\bar{\nabla}_{\tilde{X}} \tilde{Y}, \tilde{Z}) + h(\tilde{Y}, \bar{\nabla}_{\tilde{X}} \tilde{Z}),$$

for all $\tilde{X} \in C^\infty(TM)$ and $\tilde{Y}, \tilde{Z} \in C^\infty(NM)$.

PROOF. See Exercise 6.10. □

Exercises

Exercise 6.1. Let M be a smooth manifold and $\hat{\nabla}$ be a connection on the tangent bundle (TM, M, π) . Prove that the torsion of $\hat{\nabla}$

$$T : C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(TM),$$

given by

$$T(X, Y) = \hat{\nabla}_X Y - \hat{\nabla}_Y X - [X, Y],$$

is a tensor field of type $(2, 1)$.

Exercise 6.2. Find a proof of Proposition 6.11.

Exercise 6.3. Find a proof of Proposition 6.12.

Exercise 6.4. Let Sol be the 3-dimensional subgroup of $\mathbf{SL}_3(\mathbb{R})$ given by

$$\text{Sol} = \left\{ \begin{pmatrix} e^z & 0 & x \\ 0 & e^{-z} & y \\ 0 & 0 & 1 \end{pmatrix} \mid p = (x, y, z) \in \mathbb{R}^3 \right\}.$$

Let $X, Y, Z \in \mathfrak{g}$ be left-invariant vector fields on Sol such that

$$X_e = \frac{\partial}{\partial x}|_{p=0}, \quad Y_e = \frac{\partial}{\partial y}|_{p=0} \quad \text{and} \quad Z_e = \frac{\partial}{\partial z}|_{p=0}.$$

Show that

$$[X, Y] = 0, \quad [Z, X] = X \quad \text{and} \quad [Z, Y] = -Y.$$

Let g be the left-invariant Riemannian metric on G such that $\{X, Y, Z\}$ is an orthonormal basis for the Lie algebra \mathfrak{g} . Calculate the following vector fields:

$$\nabla_X Y, \quad \nabla_Y X, \quad \nabla_X Z, \quad \nabla_Z X, \quad \nabla_Y Z \quad \text{and} \quad \nabla_Z Y.$$

Exercise 6.5. Let $\mathbf{SO}(m)$ be the special orthogonal group equipped with the metric

$$\langle X, Y \rangle = \frac{1}{2} \text{trace}(X^t \cdot Y).$$

Prove that \langle, \rangle is left-invariant and that for left-invariant vector fields $X, Y \in \mathfrak{so}(m)$ we have

$$\nabla_X Y = \frac{1}{2} [X, Y].$$

Let A, B, C be the elements of the Lie algebra $\mathfrak{so}(3)$ with

$$A_e = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad B_e = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad C_e = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Prove that $\{A, B, C\}$ is an orthonormal basis for $\mathfrak{so}(3)$ and calculate

$$\nabla_A B, \quad \nabla_B C \quad \text{and} \quad \nabla_C A.$$

Exercise 6.6. Let $\mathbf{SL}_2(\mathbb{R})$ be the real special linear group equipped with the metric

$$\langle X, Y \rangle_p = \frac{1}{2} \text{trace}((p^{-1}X)^t(p^{-1}Y)).$$

Let A, B, C be the elements of the Lie algebra $\mathfrak{sl}_2(\mathbb{R})$ with

$$A_e = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad B_e = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad C_e = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Prove that $\{A, B, C\}$ is an orthonormal basis for $\mathfrak{sl}_2(\mathbb{R})$ and calculate

$$\nabla_A B, \quad \nabla_B C \quad \text{and} \quad \nabla_C A.$$

Exercise 6.7. Find a proof of Proposition 6.17.

Exercise 6.8. Find a proof of Theorem 6.20.

Exercise 6.9. Find a proof of Proposition 6.21.

Exercise 6.10. Find a proof of Proposition 6.24.

CHAPTER 7

Geodesics

The main purpose of this chapter is to introduce the important notion of a geodesic in a Riemannian manifold. Geodesics are solutions to a second order system of non-linear ordinary differential equations, depending on the geometry of the manifold. We develop the idea of a parallel vector field along a given curve in the manifold. We show that geodesics are solutions to two different variational problems. They are both critical points of the so called energy functional and locally the shortest paths between their endpoints. We then study totally geodesic submanifolds.

Definition 7.1. Let M be a smooth manifold and (TM, M, π) be its tangent bundle. A **vector field X along a curve $\gamma : I \rightarrow M$** is a curve $X : I \rightarrow TM$ such that $\pi \circ X = \gamma$. By $C_\gamma^\infty(TM)$ we denote the set of all smooth vector fields along γ . For $X, Y \in C_\gamma^\infty(TM)$ and $f \in C^\infty(I)$ we define the addition $+$ and the multiplication \cdot by

- (i) $(X + Y)(t) = X(t) + Y(t)$,
- (ii) $(f \cdot X)(t) = f(t) \cdot X(t)$.

This turns $(C_\gamma^\infty(TM), +, \cdot)$ into a module over $C^\infty(I)$ and a real vector space over the constant functions in particular. For a given smooth curve $\gamma : I \rightarrow M$ in M the smooth vector field $X : I \rightarrow TM$ with $X : t \mapsto (\gamma(t), \dot{\gamma}(t))$ is called the **tangent field** along γ .

The next result provides us with a differential operator for vector fields along a given curve and shows how this is closely related to the Levi-Civita connection.

Proposition 7.2. *Let (M, g) be a smooth Riemannian manifold and $\gamma : I \rightarrow M$ be a curve in M . Then there exists a unique operator*

$$\frac{D}{dt} : C_\gamma^\infty(TM) \rightarrow C_\gamma^\infty(TM)$$

such that for all $\lambda, \mu \in \mathbb{R}$ and $f \in C^\infty(I)$,

- (i) $D(\lambda \cdot X + \mu \cdot Y)/dt = \lambda \cdot (DX/dt) + \mu \cdot (DY/dt)$,
- (ii) $D(f \cdot Y)/dt = df/dt \cdot Y + f \cdot (DY/dt)$, and

- (iii) for each $t_0 \in I$ there exists an open subinterval J of I such that $t_0 \in J$ and if $X \in C^\infty(TM)$ is a vector field with $X_{\gamma(t)} = Y(t)$ for all $t \in J$ then

$$\left(\frac{DY}{dt}\right)(t_0) = (\nabla_{\dot{\gamma}} X)_{\gamma(t_0)}.$$

PROOF. Let us first prove the uniqueness, so for the moment we assume that such an operator exists. For a point $t_0 \in I$ choose a chart (U, x) on M and an open subinterval $J \subset I$ such that $t_0 \in J$, $\gamma(J) \subset U$ and put $X_i = \partial/\partial x_i \in C^\infty(TU)$. Then any vector field Y along the restriction of γ to J can be written in the form

$$Y(t) = \sum_{j=1}^m \alpha_j(t) \cdot (X_j)_{\gamma(t)}$$

for some functions $\alpha_j \in C^\infty(J)$. The conditions (i) and (ii) imply that

$$(1) \quad \left(\frac{DY}{dt}\right)(t) = \sum_{k=1}^m \dot{\alpha}_k(t) \cdot (X_k)_{\gamma(t)} + \sum_{j=1}^m \alpha_j(t) \cdot \left(\frac{DX_j}{dt}\right)_{\gamma(t)}.$$

For the local chart (U, x) , the composition

$$x \circ \gamma(t) = (\gamma_1(t), \dots, \gamma_m(t)) = \sum_{i=1}^m \gamma_i(t) \cdot e_i$$

parametrises a curve in \mathbb{R}^m contained in $x(U)$. Hence the derivative satisfies

$$dx_{\gamma(t)}(\dot{\gamma}(t)) = \frac{d}{dt}(x \circ \gamma(t)) = (\dot{\gamma}_1(t), \dots, \dot{\gamma}_m(t)).$$

Because the local coordinate is a diffeomorphism, its linear differential $dx : TU \rightarrow T\mathbb{R}^m$ is bijective satisfying

$$dx\left(\frac{\partial}{\partial x_i}\right) = e_i,$$

for $i = 1, \dots, m$. This immediately implies that

$$\dot{\gamma}(t) = \sum_{i=1}^m \dot{\gamma}_i(t) \cdot (X_i)_{\gamma(t)}$$

and the condition (iii) shows that

$$(2) \quad \left(\frac{DX_j}{dt}\right)_{\gamma(t)} = (\nabla_{\dot{\gamma}} X_j)_{\gamma(t)} = \sum_{i=1}^m \dot{\gamma}_i(t) \cdot (\nabla_{X_i} X_j)_{\gamma(t)}.$$

Together equations (1) and (2) give

$$(3) \quad \left(\frac{DY}{dt}\right)(t) = \sum_{k=1}^m \left\{ \dot{\alpha}_k(t) + \sum_{i,j=1}^m \alpha_j(t) \cdot \dot{\gamma}_i(t) \cdot \Gamma_{ij}^k(\gamma(t)) \right\} \cdot (X_k)_{\gamma(t)}.$$

This shows that there exists **at most** one such differential operator.

It is easily seen that if we use equation (3) for defining an operator D/dt then this satisfies the necessary conditions of Proposition 7.2. This proves the existence part of the stated result. \square

The equation (3) has the following important consequence.

Remark 7.3. It follows from the fact that the Levi-Civita connection is tensorial in its first argument i.e.

$$\nabla_f \cdot Z X = f \cdot \nabla_Z X$$

and the equation

$$(\nabla_{\dot{\gamma}} X)_{\gamma(t_0)} = \left(\frac{DY}{dt}\right)(t_0)$$

in Proposition 7.2 that the value $(\nabla_Z X)_p$ of $\nabla_Z X$ at p only depends on the value of Z_p of Z at p and the values of Y along some curve γ satisfying $\gamma(0) = p$ and $\dot{\gamma}(0) = Z_p$. This allows us to use the notation $\nabla_{\dot{\gamma}} Y$ for DY/dt .

The Levi-Civita connection can now be used to define the notions of parallel vector fields and geodesics on a Riemannian manifold. We will show that they are solutions to ordinary differential equations.

Definition 7.4. Let (M, g) be a Riemannian manifold and let $\gamma : I \rightarrow M$ be a C^1 -curve. A vector field X along γ is said to be **parallel** if

$$\nabla_{\dot{\gamma}} X = 0.$$

A C^2 -curve $\gamma : I \rightarrow M$ is said to be a **geodesic** if its tangent field $\dot{\gamma}$ is parallel along γ i.e.

$$\nabla_{\dot{\gamma}} \dot{\gamma} = 0.$$

The next result shows that for a given initial value at a point we yield a parallel vector field globally defined along any curve through that point.

Theorem 7.5. Let (M, g) be a Riemannian manifold and $I = (a, b)$ be an open interval on the real line \mathbb{R} . Further let $\gamma : [a, b] \rightarrow M$ be a continuous curve which is C^1 on I , $t_0 \in I$ and $v \in T_{\gamma(t_0)} M$. Then there exists a unique parallel vector field Y along γ such that $Y(t_0) = v$.

PROOF. Let (U, x) be a chart on M such that $\gamma(t_0) \in U$ and put $X_i = \partial/\partial x_i \in C^\infty(TU)$. Let J be an open subset of I such that the image $\gamma(J)$ is contained in U . Then the tangent of the restriction of γ to J can be written as

$$\dot{\gamma}(t) = \sum_{i=1}^m \dot{\gamma}_i(t) \cdot (X_i)_{\gamma(t)}.$$

Similarly, let Y be a vector field along γ represented by

$$Y(t) = \sum_{j=1}^m \alpha_j(t) \cdot (X_j)_{\gamma(t)}.$$

Then

$$\begin{aligned} (\nabla_{\dot{\gamma}} Y)(t) &= \sum_{j=1}^m \{ \dot{\alpha}_j(t) \cdot (X_j)_{\gamma(t)} + \alpha_j(t) \cdot (\nabla_{\dot{\gamma}} X_j)_{\gamma(t)} \} \\ &= \sum_{k=1}^m \left\{ \dot{\alpha}_k(t) + \sum_{i,j=1}^m \alpha_j(t) \cdot \dot{\gamma}_i(t) \cdot \Gamma_{ij}^k(\gamma(t)) \right\} (X_k)_{\gamma(t)}. \end{aligned}$$

This implies that the vector field Y is parallel i.e. $\nabla_{\dot{\gamma}} Y \equiv 0$ if and only if the following first order **linear** system of ordinary differential equations is satisfied

$$\dot{\alpha}_k(t) + \sum_{i,j=1}^m \alpha_j(t) \cdot \dot{\gamma}_i(t) \cdot \Gamma_{ij}^k(\gamma(t)) = 0,$$

for all $k = 1, \dots, m$. It follows from Fact 7.6 that to each initial value $\alpha(t_0) = (v_1, \dots, v_m) \in \mathbb{R}^m$, with

$$Y_0 = \sum_{k=1}^m v_k \cdot (X_k)_{\gamma(t_0)},$$

there exists a unique solution $\alpha = (\alpha_1, \dots, \alpha_m)$ to the above system. This gives us the unique parallel vector field Y

$$Y(t) = \sum_{k=1}^m \alpha_k(t) \cdot (X_k)_{\gamma(t)}$$

along J . Since the Christoffel symbols are bounded along the compact set $[a, b]$ it is clear that the parallel vector field can be extended to the whole of $I = (a, b)$. \square

The following result is the well-known theorem of Picard-Lindelöf.

Fact 7.6. *Let $f : U \rightarrow \mathbb{R}^n$ be a continuous map defined on an open subset U of $\mathbb{R} \times \mathbb{R}^n$ and $L \in \mathbb{R}^+$ such that*

$$|f(t, y_1) - f(t, y_2)| \leq L \cdot |y_1 - y_2|$$

for all $(t, y_1), (t, y_2) \in U$. If $(t_0, x_0) \in U$ then there exists a unique local solution $x : I \rightarrow \mathbb{R}^n$ to the following initial value problem

$$x'(t) = f(t, x(t)), \quad x(t_0) = x_0.$$

For parallel vector fields we have the following important result.

Lemma 7.7. *Let (M, g) be a Riemannian manifold, $\gamma : I \rightarrow M$ be a C^1 -curve and X, Y be parallel vector fields along γ . Then the function $g(X, Y) : I \rightarrow \mathbb{R}$ given by*

$$g(X, Y) : t \mapsto g_{\gamma(t)}(X_{\gamma(t)}, Y_{\gamma(t)})$$

is constant. In particular, if γ is a geodesic then $g(\dot{\gamma}, \dot{\gamma})$ is constant along γ .

PROOF. Using the fact that the Levi-Civita connection is metric we obtain

$$\frac{d}{dt}(g(X, Y)) = g(\nabla_{\dot{\gamma}} X, Y) + g(X, \nabla_{\dot{\gamma}} Y) = 0.$$

This proves that the function $g(X, Y)$ is constant along γ . □

The following result is a very useful tool. We will employ this in Chapter 9.

Proposition 7.8. *Let (M, g) be a Riemannian manifold, $p \in M$ and $\{v_1, \dots, v_m\}$ be an orthonormal basis for the tangent space $T_p M$. Let $\gamma : I \rightarrow M$ be a C^1 -curve such that $\gamma(0) = p$ and X_1, \dots, X_m be parallel vector fields along γ such that $X_k(0) = v_k$ for $k = 1, 2, \dots, m$. Then the set $\{X_1(t), \dots, X_m(t)\}$ is a orthonormal basis for the tangent space $T_{\gamma(t)} M$ for all $t \in I$.*

PROOF. This is a direct consequence of Lemma 7.7. □

Geodesics play a very important role in Riemannian geometry. For these we have the following fundamental existence and uniqueness result.

Theorem 7.9. *Let (M, g) be a Riemannian manifold. If $p \in M$ and $v \in T_p M$ then there exists an open interval $I = (-\epsilon, \epsilon)$ and a unique geodesic $\gamma : I \rightarrow M$ such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v$.*

PROOF. Let (U, x) be a chart on M such that $p \in U$ and put $X_i = \partial/\partial x_i \in C^\infty(TU)$. Let J be an open subset of I such that the image $\gamma(J)$ is contained in U . Then the tangent of the restriction of γ to J can be written as

$$\dot{\gamma}(t) = \sum_{i=1}^m \dot{\gamma}_i(t) \cdot (X_i)_{\gamma(t)}.$$

By differentiation we then obtain

$$\begin{aligned} \nabla_{\dot{\gamma}} \dot{\gamma} &= \sum_{j=1}^m \nabla_{\dot{\gamma}} (\dot{\gamma}_j(t) \cdot (X_j)_{\gamma(t)}) \\ &= \sum_{j=1}^m \{ \ddot{\gamma}_j(t) \cdot (X_j)_{\gamma(t)} + \sum_{i=1}^m \dot{\gamma}_i(t) \cdot \dot{\gamma}_j(t) \cdot (\nabla_{X_i} X_j)_{\gamma(t)} \} \\ &= \sum_{k=1}^m \{ \ddot{\gamma}_k(t) + \sum_{i,j=1}^m \dot{\gamma}_i(t) \cdot \dot{\gamma}_j(t) \cdot \Gamma_{ij}^k(\gamma(t)) \} \cdot (X_k)_{\gamma(t)}. \end{aligned}$$

Hence the curve γ is a geodesic if and only if

$$\ddot{\gamma}_k(t) + \sum_{i,j=1}^m \dot{\gamma}_i(t) \cdot \dot{\gamma}_j(t) \cdot \Gamma_{ij}^k(\gamma(t)) = 0$$

for all $k = 1, \dots, m$. It follows from Fact 7.10 that for initial values $q = x(p)$ and $w = (dx)_p(v)$ there exists an open interval $(-\epsilon, \epsilon)$ and a unique solution $(\gamma_1, \dots, \gamma_m)$ satisfying the initial conditions

$$(\gamma_1(0), \dots, \gamma_m(0)) = q \quad \text{and} \quad (\dot{\gamma}_1(0), \dots, \dot{\gamma}_m(0)) = w.$$

□

The following result is a second order consequence of the well-known theorem of Picard-Lindelöf.

Fact 7.10. *Let $f : U \rightarrow \mathbb{R}^n$ be a continuous map defined on an open subset U of $\mathbb{R} \times \mathbb{R}^{2n}$ and $L \in \mathbb{R}^+$ such that*

$$|f(t, y_1) - f(t, y_2)| \leq L \cdot |y_1 - y_2|$$

for all $(t, y_1), (t, y_2) \in U$. If $(t_0, (x_0, x_1)) \in U$ and $x_0, x_1 \in \mathbb{R}^n$ then there exists a unique local solution $x : I \rightarrow \mathbb{R}^n$ to the following initial value problem

$$x''(t) = f(t, x(t), x'(t)), \quad x(t_0) = x_0, \quad x'(t_0) = x_1.$$

Remark 7.11. The Levi-Civita connection ∇ on a given Riemannian manifold (M, g) is an inner object i.e. completely determined by

the differentiable structure on M and the Riemannian metric g , see Remark 6.6. Hence the same applies for the condition

$$\nabla_{\dot{\gamma}} \dot{\gamma} = 0$$

for a given curve $\gamma : I \rightarrow M$. This means that the image of a geodesic under a local isometry is again a geodesic.

We can now determine the geodesics in the Euclidean spaces.

Example 7.12. Let $E^m = (\mathbb{R}^m, \langle \cdot, \cdot \rangle_{\mathbb{R}^m})$ be the Euclidean space of dimension m . For the trivial chart $\text{id}_{\mathbb{R}^m} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ the metric on E^m is given by $g_{ij} = \delta_{ij}$. As a direct consequence of Example 6.13 we see that the corresponding Christoffel symbols satisfy

$$\Gamma_{ij}^k = 0 \quad \text{for all } i, j, k = 1, \dots, m.$$

Hence $\gamma : I \rightarrow \mathbb{R}^m$ is a geodesic if and only if $\ddot{\gamma}(t) = 0$. For $p \in \mathbb{R}^m$ and $v \in T_p \mathbb{R}^m \cong \mathbb{R}^m$ define

$$\gamma_{(p,v)} : \mathbb{R} \rightarrow \mathbb{R}^m \quad \text{by} \quad \gamma_{(p,v)}(t) = p + t \cdot v.$$

Then $\gamma_{(p,v)}(0) = p$, $\dot{\gamma}_{(p,v)}(0) = v$ and $\ddot{\gamma}_{(p,v)} = 0$. It now follows from Theorem 7.9 that the geodesics in E^m are the straight lines.

For the classical situation of surfaces in Euclidean \mathbb{R}^3 we have the following.

Example 7.13. Let us now consider the classical situation of a regular surface Σ as a submanifold of the three dimensional Euclidean space \mathbb{R}^3 . If $\gamma : I \rightarrow \Sigma$ is a C^2 -curve, then Theorem 6.20 tells us that

$$\nabla_{\dot{\gamma}} \dot{\gamma} = (\partial_{\dot{\gamma}} \dot{\gamma})^\top = \ddot{\gamma}^\top.$$

This means that γ is a geodesic if and only if the tangential part $\ddot{\gamma}^\top$ of the second derivative $\ddot{\gamma}$ vanishes.

Definition 7.14. A geodesic $\gamma : J \rightarrow (M, g)$ in a Riemannian manifold is said to be **maximal** if it can not be extended to a geodesic defined on an interval I strictly containing J . The manifold (M, g) is said to be **complete** if for each point $(p, v) \in TM$ there exists a geodesic $\gamma : \mathbb{R} \rightarrow M$ defined on the whole of \mathbb{R} such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v$.

Proposition 7.15. Let (N, h) be a Riemannian manifold with Levi-Civita connection ∇ and M be a submanifold equipped with the induced metric g . A curve $\gamma : I \rightarrow M$ is a geodesic in M if and only if

$$(\nabla_{\dot{\gamma}} \dot{\gamma})^\top = 0.$$

PROOF. The result follows directly from Theorem 6.20 stating that the Levi-Civita connection $\tilde{\nabla}$ on (M, g) satisfies

$$\tilde{\nabla}_{\dot{\gamma}}\dot{\gamma} = (\nabla_{\dot{\gamma}}\dot{\gamma})^\top.$$

□

The geodesics on the standard spheres can be determined as follows.

Example 7.16. Let E^{m+1} be the Euclidean space of dimension $(m+1)$ and S^m be the unit sphere in E^{m+1} with the induced metric. At a point $p \in S^m$ the normal space $N_p S^m$ of S^m in E^{m+1} is simply the line generated by p . If $\gamma : I \rightarrow S^m$ is a C^2 -curve on the sphere, then

$$\tilde{\nabla}_{\dot{\gamma}}\dot{\gamma} = (\nabla_{\dot{\gamma}}\dot{\gamma})^\top = (\partial_{\dot{\gamma}}\dot{\gamma})^\top = \ddot{\gamma}^\top = \ddot{\gamma} - \ddot{\gamma}^\perp = \ddot{\gamma} - \langle \ddot{\gamma}, \gamma \rangle \gamma.$$

This shows that γ is a geodesic on the sphere S^m if and only if

$$(4) \quad \ddot{\gamma} = \langle \ddot{\gamma}, \gamma \rangle \gamma.$$

For a point $(p, X) \in TS^m$ define the curve $\gamma = \gamma_{(p,X)} : \mathbb{R} \rightarrow S^m$ by

$$\gamma : t \mapsto \begin{cases} p & \text{if } X = 0 \\ \cos(|X|t) \cdot p + \sin(|X|t) \cdot X/|X| & \text{if } X \neq 0. \end{cases}$$

Then one easily checks that $\gamma(0) = p$, $\dot{\gamma}(0) = X$ and that γ satisfies the geodesic equation (4). This shows that the non-constant geodesics on S^m are precisely the great circles and the sphere is complete.

Having determined the geodesics on the standard spheres, we can now easily find the geodesics on the real projective spaces.

Example 7.17. Let $\text{Sym}(\mathbb{R}^{m+1})$ be equipped with the metric

$$g(A, B) = \frac{1}{8} \text{trace}(A^t \cdot B).$$

Then we know from Example 5.25 that the map $\phi : S^m \rightarrow \text{Sym}(\mathbb{R}^{m+1})$ with

$$\phi : p \mapsto (2pp^t - e)$$

is an isometric immersion and that the image $\phi(S^m)$ is isometric to the m -dimensional real projective space $\mathbb{R}P^m$. This means that the geodesics on $\mathbb{R}P^m$ are exactly the images of geodesics on S^m . This shows that the real projective spaces are complete.

We will now show that the geodesics are critical points of the so called energy functional. For this we need the following two definitions.

Definition 7.18. Let (M, g) be a Riemannian manifold and $\gamma : I \rightarrow M$ be a C^r -curve on M . A **variation** of γ is a C^r -map

$$\Phi : (-\epsilon, \epsilon) \times I \rightarrow M$$

such that for all $s \in I$, $\Phi_0(s) = \Phi(0, s) = \gamma(s)$. If the interval is compact i.e. of the form $I = [a, b]$, then the variation Φ is called **proper** if for all $t \in (-\epsilon, \epsilon)$, $\Phi_t(a) = \gamma(a)$ and $\Phi_t(b) = \gamma(b)$.

Definition 7.19. Let (M, g) be a Riemannian manifold and $\gamma : I \rightarrow M$ be a C^2 -curve on M . For every compact interval $[a, b] \subset I$ we define the **energy functional** $E_{[a,b]}$ by

$$E_{[a,b]}(\gamma) = \frac{1}{2} \int_a^b g(\dot{\gamma}(t), \dot{\gamma}(t)) dt.$$

A C^2 -curve $\gamma : I \rightarrow M$ is called a **critical point** for the energy functional if every proper variation Φ of $\gamma|_{[a,b]}$ satisfies

$$\frac{d}{dt}(E_{[a,b]}(\Phi_t))|_{t=0} = 0.$$

We will now prove that geodesics can be characterised as the critical points of the energy functional.

Theorem 7.20. A C^2 -curve $\gamma : I = [a, b] \rightarrow M$ is a critical point for the energy functional if and only if it is a geodesic.

PROOF. For a C^2 -map $\Phi : (-\epsilon, \epsilon) \times I \rightarrow M$, $\Phi : (t, s) \mapsto \Phi(t, s)$ we define the vector fields $X = d\Phi(\partial/\partial s)$ and $Y = d\Phi(\partial/\partial t)$ along Φ . The following shows that the vector fields X and Y commute.

$$\begin{aligned} \nabla_X Y - \nabla_Y X &= [X, Y] \\ &= [d\Phi(\partial/\partial s), d\Phi(\partial/\partial t)] \\ &= d\Phi([\partial/\partial s, \partial/\partial t]) \\ &= 0, \end{aligned}$$

since $[\partial/\partial s, \partial/\partial t] = 0$. We now assume that Φ is a proper variation of γ . Then

$$\begin{aligned} \frac{d}{dt}(E_{[a,b]}(\Phi_t)) &= \frac{1}{2} \frac{d}{dt} \left(\int_a^b g(X, X) ds \right) \\ &= \frac{1}{2} \int_a^b \frac{d}{dt} (g(X, X)) ds \\ &= \int_a^b g(\nabla_Y X, X) ds \end{aligned}$$

$$\begin{aligned}
&= \int_a^b g(\nabla_X Y, X) ds \\
&= \int_a^b \left(\frac{d}{ds} (g(Y, X)) - g(Y, \nabla_X X) \right) ds \\
&= [g(Y, X)]_a^b - \int_a^b g(Y, \nabla_X X) ds.
\end{aligned}$$

The variation is proper, so $Y(t, a) = Y(t, b) = 0$. Furthermore

$$X(0, s) = \partial\Phi/\partial s(0, s) = \dot{\gamma}(s),$$

so

$$\frac{d}{dt}(E_{[a,b]}(\Phi_t))|_{t=0} = - \int_a^b g(Y(0, s), (\nabla_{\dot{\gamma}} \dot{\gamma})(s)) ds.$$

The last integral vanishes for every proper variation Φ of γ if and only if $\nabla_{\dot{\gamma}} \dot{\gamma} = 0$. \square

A geodesic $\gamma : I \rightarrow (M, g)$ is a special case of what is called a **harmonic map** $\phi : (M, g) \rightarrow (N, h)$ between Riemannian manifolds. Other examples are the conformal immersions $\psi : (M^2, g) \rightarrow (N, h)$ which parametrise the minimal surfaces in (N, h) . This study was initiated by the seminar paper: J. Eells, J. H. Sampson, *Harmonic Mappings of Riemannian Manifolds*, Amer. J. Math. **86**, (1964), 109-160. For a modern reference on harmonic maps see H. Urakawa, *Calculus of Variations and Harmonic Maps*, Translations of Mathematical Monographs **132**, AMS (1993).

Our next goal is to prove the important result of Theorem 7.22. For this we need to introduce the exponential map. This is a fundamental tool in Riemannian geometry.

Definition 7.21. Let (M^m, g) be an m -dimensional Riemannian manifold, $p \in M$ and

$$S_p^{m-1} = \{v \in T_p M \mid g_p(v, v) = 1\}$$

be the unit sphere in the tangent space $T_p M$ at p . Then every point $w \in T_p M \setminus \{0\}$ can be written as $w = r_w \cdot v_w$, where $r_w = |w|$ and $v_w = w/|w| \in S_p^{m-1}$. For $v \in S_p^{m-1}$ let $\gamma_v : (-\alpha_v, \beta_v) \rightarrow M$ be the maximal geodesic such that $\alpha_v, \beta_v \in \mathbb{R}^+ \cup \{\infty\}$, $\gamma_v(0) = p$ and $\dot{\gamma}_v(0) = v$. It can be shown that the real number

$$\epsilon_p = \inf\{\alpha_v, \beta_v \mid v \in S_p^{m-1}\}$$

is positive so the open ball

$$B_{\epsilon_p}^m(0) = \{v \in T_p M \mid g_p(v, v) < \epsilon_p^2\}$$

is non-empty. The **exponential map** $\exp_p : B_{\epsilon_p}^m(0) \rightarrow M$ at p is defined by

$$\exp_p : w \mapsto \begin{cases} p & \text{if } w = 0 \\ \gamma_{v_w}(r_w) & \text{if } w \neq 0. \end{cases}$$

Note that for $v \in S_p^{m-1}$ the line segment $\lambda_v : (-\epsilon_p, \epsilon_p) \rightarrow T_p M$ with $\lambda_v : t \mapsto t \cdot v$ is mapped onto the geodesic γ_v i.e. locally we have $\gamma_v = \exp_p \circ \lambda_v$. One can prove that the map \exp_p is differentiable and it follows from its definition that the differential

$$d(\exp_p)_0 : T_p M \rightarrow T_p M$$

is the identity map for the tangent space $T_p M$. Then the inverse mapping theorem tells us that there exists an $r_p \in \mathbb{R}^+$ such that if $U_p = B_{r_p}^m(0)$ and $V_p = \exp_p(U_p)$ then $\exp_p|_{U_p} : U_p \rightarrow V_p$ is a diffeomorphism parametrising the open subset V_p of M .

The next result shows that the geodesics are locally the shortest paths between their endpoints.

Theorem 7.22. *Let (M, g) be a Riemannian manifold. Then the geodesics are locally the shortest paths between their end points.*

PROOF. Let $p \in M$, $U = B_r^m(0)$ in $T_p M$ and $V = \exp_p(U)$ be such that the restriction

$$\phi = \exp_p|_U : U \rightarrow V$$

of the exponential map at p is a diffeomorphism. We define a metric \tilde{g} on U such that for each $X, Y \in C^\infty(TU)$ we have

$$\tilde{g}(X, Y) = g(d\phi(X), d\phi(Y)).$$

This turns $\phi : (U, \tilde{g}) \rightarrow (V, g)$ into an isometry. It then follows from the construction of the exponential map, that the geodesics in (U, \tilde{g}) through the point $0 = \phi^{-1}(p)$ are exactly the lines $\lambda_v : t \mapsto t \cdot v$ where $v \in T_p M$.

Now let $q \in B_r^m(0) \setminus \{0\}$ and $\lambda_q : [0, 1] \rightarrow B_r^m(0)$ be the curve $\lambda_q : t \mapsto t \cdot q$. Further let $\sigma : [0, 1] \rightarrow U$ be any C^1 -curve such that $\sigma(0) = 0$ and $\sigma(1) = q$. Along the curve σ we define the vector field X with $X : t \mapsto \dot{\sigma}(t)$ and the tangent field $\dot{\sigma} : t \mapsto \dot{\sigma}(t)$ to σ . Then the radial component $\dot{\sigma}_{\text{rad}}$ of $\dot{\sigma}$ is the orthogonal projection of $\dot{\sigma}$ onto the line generated by X i.e.

$$\dot{\sigma}_{\text{rad}} : t \mapsto \frac{\tilde{g}(\dot{\sigma}(t), X(t))}{\tilde{g}(X(t), X(t))} X(t).$$

Then it is easily checked that

$$|\dot{\sigma}_{\text{rad}}(t)| = \frac{|\tilde{g}(\dot{\sigma}(t), X(t))|}{|X(t)|}$$

and

$$\frac{d}{dt}|X(t)| = \frac{d}{dt}\sqrt{\tilde{g}(X(t), X(t))} = \frac{\tilde{g}(\dot{\sigma}(t), X(t))}{|X(t)|}.$$

Combining these two relations we yield

$$|\dot{\sigma}_{\text{rad}}(t)| \geq \frac{d}{dt}|X(t)|.$$

This means that

$$\begin{aligned} L(\sigma) &= \int_0^1 |\dot{\sigma}(t)| dt \\ &\geq \int_0^1 |\dot{\sigma}_{\text{rad}}(t)| dt \\ &\geq \int_0^1 \frac{d}{dt}|X(t)| dt \\ &= |X(1)| - |X(0)| \\ &= |q| \\ &= L(\lambda_q). \end{aligned}$$

This proves that in fact γ is the shortest path connecting p and q . \square

We now introduce the important notion of totally geodesic submanifolds of a Riemannian manifold.

Definition 7.23. Let (N, h) be a Riemannian manifold and M be a submanifold with the induced metric. Then M is said to be **totally geodesic** in N if its second fundamental form vanishes identically i.e. $B \equiv 0$.

For the totally geodesic submanifolds we have the following important characterisation.

Proposition 7.24. *Let (N, h) be a Riemannian manifold and M be a submanifold equipped with the induced metric. Then the following conditions are equivalent*

- (i) M is totally geodesic in N
- (ii) if $\gamma : I \rightarrow M$ is a geodesic in M then it is also geodesic in N .

PROOF. The result is a direct consequence of the decomposition formula

$$\nabla_{\dot{\gamma}} \dot{\gamma} = (\nabla_{\dot{\gamma}} \dot{\gamma})^{\top} + (\nabla_{\dot{\gamma}} \dot{\gamma})^{\perp} = \tilde{\nabla}_{\dot{\gamma}} \dot{\gamma} + B(\dot{\gamma}, \dot{\gamma})$$

and the polar identity for the symmetric second fundamental form

$$4 \cdot B(X, Y) = B(X + Y, X + Y) - B(X - Y, X - Y).$$

□

Proposition 7.25. *Let (N, h) be a Riemannian manifold and M be a submanifold of N . For a point (p, v) of the tangent bundle TM let $\gamma_{(p,v)} : I \rightarrow N$ be the maximal geodesic in N with $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. Then M is totally geodesic in (N, h) if $\gamma_{(p,v)}(I)$ is contained in M for all $(p, v) \in TM$. The converse is true if M is complete.*

PROOF. See Exercise 7.5.

□

Corollary 7.26. *Let (N, h) be a Riemannian manifold, $p \in N$ and V be an m -dimensional linear subspace of the tangent space $T_p N$ of N at p . Then there exists (locally) at most one totally geodesic submanifold M of (N, h) such that $T_p M = V$.*

PROOF. See Exercise 7.6.

□

Proposition 7.27. *Let (N, h) be a Riemannian manifold and M be a submanifold of N which is the fixpoint set of an isometry $\phi : N \rightarrow N$. Then M is totally geodesic in N .*

PROOF. Let $p \in M$, $v \in T_p M$ and $c : J \rightarrow M$ be a curve such that $c(0) = p$ and $\dot{c}(0) = v$. Since M is the fix point set of ϕ we have $\phi(p) = p$ and $d\phi_p(v) = v$. Further let $\gamma : I \rightarrow N$ be the maximal geodesic in N with $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. The map $\phi : N \rightarrow N$ is an isometry so the curve $\phi \circ \gamma : I \rightarrow N$ is also a geodesic. The uniqueness result of Theorem 7.9, $\phi(\gamma(0)) = \gamma(0)$ and $d\phi(\dot{\gamma}(0)) = \dot{\gamma}(0)$ then imply that $\phi(\gamma) = \gamma$. Hence the image of the geodesic $\gamma : I \rightarrow N$ is contained in M , so following Proposition 7.25 the submanifold M is totally geodesic in N .

□

Corollary 7.28. *Let $m < n$ be positive integers. Then the m -dimensional sphere*

$$S^m = \{(x, 0) \in \mathbb{R}^{m+1} \times \mathbb{R}^{n-m} \mid |x|^2 = 1\}$$

is a totally geodesic submanifold of

$$S^n = \{(x, y) \in \mathbb{R}^{m+1} \times \mathbb{R}^{n-m} \mid |x|^2 + |y|^2 = 1\}.$$

PROOF. The statement is a direct consequence of the fact that S^m is the fixpoint set of the isometry $\phi : S^n \rightarrow S^n$ of S^n with $(x, y) \mapsto (x, -y)$.

□

Corollary 7.29. *Let $m < n$ be positive integers. Let H^n be the n -dimensional hyperbolic space modelled on the upper half space $\mathbb{R}^+ \times \mathbb{R}^{n-1}$ equipped with the Riemannian metric*

$$g(X, Y) = \frac{1}{x_1^2} \langle X, Y \rangle_{\mathbb{R}^n},$$

where $x = (x_1, \dots, x_n) \in H^n$. Then the m -dimensional hyperbolic space

$$H^m = \{(x, 0) \in H^n \mid x \in \mathbb{R}^m\}$$

is totally geodesic in H^n .

PROOF. See Exercise 7.8. □

We conclude this chapter by introducing the important Riemannian symmetric spaces. These were classified by Elie Cartan in his fundamental study from 1926. For this see the standard reference, Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Academic Press (1978).

Definition 7.30. A **symmetric space** is a Riemannian manifold (M, g) such that for each point $p \in M$ there exists a global isometry $\phi_p : M \rightarrow M$ which is a geodesic symmetry fixing p . By this we mean that $\phi_p(p) = p$ and the tangent map $d\phi_p : T_p M \rightarrow T_p M$ satisfies $d\phi_p(X) = -X$ for all $X \in T_p M$.

The standard sphere is an important example of a Riemannian symmetric space.

Example 7.31. Let p be an arbitrary point on the standard sphere S^m as a subset of \mathbb{R}^{n+1} . Then the reflection $\rho_p : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ about the line generated by p is given by

$$\rho_p : q \mapsto 2\langle q, p \rangle p - q.$$

This is a linear map hence identical to its differential $\rho_p : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$. The restriction $\phi_p = \rho_p|_{S^m} : S^m \rightarrow S^m$ is an isometry that fixes p . Its tangent map $d\phi_p : T_p S^m \rightarrow T_p S^m$ satisfies $d\phi_p(X) = -X$ for all $X \in T_p S^m$. This shows that the homogeneous space S^m is symmetric.

Proposition 7.32. *Every Riemannian symmetric space is complete.*

PROOF. See Exercise 7.10. □

The following important result is a direct consequence of the famous Hopf-Rinow theorem.

Theorem 7.33. *Let (M, g) be a complete Riemannian manifold which is path-connected. If $p, q \in M$ then there exists a geodesic $\gamma : \mathbb{R} \rightarrow M$ such that $\gamma(0) = p$ and $\gamma(1) = q$.*

PROOF. See Exercise 7.11. □

The following shows that every Riemannian symmetric space is homogeneous, see Definition 5.19.

Theorem 7.34. *Every Riemannian symmetric space is homogeneous.*

PROOF. See Exercise 7.12. □

Exercises

Exercise 7.1. The result of Exercise 5.3 shows that the two dimensional hyperbolic disc H^2 introduced in Example 5.9 is isometric to the upper half plane $M = (\{(x, y) \in \mathbb{R}^2 \mid y \in \mathbb{R}^+\})$ equipped with the Riemannian metric

$$g(X, Y) = \frac{1}{y^2} \langle X, Y \rangle_{\mathbb{R}^2}.$$

Use your local library to find all geodesics in (M, g) .

Exercise 7.2. Let the orthogonal group $\mathbf{O}(m)$ be equipped with the standard left-invariant metric

$$g(A, B) = \text{trace}(A^t \cdot B).$$

Prove that a C^2 -curve $\gamma : (-\epsilon, \epsilon) \rightarrow \mathbf{O}(m)$ is a geodesic if and only if

$$\gamma^t \cdot \ddot{\gamma} = \ddot{\gamma}^t \cdot \gamma.$$

Exercise 7.3. Let the orthogonal group $\mathbf{O}(m)$ be equipped with the standard left-invariant metric

$$g(A, B) = \text{trace}(A^t \cdot B).$$

Let $p \in \mathbf{O}(m)$ and $X \in T_e \mathbf{O}(m)$. Use the result of Exercise 7.2 to show that the curve $\gamma : \mathbb{R} \rightarrow \mathbf{O}(m)$ given by $\gamma(s) = p \cdot \text{Exp}(s \cdot X)$ is a geodesic satisfying $\gamma(0) = p$ and $\dot{\gamma}(0) = p \cdot X$.

Exercise 7.4. For the real parameter $\theta \in (0, \pi/2)$ define the 2-dimensional torus T_θ^2 by

$$T_\theta^2 = \{(\cos \theta \cdot e^{i\alpha}, \sin \theta \cdot e^{i\beta}) \in S^3 \mid \alpha, \beta \in \mathbb{R}\}.$$

Determine for which $\theta \in (0, \pi/2)$ the torus T_θ^2 is a minimal submanifold of the 3-dimensional sphere

$$S^3 = \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_1|^2 + |z_2|^2 = 1\}.$$

Exercise 7.5. Find a proof of Proposition 7.25.

Exercise 7.6. Find a proof of Corollary 7.26.

Exercise 7.7. Determine the totally geodesic submanifolds of the m -dimensional real projective space $\mathbb{R}P^m$.

Exercise 7.8. Find a proof of Corollary 7.29.

Exercise 7.9. Let the orthogonal group $\mathbf{O}(m)$ be equipped with the left-invariant metric

$$g(A, B) = \text{trace}(A^t \cdot B)$$

and let K be a Lie subgroup of $\mathbf{O}(m)$. Prove that K is totally geodesic in $\mathbf{O}(m)$.

Exercise 7.10. Find a proof of Proposition 7.32.

Exercise 7.11. Use your local library to find a proof of Theorem 7.33.

Exercise 7.12. Find a proof of Theorem 7.34.

The Riemann Curvature Tensor

In this chapter we introduce the Riemann curvature tensor and the sectional curvatures of a Riemannian manifold. These notions generalise the Gaussian curvature playing a central role in the classical differential geometry of surfaces. We derive the important Gauss equation comparing the sectional curvatures of a submanifold and that of its ambient space. We prove that the Euclidean spaces, the standard spheres and the hyperbolic spaces all have constant sectional curvature. We then determine the Riemannian curvature tensor for manifolds of constant sectional curvature and also for an important class of Lie groups.

Let (M, g) be a Riemannian manifold and ∇ be its Levi-Civita connection. Then for each vector field $X \in C^\infty(TM)$ we have the **first order covariant derivative**

$$\nabla_X : C^\infty(TM) \rightarrow C^\infty(TM)$$

of a vector field in the direction of X satisfying

$$\nabla_X : Z \mapsto \nabla_X Z.$$

We will now generalise this and introduce the covariant derivative of tensor fields of type $(r, 0)$ and $(r, 1)$.

As a motivation, let us assume that A is a tensor field of type $(2, 1)$. If we differentiate $A(Y, Z)$ in the direction of X applying the naive "product rule"

$$\nabla_X(A(Y, Z)) = (\nabla_X A)(Y, Z) + A(\nabla_X Y, Z) + A(Y, \nabla_X Z)$$

we yield

$$(\nabla_X A)(Y, Z) = \nabla_X(A(Y, Z)) - A(\nabla_X Y, Z) - A(Y, \nabla_X Z),$$

where $\nabla_X A$ is the "covariant derivative" of the tensor field A in the direction of X . This idea turns out to be very useful and leads to the following formal definition.

Definition 8.1. Let (M, g) be a Riemannian manifold with Levi-Civita connection ∇ . For a tensor field $A : C_r^\infty(TM) \rightarrow C_0^\infty(TM)$ of type $(r, 0)$ we define its **covariant derivative**

$$\nabla A : C_{r+1}^\infty(TM) \rightarrow C_0^\infty(TM)$$

by

$$\nabla A : (X, X_1, \dots, X_r) \mapsto (\nabla_X A)(X_1, \dots, X_r) =$$

$$X(A(X_1, \dots, X_r)) - \sum_{k=1}^r A(X_1, \dots, X_{k-1}, \nabla_X X_k, X_{k+1}, \dots, X_r).$$

A tensor field A of type $(r, 0)$ is said to be **parallel** if $\nabla A \equiv 0$.

The following result can be seen as, yet another, compatibility of the Levi-Civita connection ∇ of (M, g) with the Riemannian metric g .

Proposition 8.2. *Let (M, g) be a Riemannian manifold. Then the metric g is a parallel tensor field of type $(2, 0)$.*

PROOF. See Exercise 8.1. □

Example 8.3. Let (M, g) be a Riemannian manifold. Then we already know that its Levi-Civita connection ∇ is tensorial in its first argument i.e. if $X, Y \in C^\infty(TM)$ and $f, g \in C^\infty(M)$ then

$$\nabla(fX + gY)Z = f\nabla_X Z + g\nabla_Y Z.$$

This means that a vector field $Z \in C^\infty(TM)$ on M induces a **natural tensor field** $\mathcal{Z} : C_1^\infty(TM) \rightarrow C_1^\infty(TM)$ of type $(1, 1)$ given by

$$\mathcal{Z} : X \mapsto \nabla_X Z.$$

Definition 8.4. Let (M, g) be a Riemannian manifold with Levi-Civita connection ∇ . For a tensor field $B : C_r^\infty(TM) \rightarrow C_1^\infty(TM)$ of type $(r, 1)$ we define its **covariant derivative**

$$\nabla B : C_{r+1}^\infty(TM) \rightarrow C_1^\infty(TM)$$

by

$$\nabla B : (X, X_1, \dots, X_r) \mapsto (\nabla_X B)(X_1, \dots, X_r) =$$

$$\nabla_X (B(X_1, \dots, X_r)) - \sum_{k=1}^r B(X_1, \dots, X_{k-1}, \nabla_X X_k, X_{k+1}, \dots, X_r).$$

A tensor field B of type $(r, 1)$ is said to be **parallel** if $\nabla B \equiv 0$.

Definition 8.5. Let $X, Y \in C^\infty(TM)$ be two vector fields on the Riemannian manifold (M, g) with Levi-Civita connection ∇ . Then the **second order covariant derivative**

$$\nabla^2_{X,Y} : C^\infty(TM) \rightarrow C^\infty(TM)$$

is defined by

$$\nabla^2_{X,Y} : Z \mapsto (\nabla_X \mathcal{Z})(Y),$$

where \mathcal{Z} is the natural tensor field of type $(1,1)$ induced by $Z \in C^\infty(TM)$, see Example 8.3.

As a direct consequence of Definitions 8.4 and 8.5 we see that if $X, Y, Z \in C^\infty(TM)$ are vector fields on M , then the second order covariant derivative $\nabla^2_{X,Y}$ satisfies

$$\nabla^2_{X,Y} Z = \nabla_X(\mathcal{Z}(Y)) - \mathcal{Z}(\nabla_X Y) = \nabla_X \nabla_Y Z - \nabla_{\nabla_X Y} Z.$$

This leads us to the following important definition.

Definition 8.6. Let (M, g) be a Riemannian manifold with Levi-Civita connection ∇ . Then its **Riemann curvature operator**

$$R : C^\infty(TM) \times C^\infty(TM) \times C^\infty(TM) \rightarrow C^\infty(TM)$$

is defined as twice the skew-symmetric part of the second covariant derivative ∇^2 i.e.

$$R(X, Y)Z = \nabla^2_{X,Y} Z - \nabla^2_{Y,X} Z.$$

The next remarkable result shows that the curvature operator is actually a tensor field.

Theorem 8.7. *Let (M, g) be a Riemannian manifold with Levi-Civita connection ∇ . Then the Riemann curvature operator*

$$R : C_3^\infty(TM) \rightarrow C_1^\infty(TM)$$

satisfying

$$R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z$$

is a tensor field on M of type $(3,1)$.

PROOF. See Exercise 8.2. □

The reader should note that the Riemann curvature tensor is an intrinsic object since it only depends on the intrinsic Levi-Civita connection. The following result shows that the curvature tensor has many beautiful symmetries.

Proposition 8.8. *Let (M, g) be a Riemannian manifold. For vector fields $X, Y, Z, W \in C^\infty(TM)$ on M we then have*

- (i) $R(X, Y)Z = -R(Y, X)Z$,
- (ii) $g(R(X, Y)Z, W) = -g(R(X, Y)W, Z)$,
- (iii) $R(X, Y)Z + R(Z, X)Y + R(Y, Z)X = 0$,
- (iv) $g(R(X, Y)Z, W) = g(R(Z, W)X, Y)$,
- (v) $6 \cdot R(X, Y)Z = R(X, Y + Z)(Y + Z) - R(X, Y - Z)(Y - Z) + R(X + Z, Y)(X + Z) - R(X - Z, Y)(X - Z)$.

PROOF. See Exercise 8.3. \square

Part (iii) of Proposition 8.8 is the so called first **Bianchi identity**. The second Bianchi identity is a similar result concerning the covariant derivative ∇R of the curvature tensor. This will not be treated here.

Definition 8.9. Let (M, g) be a Riemannian manifold and $p \in M$. Then a **section** V at p is a 2-dimensional subspace of the tangent space $T_p M$. The set

$$G_2(T_p M) = \{V \mid V \text{ is a section of } T_p M\}$$

of sections is called the **Grassmannian** of 2-planes at p .

Remark 8.10. In Gaussian geometry the tangent space $T_p \Sigma$ of a surface Σ in the Euclidean \mathbb{R}^3 is two dimensional. This means that in this case there is only one section at $p \in \Sigma$ namely the full two dimensional tangent plane $T_p \Sigma$.

Before introducing the notion of the sectional curvature we need the following technical lemma.

Lemma 8.11. *Let (M, g) be a Riemannian manifold, $p \in M$ and $X, Y, Z, W \in T_p M$ be tangent vectors at p such that the two sections $\text{span}_{\mathbb{R}}\{X, Y\}$ and $\text{span}_{\mathbb{R}}\{Z, W\}$ are identical. Then*

$$\frac{g(R(X, Y)Y, X)}{|X|^2|Y|^2 - g(X, Y)^2} = \frac{g(R(Z, W)W, Z)}{|Z|^2|W|^2 - g(Z, W)^2}.$$

PROOF. See Exercise 8.4. \square

Definition 8.12. Let (M, g) be a Riemannian manifold and $p \in M$. Then the function $K_p : G_2(T_p M) \rightarrow \mathbb{R}$ given by

$$K_p : \text{span}_{\mathbb{R}}\{X, Y\} \mapsto \frac{g(R(X, Y)Y, X)}{|X|^2|Y|^2 - g(X, Y)^2}$$

is called the **sectional curvature** at p . We usually write $K(X, Y)$ for $K(\text{span}_{\mathbb{R}}\{X, Y\})$.

Definition 8.13. Let (M, g) be a Riemannian manifold, $p \in M$ and $K_p : G_2(T_p M) \rightarrow \mathbb{R}$ be the sectional curvature at p . Then we define the functions $\delta, \Delta : M \rightarrow \mathbb{R}$ by

$$\delta : p \mapsto \min_{V \in G_2(T_p M)} K_p(V) \quad \text{and} \quad \Delta : p \mapsto \max_{V \in G_2(T_p M)} K_p(V).$$

The Riemannian manifold (M, g) is said to be

- (i) of **positive curvature** if $\delta(p) \geq 0$ for all p ,
- (ii) of **strictly positive curvature** if $\delta(p) > 0$ for all p ,
- (iii) of **negative curvature** if $\Delta(p) \leq 0$ for all p ,
- (iv) of **strictly negative curvature** if $\Delta(p) < 0$ for all p ,
- (v) of **constant curvature** if $\delta = \Delta$ is constant,
- (vi) **flat** if $\delta \equiv \Delta \equiv 0$.

The next example shows how the Riemann curvature tensor can be presented by means local coordinates. Hopefully this will convince the reader that those should be avoided whenever possible.

Example 8.14. Let (M, g) be a Riemannian manifold and let (U, x) be a chart on M . For $i, j, k, l = 1, \dots, m$ put

$$X_i = \frac{\partial}{\partial x_i}, \quad g_{ij} = g(X_i, X_j) \quad \text{and} \quad R_{ijkl} = g(R(X_i, X_j)X_k, X_l).$$

Then

$$R_{ijkl} = \sum_{s=1}^m g_{sl} \left(\frac{\partial \Gamma_{jk}^s}{\partial x_i} - \frac{\partial \Gamma_{ik}^s}{\partial x_j} + \sum_{r=1}^m \{ \Gamma_{jk}^r \cdot \Gamma_{ir}^s - \Gamma_{ik}^r \cdot \Gamma_{jr}^s \} \right),$$

where the functions Γ_{ij}^k are the Christoffel symbols of the Levi-Civita connection ∇ of (M, g) with respect to (U, x) .

PROOF. Using the fact that $[X_i, X_j] = 0$, see Proposition 4.29, we obtain

$$\begin{aligned} R(X_i, X_j)X_k &= \nabla_{X_i} \nabla_{X_j} X_k - \nabla_{X_j} \nabla_{X_i} X_k \\ &= \sum_{s=1}^m \{ \nabla_{X_i} (\Gamma_{jk}^s \cdot X_s) - \nabla_{X_j} (\Gamma_{ik}^s \cdot X_s) \} \\ &= \sum_{s=1}^m \left(\frac{\partial \Gamma_{jk}^s}{\partial x_i} \cdot X_s + \sum_{r=1}^m \Gamma_{jk}^s \Gamma_{ir}^r X_r - \frac{\partial \Gamma_{ik}^s}{\partial x_j} \cdot X_s - \sum_{r=1}^m \Gamma_{ik}^s \Gamma_{jr}^r X_r \right) \\ &= \sum_{s=1}^m \left(\frac{\partial \Gamma_{jk}^s}{\partial x_i} - \frac{\partial \Gamma_{ik}^s}{\partial x_j} + \sum_{r=1}^m \{ \Gamma_{jk}^r \Gamma_{ir}^s - \Gamma_{ik}^r \Gamma_{jr}^s \} \right) X_s. \end{aligned}$$

□

Example 8.15. Let E^m be the m -dimensional vector space \mathbb{R}^m equipped with the Euclidean metric $\langle, \rangle_{\mathbb{R}^m}$. The set $\{\partial/\partial x_1, \dots, \partial/\partial x_m\}$ is a global frame for the tangent bundle $T\mathbb{R}^m$. In this situation we have $g_{ij} = \delta_{ij}$, so $\Gamma_{ij}^k \equiv 0$ by Example 6.13. This implies that $R \equiv 0$ so E^m is flat.

We will now prove the important **Gauss equation** comparing the curvature tensor of a submanifold and that of its ambient space in terms of the second fundamental form.

Theorem 8.16. *Let (N, h) be a Riemannian manifold and M be a submanifold of N equipped with the induced metric g . Let $X, Y, Z, W \in C^\infty(TN)$ be vector fields extending $\tilde{X}, \tilde{Y}, \tilde{Z}, \tilde{W} \in C^\infty(TM)$. Then at a point $p \in M$ we have*

$$\begin{aligned} & g(\tilde{R}(\tilde{X}, \tilde{Y})\tilde{Z}, \tilde{W}) - h(R(X, Y)Z, W) \\ &= h(B(\tilde{Y}, \tilde{Z}), B(\tilde{X}, \tilde{W})) - h(B(\tilde{X}, \tilde{Z}), B(\tilde{Y}, \tilde{W})). \end{aligned}$$

Here \tilde{R} and R are the Riemann curvature tensors of (M, g) and (N, h) , respectively, and B the second fundamental of M as a submanifold of N .

PROOF. Employing the definitions of the curvature tensors \tilde{R} , R , the Levi-Civita connection $\tilde{\nabla}$ and the second fundamental form B of M as a submanifold of N we obtain the following:

$$\begin{aligned} & g(\tilde{R}(\tilde{X}, \tilde{Y})\tilde{Z}, \tilde{W}) \\ &= g(\tilde{\nabla}_{\tilde{X}}\tilde{\nabla}_{\tilde{Y}}\tilde{Z} - \tilde{\nabla}_{\tilde{Y}}\tilde{\nabla}_{\tilde{X}}\tilde{Z} - \tilde{\nabla}_{[\tilde{X}, \tilde{Y}]\tilde{Z}}, \tilde{W}) \\ &= g((\nabla_X(\nabla_Y Z)^\top)^\top - (\nabla_Y(\nabla_X Z)^\top)^\top - (\nabla_{[X, Y]}Z)^\top, W) \\ &= h((\nabla_X(\nabla_Y Z - (\nabla_Y Z)^\perp))^\top - (\nabla_Y(\nabla_X Z - (\nabla_X Z)^\perp))^\top, W) \\ &\quad - h((\nabla_{[X, Y]}Z - (\nabla_{[X, Y]}Z)^\perp)^\top, W) \\ &= h(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]}Z, W) \\ &\quad - h((\nabla_X(\nabla_Y Z)^\perp, W) + h(\nabla_Y(\nabla_X Z)^\perp, W) \\ &= h(R(X, Y)Z, W) \\ &\quad + h((\nabla_Y Z)^\perp, (\nabla_X W)^\perp) - h((\nabla_X Z)^\perp, (\nabla_Y W)^\perp) \\ &= h(R(X, Y)Z, W) \\ &\quad + h(B(\tilde{Y}, \tilde{Z}), B(\tilde{X}, \tilde{W})) - h(B(\tilde{X}, \tilde{Z}), B(\tilde{Y}, \tilde{W})). \end{aligned}$$

□

We will now employ the Gauss equation to the classical situation of a surface in the three dimensional Euclidean space.

Example 8.17. Let Σ be a regular surface in the Euclidean \mathbb{R}^3 . Let $\{\tilde{X}, \tilde{Y}\}$ be a local orthonormal frame for the tangent bundle $T\Sigma$ of Σ around a point $p \in \Sigma$ and \tilde{N} be the local Gauss map with $\tilde{N} = \tilde{X} \times \tilde{Y}$. If X, Y, N are local extensions of $\tilde{X}, \tilde{Y}, \tilde{N}$, such that $\{X, Y, N\}$ is a local orthonormal frame for $T\mathbb{R}^3$, then at $p \in M$ the second fundamental form B of Σ in \mathbb{R}^3 satisfies

$$\begin{aligned} B(\tilde{X}, \tilde{Y}) &= (\partial_X Y)^\perp \\ &= \langle \partial_X Y, N \rangle N \\ &= -\langle Y, \partial_X N \rangle N \\ &= -\langle Y, dN(X) \rangle N \\ &= \langle \tilde{Y}, S_p(\tilde{X}) \rangle \tilde{N}, \end{aligned}$$

where $S_p : T_p\Sigma \rightarrow T_p\Sigma$ is the shape operator at p . If we now apply the fact that \mathbb{R}^3 is flat, then the Gauss equation tells us that the sectional curvature $K(\tilde{X}, \tilde{Y})$ of Σ at p satisfies

$$\begin{aligned} K(\tilde{X}, \tilde{Y}) &= \langle \tilde{R}(\tilde{X}, \tilde{Y})\tilde{Y}, \tilde{X} \rangle \\ &= \langle B(\tilde{Y}, \tilde{Y}), B(\tilde{X}, \tilde{X}) \rangle - \langle B(\tilde{X}, \tilde{Y}), B(\tilde{Y}, \tilde{X}) \rangle \\ &= \det S_p. \end{aligned}$$

In other words, the sectional curvature $K(\tilde{X}, \tilde{Y})$ is the determinant of the shape operator S_p i.e. the classical **Gaussian curvature**.

An interesting consequence of the Gauss equation is the following useful result. For important applications see Exercises 8.7 and 8.8.

Corollary 8.18. *Let (N, h) be a Riemannian manifold and M be a totally geodesic submanifold of N equipped with the induced metric g . Let $X, Y, Z, W \in C^\infty(TN)$ be vector fields extending $\tilde{X}, \tilde{Y}, \tilde{Z}, \tilde{W} \in C^\infty(TM)$. Then at a point $p \in M$ we have*

$$g(\tilde{R}(\tilde{X}, \tilde{Y})\tilde{Z}, \tilde{W}) = h(R(X, Y)Z, W).$$

PROOF. This follows directly from the fact that the second fundamental for B of M in N vanishes identically. \square

Example 8.19. The standard sphere S^m has constant sectional curvature $+1$ (see Exercises 8.6 and 8.7) and the hyperbolic space H^m has constant sectional curvature -1 (see Exercise 8.8).

Our next aim is a formula for the curvature tensor for manifolds of constant sectional curvature. This we present in Corollary 8.23. First we need some preparations.

Lemma 8.20. *Let (M, g) be a Riemannian manifold, $p \in M$ and $Y \in T_p M$. Then the map $\bar{Y} : T_p M \rightarrow T_p M$ given by*

$$\bar{Y} : X \mapsto R(X, Y)Y$$

is a symmetric endomorphism of the tangent space $T_p M$.

PROOF. If $X, Y, Z \in T_p M$ then it follows from Proposition 8.8 that

$$\begin{aligned} g(\bar{Y}(X), Z) &= g(R(X, Y)Y, Z) \\ &= g(R(Y, Z)X, Y) \\ &= g(R(Z, Y)Y, X) \\ &= g(X, \bar{Y}(Z)). \end{aligned}$$

□

For a tangent vector $Y \in T_p M$ with $|Y| = 1$ let $\mathcal{N}(Y)$ be the normal space to Y

$$\mathcal{N}(Y) = \{X \in T_p M \mid g(X, Y) = 0\}.$$

The fact that $\bar{Y}(Y) = 0$ and Lemma 8.20 ensure the existence of an orthonormal basis of eigenvectors X_1, \dots, X_{m-1} for the restriction of the symmetric endomorphism \bar{Y} to $\mathcal{N}(Y)$. Without loss of generality, we can assume that the corresponding eigenvalues satisfy

$$\lambda_1(p) \leq \dots \leq \lambda_{m-1}(p).$$

If $X \in \mathcal{N}(Y)$, $|X| = 1$ and $\bar{Y}(X) = \lambda X$ then

$$K_p(X, Y) = g(R(X, Y)Y, X) = g(\bar{Y}(X), X) = \lambda.$$

This means that the eigenvalues satisfy

$$\delta(p) \leq \lambda_1(p) \leq \dots \leq \lambda_{m-1}(p) \leq \Delta(p).$$

Definition 8.21. For a Riemannian manifold (M, g) define the tensor field $R_1 : C_3^\infty(TM) \rightarrow C_1^\infty(TM)$ of type $(3, 1)$ by

$$R_1(X, Y)Z = g(Y, Z)X - g(X, Z)Y.$$

Proposition 8.22. *Let (M, g) be a Riemannian manifold and $X, Y, Z \in C^\infty(TM)$ be vector fields on M . Then*

- (i) $|R(X, Y)Y - \frac{\delta+\Delta}{2}R_1(X, Y)Y| \leq \frac{1}{2}(\Delta - \delta)|X||Y|^2$
- (ii) $|R(X, Y)Z - \frac{\delta+\Delta}{2}R_1(X, Y)Z| \leq \frac{2}{3}(\Delta - \delta)|X||Y||Z|$

PROOF. Without loss of generality we can assume that $|X| = |Y| = |Z| = 1$. If $X = X^\perp + X^\top$ with $X^\perp \perp Y$ and X^\top is a multiple of Y then $R(X, Y)Z = R(X^\perp, Y)Z$ and $|X^\perp| \leq |X|$ so we can also assume that $X \perp Y$. Then $R_1(X, Y)Y = \langle Y, Y \rangle X - \langle X, Y \rangle Y = X$.

The first statement follows from the fact that the symmetric endomorphism of $T_p M$ with

$$X \mapsto \{R(X, Y)Y - \frac{\Delta + \delta}{2} \cdot X\}$$

restricted to $\mathcal{N}(Y)$ has eigenvalues in the interval $[\frac{\delta - \Delta}{2}, \frac{\Delta - \delta}{2}]$.

It is easily checked that the operator R_1 satisfies the conditions of Proposition 8.8 and hence $D = R - \frac{\Delta + \delta}{2} \cdot R_1$ as well. This implies that

$$\begin{aligned} 6 \cdot D(X, Y)Z &= D(X, Y + Z)(Y + Z) - D(X, Y - Z)(Y - Z) \\ &\quad + D(X + Z, Y)(X + Z) - D(X - Z, Y)(X - Z). \end{aligned}$$

The second statement then follows from (i) and

$$\begin{aligned} 6|D(X, Y)Z| &\leq \frac{1}{2}(\Delta - \delta)\{|X|(|Y + Z|^2 + |Y - Z|^2) \\ &\quad + |Y|(|X + Z|^2 + |X - Z|^2)\} \\ &= \frac{1}{2}(\Delta - \delta)\{2|X|(|Y|^2 + |Z|^2) + 2|Y|(|X|^2 + |Z|^2)\} \\ &= 4(\Delta - \delta). \end{aligned}$$

□

As a direct consequence of Proposition 8.22 we have the following useful result.

Corollary 8.23. *Let (M, g) be a Riemannian manifold of constant curvature κ . Then the curvature tensor R is given by*

$$R(X, Y)Z = \kappa \cdot (g(Y, Z)X - g(X, Z)Y).$$

PROOF. The result follows immediately from $\kappa = \delta = \Delta$. □

The following result shows that the curvature tensor takes a rather simple form in the important class of Lie groups treated in Proposition 6.12.

Proposition 8.24. *Let (G, g) be a Lie group equipped with a left-invariant metric such that for all $X \in \mathfrak{g}$ the endomorphism*

$$\text{ad}_X : \mathfrak{g} \rightarrow \mathfrak{g}$$

is skew-symmetric with respect to g . Then for left-invariant vector fields $X, Y, Z \in \mathfrak{g}$ the curvature tensor R is given by

$$R(X, Y)Z = -\frac{1}{4} \cdot [[X, Y], Z].$$

PROOF. See Exercise 8.9. \square

Corollary 8.25. Let (G, g) be a Lie group equipped with a left-invariant metric such that for all $Z \in \mathfrak{g}$ the endomorphism

$$\text{ad}_Z : \mathfrak{g} \rightarrow \mathfrak{g}$$

is skew-symmetric with respect to g . Let $X, Y \in \mathfrak{g}$ be left-invariant vector fields such that $|X| = |Y| = 1$ and $g(X, Y) = 0$. Then the sectional curvature $K(X, Y)$ satisfies

$$K(X, Y) = \frac{1}{4} \cdot |[X, Y]|^2.$$

PROOF. See Exercise 8.10. \square

We conclude this chapter by defining the Ricci and scalar curvatures of a Riemannian manifold. These are obtained by taking traces over the curvature tensor and play an important role in Riemannian geometry.

Definition 8.26. Let (M, g) be a Riemannian manifold, then we define

(i) the **Ricci operator** $r : C_1^\infty(TM) \rightarrow C_1^\infty(M)$ by

$$r(X) = \sum_{i=1}^m R(X, e_i)e_i,$$

(ii) the **Ricci curvature** $Ric : C_2^\infty(TM) \rightarrow C_0^\infty(TM)$ by

$$Ric(X, Y) = \sum_{i=1}^m g(R(X, e_i)e_i, Y),$$

(iii) the **scalar curvature** $s \in C^\infty(M)$ by

$$s = \sum_{j=1}^m Ric(e_j, e_j) = \sum_{j=1}^m \sum_{i=1}^m g(R(e_i, e_j)e_j, e_i).$$

Here $\{e_1, \dots, e_m\}$ is any local orthonormal frame for the tangent bundle.

In the case of constant sectional curvature we have the following result.

Corollary 8.27. *Let (M^m, g) be a Riemannian manifold of constant sectional curvature κ . Then its scalar curvature satisfies the following*

$$s = m \cdot (m - 1) \cdot \kappa.$$

PROOF. Let $\{e_1, \dots, e_m\}$ be any local orthonormal frame. Then Corollary 8.23 implies that

$$\begin{aligned} Ric(e_j, e_j) &= \sum_{i=1}^m g(R(e_j, e_i)e_i, e_j) \\ &= \sum_{i=1}^m g(\kappa(g(e_i, e_i)e_j - g(e_j, e_i)e_i), e_j) \\ &= \kappa\left(\sum_{i=1}^m g(e_i, e_i)g(e_j, e_j) - \sum_{i=1}^m g(e_i, e_j)g(e_i, e_j)\right) \\ &= \kappa\left(\sum_{i=1}^m 1 - \sum_{i=1}^m \delta_{ij}\right) = (m - 1) \cdot \kappa. \end{aligned}$$

To obtain the formula for the scalar curvature s we only need to multiply the constant Ricci curvature $Ric(e_j, e_j)$ by m . \square

As a reference on further notions of curvature we recommend the interesting book, Wolfgang Kühnel, Differential Geometry: Curves - Surfaces - Manifolds, American Mathematical Society (2002).

Exercises

Exercise 8.1. Let (M, g) be a Riemannian manifold. Prove that the tensor field g of type $(2, 0)$ is parallel with respect to the Levi-Civita connection.

Exercise 8.2. Let (M, g) be a Riemannian manifold. Prove that the Riemann curvature operator R is a tensor field of type $(3, 1)$.

Exercise 8.3. Find a proof for Proposition 8.8.

Exercise 8.4. Find a proof for Lemma 8.11.

Exercise 8.5. Let \mathbb{R}^m and \mathbb{C}^m be equipped with their standard Euclidean metric g given by

$$g(z, w) = \operatorname{Re} \sum_{k=1}^m z_k \bar{w}_k$$

and let $T^m = \{z \in \mathbb{C}^m \mid |z_1| = \dots = |z_m| = 1\}$ be the m -dimensional torus in \mathbb{C}^m with the induced metric. Find an isometric immersion $\phi : \mathbb{R}^m \rightarrow T^m$, determine all geodesics on T^m and prove that the torus is flat.

Exercise 8.6. Let the Lie group $S^3 \cong \mathbf{SU}(2)$ be equipped with the metric

$$g(Z, W) = \frac{1}{2} \operatorname{Re}(\operatorname{trace}(\bar{Z}^t W)).$$

- (i) Find an orthonormal basis for $T_e \mathbf{SU}(2)$.
- (ii) Prove that $(\mathbf{SU}(2), g)$ has constant sectional curvature $+1$.

Exercise 8.7. Let S^m be the unit sphere in \mathbb{R}^{m+1} equipped with the standard Euclidean metric $\langle \cdot, \cdot \rangle_{\mathbb{R}^{m+1}}$. Use the results of Corollaries 7.28, 8.18 and Exercise 8.6 to prove that $(S^m, \langle \cdot, \cdot \rangle_{\mathbb{R}^{m+1}})$ has constant sectional curvature $+1$.

Exercise 8.8. Let H^m be the m -dimensional hyperbolic space modelled on the upper half space $\mathbb{R}^+ \times \mathbb{R}^{m-1}$ equipped with the Riemannian metric

$$g(X, Y) = \frac{1}{x_1^2} \langle X, Y \rangle_{\mathbb{R}^m},$$

where $x = (x_1, \dots, x_m) \in H^m$. For $k = 1, \dots, m$ let the vector fields $X_k \in C^\infty(TH^m)$ be given by

$$(X_k)_x = x_1 \cdot \frac{\partial}{\partial x_k}$$

and define the operation $*$ on H^m by

$$(\alpha, x) * (\beta, y) = (\alpha \cdot \beta, \alpha \cdot y + x).$$

Prove that

- (i) $(H^m, *)$ is a Lie group,
- (ii) the vector fields X_1, \dots, X_m are left-invariant,
- (iii) $[X_k, X_l] = 0$ and $[X_1, X_k] = X_k$ for $k, l = 2, \dots, m$,
- (iv) the metric g is left-invariant,
- (v) (H^m, g) has constant curvature -1 .

Compare with Exercises 6.4 and 7.1.

Exercise 8.9. Find a proof for Proposition 8.24.

Exercise 8.10. Find a proof for Corollary 8.25.

Curvature and Local Geometry

This chapter is devoted to the study of the local geometry of a Riemannian manifold and how this is controlled by its curvature tensor. For this we introduce the notion of a Jacobi field which is a standard tool in differential geometry. With this at hand we obtain a fundamental comparison result describing the curvature dependence of local distances.

Definition 9.1. Let (M, g) be a Riemannian manifold. By a 1-parameter **family of geodesics** we mean a C^3 -map

$$\Phi : (-\epsilon, \epsilon) \times I \rightarrow M$$

such that the curve $\gamma_t : I \rightarrow M$ given by $\gamma_t : s \mapsto \Phi(t, s)$ is a geodesic for all $t \in (-\epsilon, \epsilon)$. The variable $t \in (-\epsilon, \epsilon)$ is called the **family parameter** of Φ .

The following result suggests that the Riemann curvature tensor is closely related to the local behaviour of geodesics.

Proposition 9.2. *Let (M, g) be a Riemannian manifold and $\Phi : (-\epsilon, \epsilon) \times I \rightarrow M$ be a 1-parameter family of geodesics. Then for each $t \in (-\epsilon, \epsilon)$ the vector field $J_t : I \rightarrow TM$ along γ_t , given by*

$$J_t(s) = \frac{\partial \Phi}{\partial t}(t, s),$$

satisfies the second order linear ordinary differential equation

$$\nabla_{\dot{\gamma}_t} \nabla_{\dot{\gamma}_t} J_t + R(J_t, \dot{\gamma}_t) \dot{\gamma}_t = 0.$$

PROOF. Along Φ we define the vector fields $X(t, s) = \partial \Phi / \partial s$ and $J(t, s) = \partial \Phi / \partial t$. The fact that $[\partial / \partial t, \partial / \partial s] = 0$ implies that

$$[J, X] = [d\Phi(\partial / \partial t), d\Phi(\partial / \partial s)] = d\Phi([\partial / \partial t, \partial / \partial s]) = 0.$$

Since Φ is a family of geodesics we have $\nabla_X X = 0$ and the definition of the curvature tensor then implies that

$$\begin{aligned} R(J, X)X &= \nabla_J \nabla_X X - \nabla_X \nabla_J X - \nabla_{[J, X]} X \\ &= -\nabla_X \nabla_J X \end{aligned}$$

$$= -\nabla_X \nabla_X J.$$

Hence for each $t \in (-\epsilon, \epsilon)$ we have

$$\nabla_{\dot{\gamma}_t} \nabla_{\dot{\gamma}_t} J_t + R(J_t, \dot{\gamma}_t) \dot{\gamma}_t = 0.$$

□

The result of Proposition 9.2 leads to the following natural notion.

Definition 9.3. Let (M, g) be a Riemannian manifold, $\gamma : I \rightarrow M$ be a geodesic and $X = \dot{\gamma}$ be the tangent vector field along γ . A C^2 vector field J along γ is called a **Jacobi field** if and only if

$$(5) \quad \nabla_X \nabla_X J + R(J, X)X = 0$$

along γ . We denote the space of all Jacobi fields along γ by $\mathcal{J}_\gamma(TM)$.

We now give an example of a 1-parameter family of geodesics in the Euclidean space E^{m+1} .

Example 9.4. Let $c, n : \mathbb{R} \rightarrow E^{m+1}$ be smooth curves such that the image $n(\mathbb{R})$ of n is contained in the unit sphere S^m . If we define a map $\Phi : \mathbb{R} \times \mathbb{R} \rightarrow E^{m+1}$ by

$$\Phi : (t, s) \mapsto c(t) + s \cdot n(t)$$

then for each $t \in \mathbb{R}$ the curve $\gamma_t : s \mapsto \Phi(t, s)$ is a straight line and hence a geodesic in E^{m+1} . By differentiating this with respect to the family parameter t we yield the Jacobi field $J \in \mathcal{J}_{\gamma_0}(TE^{m+1})$ along γ_0 satisfying

$$J(s) = \frac{d}{dt} \Phi(t, s)|_{t=0} = \dot{c}(0) + s \cdot \dot{n}(0).$$

The Jacobi equation (5) is linear in J . This means that the space of Jacobi fields $\mathcal{J}_\gamma(TM)$ along γ is a vector space. We are now interested in determining its dimension.

Proposition 9.5. Let (M^m, g) be a Riemannian manifold, $p \in M$, $\gamma : I \rightarrow M$ be a geodesic with $\gamma(0) = p$ and $X = \dot{\gamma}$ be the tangent vector field along γ . If $v, w \in T_p M$ are two tangent vectors at p then there exists a unique Jacobi field J along γ such that

$$J_p = v \quad \text{and} \quad (\nabla_X J)_p = w.$$

PROOF. In the spirit of Proposition 7.8 let $\{X_1, \dots, X_m\}$ be an orthonormal frame of parallel vector fields along γ . If J is a vector field along γ then

$$J = \sum_{i=1}^m a_i X_i,$$

where $a_i = g(J, X_i)$ are C^2 -functions on the real interval I . The vector fields X_1, \dots, X_m are parallel so

$$\nabla_X J = \sum_{i=1}^m \dot{a}_i X_i \quad \text{and} \quad \nabla_X \nabla_X J = \sum_{i=1}^m \ddot{a}_i X_i.$$

For the curvature tensor we have

$$R(X_i, X)X = \sum_{k=1}^m b_i^k X_k,$$

where $b_i^k = g(R(X_i, X)X, X_k)$ are smooth functions on the real interval I , heavily depending on the geometry of (M, g) . This means that $R(J, X)X$ is given by

$$R(J, X)X = \sum_{i,k=1}^m a_i b_i^k X_k$$

and that J is a Jacobi field if and only if

$$\sum_{i=1}^m (\ddot{a}_i + \sum_{k=1}^m a_k b_k^i) X_i = 0.$$

This is clearly equivalent to the following second order system of linear ordinary differential equations in $a = (a_1, \dots, a_m)$:

$$\ddot{a}_i + \sum_{k=1}^m a_k b_k^i = 0 \quad \text{for all } i = 1, 2, \dots, m.$$

A global solution will always exist and is uniquely determined by the initial values $a(0)$ and $\dot{a}(0)$. This implies that J exists globally and is uniquely determined by the initial conditions

$$J(0) = v \quad \text{and} \quad (\nabla_X J)(0) = w.$$

□

As an immediate consequence of Proposition 9.5 we have the following interesting result.

Corollary 9.6. *Let (M^m, g) be a Riemannian manifold and $\gamma : I \rightarrow M$ be a geodesic in M . Then the vector space $\mathcal{J}_\gamma(TM)$, of Jacobi fields along γ , has the dimension $2m$.*

The following Lemma 9.7 shows that when proving results about Jacobi fields along a geodesic γ we can always assume, without loss of generality, that they are parametrised by arclength i.e. $|\dot{\gamma}| = 1$.

Lemma 9.7. *Let (M, g) be a Riemannian manifold, $\gamma : I \rightarrow M$ be a geodesic and J be a Jacobi field along γ . If λ is a non-zero real number and $\sigma : \lambda I \rightarrow I$ is given by $\sigma : t \mapsto t/\lambda$, then $\gamma \circ \sigma : \lambda I \rightarrow M$ is a geodesic and $J \circ \sigma$ is a Jacobi field along $\gamma \circ \sigma$.*

PROOF. See Exercise 9.1. \square

The next result shows that both the tangential and the normal parts of a Jacobi field are again Jacobi fields. Furthermore we completely determine the tangential Jacobi fields.

Proposition 9.8. *Let (M, g) be a Riemannian manifold, $\gamma : I \rightarrow M$ be a geodesic with $|\dot{\gamma}| = 1$ and J be a Jacobi field along γ . Let J^\top be the tangential part of J given by*

$$J^\top = g(J, \dot{\gamma})\dot{\gamma} \quad \text{and} \quad J^\perp = J - J^\top$$

be its normal part. Then J^\top and J^\perp are Jacobi fields along γ and there exist $a, b \in \mathbb{R}$ such that $J^\top(s) = (as + b)\dot{\gamma}(s)$ for all $s \in I$.

PROOF. In this situation we have

$$\begin{aligned} \nabla_{\dot{\gamma}} \nabla_{\dot{\gamma}} J^\top + R(J^\top, \dot{\gamma})\dot{\gamma} &= \nabla_{\dot{\gamma}} \nabla_{\dot{\gamma}} (g(J, \dot{\gamma})\dot{\gamma}) + R(g(J, \dot{\gamma})\dot{\gamma}, \dot{\gamma})\dot{\gamma} \\ &= g(\nabla_{\dot{\gamma}} \nabla_{\dot{\gamma}} J, \dot{\gamma})\dot{\gamma} \\ &= -g(R(J, \dot{\gamma})\dot{\gamma}, \dot{\gamma})\dot{\gamma} \\ &= 0. \end{aligned}$$

This shows that the tangential part J^\top of J is a Jacobi field. The fact that $\mathcal{J}_\gamma(TM)$ is a vector space implies that the normal part $J^\perp = J - J^\top$ of J also is a Jacobi field.

By differentiating $g(J, \dot{\gamma})$ twice along γ we obtain

$$\frac{d^2}{ds^2}(g(J, \dot{\gamma})) = g(\nabla_{\dot{\gamma}} \nabla_{\dot{\gamma}} J, \dot{\gamma}) = -g(R(J, \dot{\gamma})\dot{\gamma}, \dot{\gamma}) = 0$$

so $g(J, \dot{\gamma}(s)) = (as + b)$ for some $a, b \in \mathbb{R}$. \square

Corollary 9.9. *Let (M, g) be a Riemannian manifold, $\gamma : I \rightarrow M$ be a geodesic and J be a Jacobi field along γ . If*

$$g(J(t_0), \dot{\gamma}(t_0)) = 0 \quad \text{and} \quad g((\nabla_{\dot{\gamma}} J)(t_0), \dot{\gamma}(t_0)) = 0$$

for some $t_0 \in I$, then $g(J(t), \dot{\gamma}(t)) = 0$ for all $t \in I$.

PROOF. This is a direct consequence of the fact that the function $g(J, \dot{\gamma})$ satisfies the second order ordinary differential equation $\ddot{f} = 0$ and the initial conditions $f(0) = 0$ and $\dot{f}(0) = 0$. \square

Our next aim is to show that if the Riemannian manifold (M, g) has constant sectional curvature then we can completely solve the Jacobi equation

$$\nabla_X \nabla_X J + R(J, X)X = 0$$

along any given geodesic $\gamma : I \rightarrow M$. For this we introduce the following useful notation. For a real number $\kappa \in \mathbb{R}$ we define the functions $c_\kappa, s_\kappa : \mathbb{R} \rightarrow \mathbb{R}$ by

$$c_\kappa(s) = \begin{cases} \cosh(\sqrt{|\kappa|}s) & \text{if } \kappa < 0, \\ 1 & \text{if } \kappa = 0, \\ \cos(\sqrt{\kappa}s) & \text{if } \kappa > 0. \end{cases}$$

and

$$s_\kappa(s) = \begin{cases} \sinh(\sqrt{|\kappa|}s)/\sqrt{|\kappa|} & \text{if } \kappa < 0, \\ s & \text{if } \kappa = 0, \\ \sin(\sqrt{\kappa}s)/\sqrt{\kappa} & \text{if } \kappa > 0. \end{cases}$$

It is a well known fact that the unique solution to the initial value problem

$$\ddot{f} + \kappa \cdot f = 0, \quad f(0) = a \quad \text{and} \quad \dot{f}(0) = b$$

is the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(s) = a \cdot c_\kappa(s) + b \cdot s_\kappa(s)$.

We now give examples of Jacobi fields in the three model geometries of dimension two, the Euclidean plane, the sphere and hyperbolic plane, all of constant sectional curvature.

Example 9.10. Let \mathbb{C} be the complex plane equipped with the standard Euclidean metric $\langle \cdot, \cdot \rangle_{\mathbb{R}^2}$ of constant sectional curvature $\kappa = 0$. The rotations about the origin produce a 1-parameter family of geodesics $\Phi_t : s \mapsto s \cdot e^{it}$. Along the geodesic $\gamma_0 : s \mapsto s$ we yield the Jacobi field

$$J_0(s) = \frac{\partial \Phi_t}{\partial t}(0, s) = is$$

with $|J_0(s)|^2 = s^2 = |s_\kappa(s)|^2$.

Example 9.11. Let S^2 be the unit sphere in the standard three dimensional Euclidean space $\mathbb{C} \times \mathbb{R}$ equipped with the induced metric of constant sectional curvature $\kappa = +1$. Rotations about the \mathbb{R} -axis produce a 1-parameter family of geodesics $\Phi_t : s \mapsto (\sin(s) \cdot e^{it}, \cos(s))$. Along the geodesic $\gamma_0 : s \mapsto (\sin(s), \cos(s))$ we have the Jacobi field

$$J_0(s) = \frac{\partial \Phi_t}{\partial t}(0, s) = (i \sin(s), 0)$$

with $|J_0(s)|^2 = \sin^2(s) = |s_\kappa(s)|^2$.

Example 9.12. Let $B_1^2(0)$ be the open unit disk in the complex plane equipped with the hyperbolic metric

$$g(X, Y) = \frac{4}{(1 - |z|^2)^2} \langle \cdot, \cdot \rangle_{\mathbb{R}^2}$$

of constant sectional curvature $\kappa = -1$. Rotations about the origin produce a 1-parameter family of geodesics $\Phi_t : s \mapsto \tanh(s/2) \cdot e^{it}$. Along the geodesic $\gamma_0 : s \mapsto \tanh(s/2)$ we obtain the Jacobi field

$$J_0(s) = \frac{\partial \Phi_t}{\partial t}(0, s) = i \cdot \tanh(s/2)$$

with

$$|J_0(s)|^2 = \frac{4 \cdot \tanh^2(s/2)}{(1 - \tanh^2(s/2))^2} = \sinh^2(s) = |s_\kappa(s)|^2.$$

We are now ready to show that, in the case of constant sectional curvature, we can completely solve the Jacobi equation along any geodesic.

Example 9.13. Let (M, g) be a Riemannian manifold of constant sectional curvature κ and $\gamma : I \rightarrow M$ be a geodesic with $|X| = 1$ where $X = \dot{\gamma}$ is the tangent vector field along γ . Following Proposition 7.8 let P_1, P_2, \dots, P_{m-1} be parallel vector fields along γ such that

$$g(P_i, P_j) = \delta_{ij} \quad \text{and} \quad g(P_i, X) = 0.$$

Then any vector field J along γ may be written as

$$J(s) = \sum_{i=1}^{m-1} f_i(s) P_i(s) + f_m(s) X(s).$$

Since the vector fields $P_1, P_2, \dots, P_{m-1}, X$ are parallel along the curve γ , this means that J is a Jacobi field if and only if

$$\begin{aligned} \sum_{i=1}^{m-1} \ddot{f}_i(s) P_i(s) + \ddot{f}_m(s) X(s) &= \nabla_X \nabla_X J \\ &= -R(J, X)X \\ &= -R(J^\perp, X)X \\ &= -\kappa(g(X, X)J^\perp - g(J^\perp, X)X) \\ &= -\kappa J^\perp \\ &= -\kappa \sum_{i=1}^{m-1} f_i(s) P_i(s). \end{aligned}$$

This is equivalent to the following system of ordinary differential equations

$$(6) \quad \ddot{f}_m(s) = 0 \quad \text{and} \quad \ddot{f}_i(s) + \kappa f_i(s) = 0 \quad \text{for all } i = 1, 2, \dots, m-1.$$

It is clear that for the initial values

$$\begin{aligned} J(s_0) &= \sum_{i=1}^{m-1} v_i P_i(s_0) + v_m X(s_0), \\ (\nabla_X J)(s_0) &= \sum_{i=1}^{m-1} w_i P_i(s_0) + w_m X(s_0) \end{aligned}$$

or equivalently

$$f_i(s_0) = v_i \quad \text{and} \quad \dot{f}_i(s_0) = w_i \quad \text{for all } i = 1, 2, \dots, m$$

we have a unique explicit solution to the system (6) on the whole of the interval I . It is given by

$$f_m(s) = v_m + s w_m \quad \text{and} \quad f_i(s) = v_i c_\kappa(s) + w_i s_\kappa(s)$$

for all $i = 1, 2, \dots, m-1$. It should be noted that if $g(J, X) = 0$ and $J(0) = 0$ then

$$(7) \quad |J(s)| = |(\nabla_X J)(0)| \cdot |s_\kappa(s)|.$$

In the next example we give a complete description of the Jacobi fields along a geodesic on the 2-dimensional sphere.

Example 9.14. Let S^2 be the unit sphere in the three dimensional Euclidean space $\mathbb{C} \times \mathbb{R}$ equipped with the induced metric of constant sectional curvature $\kappa = +1$. Further let $\gamma : \mathbb{R} \rightarrow S^2$ be the geodesic given by $\gamma : s \mapsto (e^{is}, 0)$. Then the tangent vector field along γ satisfies

$$\dot{\gamma}(s) = (ie^{is}, 0).$$

It then follows from Proposition 9.8 that all the Jacobi fields tangent to γ are given by

$$J_{(a,b)}^T(s) = (as + b)(ie^{is}, 0),$$

where $a, b \in \mathbb{R}$. The unit vector field $P : \mathbb{R} \rightarrow TS^2$ given by

$$s \mapsto ((e^{is}, 0), (0, 1))$$

is clearly normal along γ . In S^2 the tangent vector field $\dot{\gamma}$ is parallel along γ so P must be parallel. This implies that all the Jacobi fields orthogonal to $\dot{\gamma}$ are given by

$$J_{(a,b)}^N(s) = (0, a \cos s + b \sin s),$$

where $a, b \in \mathbb{R}$.

In the general situation, when we do assume constant sectional curvature, the exponential map can be used to produce Jacobi fields as follows.

Example 9.15. Let (M, g) be a complete Riemannian manifold, $p \in M$ and $v, w \in T_p M$. Then $s \mapsto s(v + tw)$ defines a 1-parameter family of lines in the tangent space $T_p M$ which all pass through the origin $0 \in T_p M$. Remember that the exponential map

$$\exp_p|_{B_{\varepsilon_p}^m(0)} : B_{\varepsilon_p}^m(0) \rightarrow \exp_p(B_{\varepsilon_p}^m(0))$$

maps lines in $T_p M$ through the origin onto geodesics on M . Hence the map

$$\Phi_t : s \mapsto \exp_p(s(v + tw))$$

is a 1-parameter family of geodesics through $p \in M$, as long as $s(v + tw)$ is an element of $B_{\varepsilon_p}^m(0)$. This means that

$$J(s) = \frac{\partial \Phi_t}{\partial t}(t, s)|_{t=0} = d(\exp_p)_{s(v+tw)}(sw)|_{t=0} = d(\exp_p)_{sv}(sw)$$

is a Jacobi field along the geodesic $\gamma : s \mapsto \Phi_0(s)$ with $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. Here

$$d(\exp_p)_{s(v+tw)} : T_{s(v+tw)} T_p M \rightarrow T_{\exp_p(s(v+tw))} M$$

is the linear tangent map of the exponential map \exp_p at $s(v + tw)$. Now differentiating with respect to the parameter s gives

$$(\nabla_X J)(0) = \frac{d}{ds}(d(\exp_p)_{sv}(sw))|_{s=0} = d(\exp_p)_0(w) = w.$$

The above calculations show that

$$(8) \quad J(0) = 0 \quad \text{and} \quad (\nabla_X J)(0) = w.$$

For the proof of our main result, stated in Theorem 9.17, we need the following technical lemma.

Lemma 9.16. *Let (M, g) be a Riemannian manifold with sectional curvature uniformly bounded above by Δ and $\gamma : [0, \alpha] \rightarrow M$ be a geodesic on M with $|X| = 1$ where $X = \dot{\gamma}$. Further let $J : [0, \alpha] \rightarrow TM$ be a Jacobi field along γ such that $g(J, X) = 0$ and $|J| \neq 0$ on $(0, \alpha)$. Then*

- (i) $\frac{d^2}{ds^2}|J| + \Delta \cdot |J| \geq 0$,
- (ii) if $f : [0, \alpha] \rightarrow \mathbb{R}$ is a C^2 -function such that
 - (a) $\ddot{f} + \Delta \cdot f = 0$ and $f > 0$ on $(0, \alpha)$,
 - (b) $\dot{f}(0) = |J|(0)$, and
 - (c) $\dot{f}(0) = |\nabla_X J|(0)$,

- then $f(s) \leq |J(s)|$ on $(0, \alpha)$,
 (iii) if $J(0) = 0$, then $|\nabla_X J(0)| \cdot s_\Delta(s) \leq |J(s)|$ for all $s \in (0, \alpha)$.

PROOF. (i) Using the facts that $|X| = 1$ and $\langle X, J \rangle = 0$ we obtain

$$\begin{aligned}
 \frac{d^2}{ds^2}|J| &= \frac{d^2}{ds^2}\sqrt{g(J, J)} = \frac{d}{ds}\left(\frac{g(\nabla_X J, J)}{|J|}\right) \\
 &= \frac{g(\nabla_X \nabla_X J, J)}{|J|} + \frac{|\nabla_X J|^2 |J|^2 - g(\nabla_X J, J)^2}{|J|^3} \\
 &\geq \frac{g(\nabla_X \nabla_X J, J)}{|J|} \\
 &= -\frac{g(R(J, X)X, J)}{|J|} \\
 &= -K(X, J) \cdot |J| \\
 &\geq -\Delta \cdot |J|.
 \end{aligned}$$

(ii) Define the function $h : [0, \alpha) \rightarrow \mathbb{R}$ by

$$h(s) = \begin{cases} \frac{|J(s)|}{f(s)} & \text{if } s \in (0, \alpha), \\ \lim_{s \rightarrow 0} \frac{|J(s)|}{f(s)} = 1 & \text{if } s = 0. \end{cases}$$

Then

$$\begin{aligned}
 \dot{h}(s) &= \frac{1}{f^2(s)} \left\{ \frac{d}{ds} |J(s)| \cdot f(s) - |J(s)| \cdot \dot{f}(s) \right\} \\
 &= \frac{1}{f^2(s)} \int_0^s \frac{d}{dt} \left\{ \frac{d}{dt} |J(t)| \cdot f(t) - |J(t)| \cdot \dot{f}(t) \right\} dt \\
 &= \frac{1}{f^2(s)} \int_0^s \left\{ \frac{d^2}{dt^2} |J(t)| \cdot f(t) - |J(t)| \cdot \ddot{f}(t) \right\} dt \\
 &= \frac{1}{f^2(s)} \int_0^s f(t) \cdot \left\{ \frac{d^2}{dt^2} |J(t)| + \Delta \cdot |J(t)| \right\} dt \\
 &\geq 0.
 \end{aligned}$$

This implies that $\dot{h}(s) \geq 0$ so $f(s) \leq |J(s)|$ for all $s \in (0, \alpha)$.

(iii) The function $f(s) = |\nabla_X J(0)| \cdot s_\Delta(s)$ satisfies the differential equation

$$\ddot{f}(s) + \Delta f(s) = 0$$

and the initial conditions $f(0) = |J(0)| = 0$, $\dot{f}(0) = |\nabla_X J(0)|$ so it follows from (ii) that $|\nabla_X J(0)| \cdot s_\Delta(s) = f(s) \leq |J(s)|$ for all $s \in (0, \alpha)$. \square

Let (M, g) be a Riemannian manifold of sectional curvature which is uniformly bounded above, i.e. there exists a $\Delta \in \mathbb{R}$ such that $K_p(V) \leq \Delta$ for all $V \in G_2(T_p M)$ and $p \in M$. Let (M_Δ, g_Δ) be another Riemannian manifold which is complete and of constant sectional curvature $K \equiv \Delta$. Let $p \in M$, $p_\Delta \in M_\Delta$ and identify $T_p M \cong \mathbb{R}^m \cong T_{p_\Delta} M_\Delta$.

Let U be an open neighbourhood of \mathbb{R}^m around 0 such that the exponential maps $(\exp)_p$ and $(\exp)_{p_\Delta}$ are diffeomorphisms from U onto their images $(\exp)_p(U)$ and $(\exp)_{p_\Delta}(U)$, respectively. Let (r, p, q) be a geodesic triangle i.e. a triangle with sides which are shortest paths between their endpoints. Furthermore let $c : [a, b] \rightarrow M$ be the geodesic connecting r and q and $v : [a, b] \rightarrow T_p M$ be the curve defined by $c(t) = (\exp)_p(v(t))$. Put $c_\Delta(t) = (\exp)_{p_\Delta}(v(t))$ for $t \in [a, b]$ and then it directly follows that $c(a) = r$ and $c(b) = q$. Finally put $r_\Delta = c_\Delta(a)$ and $q_\Delta = c_\Delta(b)$.

Theorem 9.17. *For the above situation the following inequality for the distance function d is satisfied*

$$d(q_\Delta, r_\Delta) \leq d(q, r).$$

PROOF. Define a 1-parameter family $s \mapsto s \cdot v(t)$ of straight lines in $T_p M$ through 0. Then

$$\Phi_t : s \mapsto (\exp)_p(s \cdot v(t)) \quad \text{and} \quad \Phi_t^\Delta : s \mapsto (\exp)_{p_\Delta}(s \cdot v(t))$$

are 1-parameter families of geodesics through $p \in M$, and $p_\Delta \in M_\Delta$, respectively. Hence

$$J_t = \partial \Phi_t / \partial t \quad \text{and} \quad J_t^\Delta = \partial \Phi_t^\Delta / \partial t$$

are Jacobi fields satisfying the initial conditions

$$J_t(0) = 0 = J_t^\Delta(0) \quad \text{and} \quad (\nabla_X J_t)(0) = \dot{v}(t) = (\nabla_X J_t^\Delta)(0).$$

Employing Equation (7), Lemma 9.16 and the fact that M_Δ has constant sectional curvature Δ we now yield

$$\begin{aligned} |\dot{c}_\Delta(t)| &= |J_t^\Delta(1)| \\ &= |(\nabla_X J_t^\Delta)(0)| \cdot s_\Delta(1) \\ &= |(\nabla_X J_t)(0)| \cdot s_\Delta(1) \\ &\leq |J_t(1)| \\ &= |\dot{c}(t)| \end{aligned}$$

The curve c is the shortest path between r and q so we have

$$d(r_\Delta, q_\Delta) \leq L(c_\Delta) \leq L(c) = d(r, q).$$

□

We now add the assumption that the sectional curvature of the manifold (M, g) is uniformly bounded below i.e. there exists a $\delta \in \mathbb{R}$ such that $\delta \leq K_p(V)$ for all $V \in G_2(T_p M)$ and $p \in M$. Let (M_δ, g_δ) be a complete Riemannian manifold of constant sectional curvature δ . Let $p \in M$ and $p_\delta \in M_\delta$ and identify $T_p M \cong \mathbb{R}^m \cong T_{p_\delta} M_\delta$. Then a similar construction as above gives two pairs of points $q, r \in M$ and $q_\delta, r_\delta \in M_\delta$ and shows that

$$d(q, r) \leq d(q_\delta, r_\delta).$$

Combining these two results we obtain **locally**

$$d(q_\Delta, r_\Delta) \leq d(q, r) \leq d(q_\delta, r_\delta).$$

Exercises

Exercise 9.1. Find a proof of Lemma 9.7.

Exercise 9.2. Let (M, g) be a Riemannian manifold and $\gamma : I \rightarrow M$ be a geodesic such that $X = \dot{\gamma} \neq 0$. Further let J be a non-vanishing Jacobi field along γ with $g(X, J) = 0$. Prove that if $g(J, J)$ is constant along γ then (M, g) does not have strictly negative curvature.