

1

A Radioactive Element Has An Unstable
Nucleus

Unstable parent
nucleus

Daughter Atomic number:
2 fewer
Atomic mass:
4 fewer
Atomic number:
2 fewer
Atomic mass:
4 fewer
Atomic number:
2 fewer
Atomic mass:
4 fewer
Atomic number:
2 fewer
Atomic mass:
4 fewer

Parent isotope

Daughter isotope

After one half-life After two half-lives

Parent is the original unstable radioactive isotope

Daughter is the new, more stable isotope formed as a result of radioactive decay of parent

Over time, unstable parent isotopes decay to more stable daughter isotopes at a rate measured in half lives

3

Types of Radioactive DecayAlpha emissionBeta emissionElectron capture

Unstable parent nucleus

Daughter nucleus

2 fewer

Atomic number: 2 fewer


Atomic number: 4 fewer


Atomic number: 2 fewer


At


5


7/22/2024


















7/22/2024







13 14





15





17 18

TABLE 9.1 Isotopes Frequently Used in Radiometric Dating		
Radioactive Parent	Stable Daughter Product	Currently Accepted Half-Life Values
Uranium-238	Lead-206	4.5 billion years
Uranium-235	Lead-207	704 million years
Thorium-232	Lead-208	14.1 billion years
Rubidium-87	Strontium-87	47.0 billion years
Potassium-40	Argon-40	1.3 billion years

**Radiometric Dating** 

- If the following are known:
  - Half life of the isotope being measured
  - Parent/daughter ratio in the sample
- Then the age of the sample can be calculated

19 20

What is the age of a sample if the half-life is 5 million years and the...

Parent:Daughter ratio is 1:1?
25% Parent & 75% Daughter?
After four half-lives?

Smillion years
10 million years
20 million years

Mare one half-life


After two half-lives

After three half-lives


21 22


Resetting Isotopic Ages

The isotopic clock can be reset by metamorphism when temperatures become high enough for daughter isotopes to escape the minerals where they accumulated





23 24







25 26





27





29 30

#### **Isotopic Dating of Igneous Rocks**

- Igneous rocks can provide an approximate crystallization age using whole rock analysis
- Minerals also provide a crystallization age provided the rock has not experienced subsequent metamorphism


### Isotopic Dating of Metamorphic Rocks

- Minerals in a metamorphic rock can provide an age for the last metamorphic event
- If the rock experienced more than one episode of metamorphism, then usually only the most recent event can be dated

31 32

## Isotopic Dating of Sedimentary Rocks?


- In general, detrital (clastic) sedimentary rocks do not give meaningful ages because the minerals contained in these rocks were derived from other sources (parent rocks):
  - Clastic minerals in detrital sedimentary rocks only provide the age(s) of their parent source rocks
  - These ages do not reflect the time of sedimentation
- Multiple parent rocks of different ages can contribute sediment into the same basin:
  - This mixture of different ages in basin is meaningless



33

## Isotopic Dating of Sedimentary Rocks?


Isotopic dating of a sedimentary rock is possible if it contains authigenic minerals or material (e.g. glauconite, K-feldspar, organic matter) that formed during burial and diagenesis in the sedimentary basin



35 36

# **Bracketing the Ages of Sedimentary Rocks**

The ages of sedimentary rock packages can be bracketed by dating underlying and overlying igneous and metamorphic rocks, interlayered volcanic ash deposits and crosscutting igneous dikes.



37 38

#### **Dating With Radioactivity**

- Importance of radiometric dating:
  - Radiometric dating is a complex procedure that requires precise measurement
  - -Rocks from several localities have been dated at more than 3 billion years
  - -Confirms the idea that geologic time is immense

The Geologic
Time Scale

The Geologic Time Scale – a
"calendar" of Earth history:

• Subdivides geologic history into units

• Originally created using relative dating
• Absolute dates assigned later after the discovery of radioactivity in 1896

39 40

# The Geologic Time Scale Geologic Time Scale is subdivided into various units of time: Eons (largest divisions) Eras Periods Epochs (smallest divisions)

The Geologic
Time Scale

Eons constitute the largest divisions:


• Phanerozoic ("visible life"), most recent Eon, began about 541 million years ago and continues to present


• Proterozoic (2500 – 541 m.y. ago)

• Archean (4000 - 2500 m.y. ago)

• Hadean – oldest (4600 - 4000 m.y. ago)

41 42





43 44

