


Dots: every possible outcome for N = 5 replays and M = 2 controls, coloured by decide().

What counts as a pass, and how sure we are
A replay passes only if every originally failing target check now passes and no check that passed before has stopped passing.
Wilson 95% interval for k/N: 5 of 5 gives 57% to 100%; 0 of 5 gives 0% to 43%.
With five replays the interval is wide, and the report says so.

1 Did every replay and every control finish?
no: ERROR, no verdict is given; the finding returns to 'hypothesized'

2 k/N ≥ 0.8 and c/M ≤ 0.2 ?
yes: CONFIRMED, the fix (not luck) changed the outcome

3 k/N ≤ 0.2 ?
yes: REFUTED, fixing this step changes nothing, so it is not the origin

4 otherwise
INCONCLUSIVE: run more replays, or review with a written reason

decide(k, 5, c, 2) for every k and c:

	k = 0	k = 1	k = 2	k = 3	k = 4	k = 5
c = 0	refuted	refuted	inconclusive	inconclusive	confirmed	confirmed
c = 1	refuted	refuted	inconclusive	inconclusive	inconclusive	inconclusive
c = 2	refuted	refuted	inconclusive	inconclusive	inconclusive	inconclusive

Limits
N is at most 10 and M at most 5. A changed threshold is a new rule version (VR-2), never an edit of VR-1.