

MAJOR TECHNIQUES IN GENE MANIPULATION OF PLANTS



INTRODUCTION



- Plant genetic engineering has become one of the most important molecular tools in the modern molecular breeding of crops.
- Over the last decade, significant progress has been made in the development of new and efficient transformation methods in plants.
- Despite a variety of available DNA delivery methods, *Agrobacterium*- and biolistic-mediated transformation remain the two predominantly employed approaches.

• Production of transgenic plants

Isolate and clone gene of interest



Add DNA segments to initiate or enhance gene expression



Add selectable markers




Introduce gene construct into plant cells (transformation)




Select transformed cells or tissues



Regenerate whole plants



Plant Transformation Methods



Techniques for plant genetic transformation



- Indirect method- *Agrobacterium* mediated gene transfer
- Direct methods-
 - Particle bombardment (biolistics)
 - Microprojectile gun method
 - Electroporation
 - Silicon carbide fibres
 - Polyethylene glycol (PEG)/protoplast fusion
 - Liposome mediated gene transfer

Transformation vector requirements



- Origin of replication
- Bacterial selectable marker
- Gene constructs of interest
- T-DNA borders and other *Agrobacterium* genes if using *Agrobacterium*
- Compatible with helper plasmid if using *Agrobacterium*

Agrobacterium mediated gene transfer




Agrobacterium-

- Soil borne, gram negative, rod shaped, motile found in rhizosphere
- Causative agents of “Crown gall” disease of dicotyledones
- Have ability transfer bacterial genes to plant genome
- Attracted to wound site via chemotaxis in response to chemicals (sugar and Phenolic molecules: acetosyringone) released from damaged plant cells
- Contains Ti plasmid which can transfer its T-DNA region into genome of host plants


Ti-plasmid features



- Two strains of Ti-plasmid:
 - Octopine strains- contains two T-DNA region: T_L (14 kb) and T_R (7 kb)
 - Nopaline strains- contain one T-DNA region(20 kb)
- Size is about 200 kb
- Has a central role in Crown-gall formation
- Contains one or more T-DNA region that is integrated into the genome of host plants
- Contain a *vir* region ~ 40 kb at least 8~11 *vir* genes
- Has origin of replication
- Contains a region enabling conjugative transfer
- Has genes for the catabolism of opines



PASSAGE OF T-DNA FROM AGROBACTERIUM INTO PLANT NUCLEAR DNA




Electroporation technique.



- Is the process where by electrical impulses of high field strength are used to reversibly permeabilize cell membrane to facilitate uptake of large molecules, including DNA.
- It has been used for long time for transient and integrative transformation of protoplasts.
- 1 to 1.5 k V. so uses low capacitance hence short decay time .

Electroporation Technique




Biolistic/Particle bombardment



- High velocity micro projectile were utilize to deliver nucleic acids into living cells.
- **Advantages :**
 - Transformation of organized tissue
 - Universal delivery system
 - Transformation of recalcitrant spp
 - Study of basic plant development processes.

Particle Gun


The Helium Gas Gun – Circa 2000



Microinjection



- Under a [microscope](#), a cell is manipulated to a blunt capillary. Gentle suction holds the cell in place.
- With a [micromanipulator](#), a very fine tipped pipet is inserted into the cytoplasm or nucleus.
- DNA or RNA is injected directly into the nucleus or cytoplasm.
- Microinjection has been successfully used with large frog eggs, cultured mammalian cells, mammalian embryos, and plant protoplasts and tissues



Thank you

Three parallel white diagonal lines are located in the bottom right corner of the image, extending from the right edge towards the center.