


Characterization of 2-connected Graphs

Theorem []:


Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.




Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.




But . . .



this graph is 2-connected

and
here is
a u, v
path
 P




yet ???

Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

Proof (\Leftarrow) In this case, no pair of vertices u, v can be disconnected by the removal of a single vertex. Thus $\kappa(G) \geq 2$.



Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.

(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.

(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)


Recall

$$\kappa(G) \leq \kappa'(G) \leq \delta(G)$$

$$\rightarrow \kappa(G) \geq 2$$

$$\therefore \kappa'(G) \geq 2$$

\therefore no cut edge



Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

Let u, v be vertices with $d(u, v) = t \geq 2$;

assume true for vertices a distance $< t$ apart.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.

(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.


(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Let u, v be vertices with $d(u, v) = t \geq 2$;

assume true for vertices a distance $< t$ apart.

There is a u, v path R of length t . Let w be the predecessor of v on R



Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.


(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Let u, v be vertices with $d(u, v) = t \geq 2$;

assume true for vertices a distance $< t$ apart.

There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,

\exists internally disjoint u, w paths P, Q

Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.


(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Let u, v be vertices with $d(u, v) = t \geq 2$;

assume true for vertices a distance $< t$ apart.

There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,

\exists internally disjoint u, w paths P, Q

Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.


(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Let u, v be vertices with $d(u, v) = t \geq 2$;

assume true for vertices a distance $< t$ apart.

There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,
 \exists internally disjoint u, w paths P, Q

Also, w is not a cut vertex (why?)
so $G - w$ has a u, v path P'

Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.


(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Let u, v be vertices with $d(u, v) = t \geq 2$;

assume true for vertices a distance $< t$ apart.

There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,
 \exists internally disjoint u, w paths P, Q

Also, w is not a cut vertex (why?)
so $G - w$ has a u, v path P'

Characterization of 2-connected Graphs

Theorem [Whitney 1932]:

Graph G with $n(G) \geq 3$ is 2-connected if and only if for each pair u, v of distinct vertices, G has at least 2 internally disjoint u, v paths.

(\Rightarrow) Assume G is 2-connected. Proof by induction on $d(u, v)$.


(Basis) If $d(u, v) = 1$, uv is not a cut edge since $\kappa'(G) \geq \kappa(G) \geq 2$. Therefore $G - uv$ contains a (u, v) -path. The edge uv is the other path in G .

(Inductive step:)

Let u, v be vertices with $d(u, v) = t \geq 2$;

assume true for vertices a distance $< t$ apart.

There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,
 \exists internally disjoint u, w paths P, Q


Also, w is not a cut vertex (why?)
 so $G - w$ has a u, v path P'

Now we can get 2 internally disjoint u, v paths:

Let x be the last vertex of p' which is also on p or q .

then do this:

There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,
 \exists internally disjoint u, w paths p, q


Also, w is not a
cut vertex (why?)
so $G - w$ has a u, v
path p'

Now we can get 2 internally disjoint u, v paths:

Let x be the last vertex of p' which is also on p or q .

then do this:

There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,
 \exists internally disjoint u, w paths p, q


Also, w is not a
cut vertex (why?)
so $G - w$ has a u, v
path p'

Now we can get 2 internally disjoint u, v paths:

Let x be the last vertex of p' which is also on p or q .

then do this:


There is a u, v path R of length t . Let w be the predecessor of v on R



$d(u, w) < t$, so by induction,
 \exists internally disjoint u, w paths p, q

Also, w is not a
cut vertex (why?)
so $G - w$ has a u, v
path p'

Expansion Lemma 4.2.3. If G is k -connected and G' is obtained from G by adding a new vertex y adjacent to at least k vertices of G , then G' is k -connected.



G' is k -connected

Expansion Lemma 4.2.3. If G is k -connected and G' is obtained from G by adding a new vertex y adjacent to at least k vertices of G , then G' is k -connected.


Proof. Suppose S separates G' .

(Show $|S| \geq k$)

If $y \in S$, then $S - y$ separates G . Thus, $|S - y| \geq k$, i.e., $|S| \geq k + 1$.


If $y \notin S$, then if $N(y) \subseteq S$ then $|S| \geq |N(y)| = k$.

Otherwise, S separates 2 vertices of G and therefore $|S| \geq k$.



G' is k -connected

If S disconnects G' then one of these:




$y \in S$

Then $S-y$
disconnects G


So $|S-y| \geq k$

$\therefore |S| \geq k+1$



$y \notin S$ but
 $N(y) \subseteq S$

Then $|S| \geq |N(y)| = k$



otherwise

Removing S
disconnects G

$\therefore |S| \geq k$

Theorem 4.2.4. If $n(G) \geq 3$, the f.a.e.

(A) G is 2-connected

(B) For all $x, y \in V(G)$, there are internally disjoint x, y -paths.

(C) For all $x, y \in V(G)$, there is a cycle containing x and y .

(D) $\delta(G) \geq 1$ and every two edges of G lie on a common cycle.

Theorem 4.2.4. If $n(G) \geq 3$, the f.a.e.

(A) G is 2-connected

(B) For all $x, y \in V(G)$, there are internally disjoint x, y -paths.

(C) For all $x, y \in V(G)$, there is a cycle containing x and y .

(D) $\delta(G) \geq 1$ and every two edges of G lie on a common cycle.

$A \Leftrightarrow B$ is
Whitney's Theorem

Theorem 4.2.4. If $n(G) \geq 3$, the f.a.e.


(A) G is 2-connected

(B) For all $x, y \in V(G)$, there are internally disjoint x, y -paths.

(C) For all $x, y \in V(G)$, there is a cycle containing x and y .

(D) $\delta(G) \geq 1$ and every two edges of G lie on a common cycle.

$B \Leftrightarrow C$



Theorem 4.2.4. If $n(G) \geq 3$, the f.a.e.

(A) G is 2-connected


(B) For all $x, y \in V(G)$, there are internally disjoint x, y -paths.

(C) For all $x, y \in V(G)$, there is a cycle containing x and y .

(D) $\delta(G) \geq 1$ and every two edges of G lie on a common cycle.

Every 2 vertices lie on a cycle
 \Updownarrow
Every 2 edges lie on a cycle

$D \rightarrow C$ is easy




For $C \rightarrow D$
need "expansion lemma"

Given $A \Leftrightarrow B \Leftrightarrow C$, show $C \Rightarrow D$.

Assume any 2 vertices of G lie on a common cycle.

Then G is 2 connected (since $A \Leftrightarrow C$).

Let uv and wx be any 2 edges of G .




Given $A \Leftrightarrow B \Leftrightarrow C$, show $C \Rightarrow D$.

Assume any 2 vertices of G lie on a common cycle.

Then G is 2 connected (since $A \Leftrightarrow C$).

Let uv and wx be any 2 edges of G .

Use expansion lemma like this:



Result is 2-connected, so a & b lie on a common cycle, which must contain (consecutively) w, x and

u, v . Now

$u, b, x \rightarrow wx$

$u, a, v \rightarrow uv$


Given $A \Leftrightarrow B \Leftrightarrow C$, show $C \Rightarrow D$.

Assume any 2 vertices of G lie on a common cycle.

Then G is 2 connected (since $A \Leftrightarrow C$).

Let uv and wx be any 2 edges of G .

Use expansion lemma like this:



Result is 2-connected, so
 a & b lie on a common
cycle, which must contain
(consecutively) w, x and

u, v . Now

$w, b, x \rightarrow wx$

$u, a, v \rightarrow uv$


Given $A \Leftrightarrow B \Leftrightarrow C$, show $C \Rightarrow D$.

Assume any 2 vertices of G lie on a common cycle.

Then G is 2 connected (since $A \Leftrightarrow C$).

Let uv and wx be any 2 edges of G .

Use expansion lemma like this:



Result is 2-connected, so
 a & b lie on a common
cycle, which must contain
(consecutively) w, x and

u, v . Now

$w, x \rightarrow wx$

$u, v \rightarrow uv$

Can generalize:

Menger's Theorem: Graph G with at least $k + 1$ vertices is k -connected iff for every pair of vertices u, v of G , there are at least k pairwise internally disjoint (u, v) -paths.

Also: k -edge-connected iff at least k pairwise edge-disjoint u, v -paths.

