

Coloring Planar Graphs

(so assume graphs are simple)

ex 1

Show if G is simple, planar then $\delta(G) \leq 5$

ex. 2 Show that planar graphs are 6-colorable

The Five Color Theorem

Theorem 6.3.1. [Heawood 1890] Every planar graph is 5-colorable.

The Five Color Theorem

Theorem 6.3.1. [Heawood 1890] Every planar graph is 5-colorable.

Pf. Induction on $n(G)$. Basis $n(G) \leq 5$, clear.

Assume G is planar and $n(G) > 5$.

The Five Color Theorem

Theorem 6.3.1. [Heawood 1890] Every planar graph is 5-colorable.

Pf. Induction on $n(G)$. Basis $n(G) \leq 5$, clear.


Assume G is planar and $n(G) > 5$.

Then G has a vertex v of degree ≤ 5 .

By induction $G - v$ is 5-colorable, so 5-color $G - v$.

The Five Color Theorem

Theorem 6.3.1. [Heawood 1890] Every planar graph is 5-colorable.



Pf. Induction on $n(G)$. Basis $n(G) \leq 5$, clear.

Assume G is planar and $n(G) > 5$.


Then G has a vertex v of degree ≤ 5 .

By induction $G - v$ is 5-colorable, so 5-color $G - v$.

If $d_G(v) = 4$ or if $d_G(v) = 5$, but neighbors of v use only 4 colors, then G is 5-colorable.

So, suppose $d_G(v) = 5$ and each nbr of v has different color.

Planar embed G and label nbrs of v clockwise v_1, v_2, v_3, v_4, v_5



Let $G_{i,j}$ be subgraph induced by vertices colored i and j




So, suppose $d_G(v) = 5$ and each nbr of v has different color.

Planar embed G and label nbrs of v clockwise v_1, v_2, v_3, v_4, v_5

Let $G_{i,j}$ be subgraph induced by vertices colored i and j

e.g. $G_{2,4}$




So, suppose $d_G(v) = 5$ and each nbr of v has different color.

Planar embed G and label nbrs of v clockwise v_1, v_2, v_3, v_4, v_5

Let $G_{i,j}$ be subgraph induced by vertices colored i and j

e.g. $G_{2,4}$

Are $v_2 + v_4$ in same component?



So, suppose $d_G(v) = 5$ and each nbr of v has different color.

Planar embed G and label nbrs of v clockwise v_1, v_2, v_3, v_4, v_5


Let $G_{i,j}$ be subgraph induced by vertices colored i and j

e.g. $G_{2,4}$

Are $v_2 + v_4$ in same component?

If not

Swap colors in one of the components




So, suppose $d_G(v) = 5$ and each nbr of v has different color.

Planar embed G and label nbrs of v clockwise v_1, v_2, v_3, v_4, v_5

Let $G_{i,j}$ be subgraph induced by vertices colored i and j

e.g. $G_{2,4}$ Are $v_2 + v_4$ in same component?




If not

Swap colors in one of the components

So, suppose $d_G(v) = 5$ and each nbr of v has different color.

Planar embed G and label nbrs of v clockwise v_1, v_2, v_3, v_4, v_5



Let $G_{i,j}$ be subgraph induced by vertices colored i and j

e.g. $G_{2,4}$

Are $v_2 + v_4$ in same component?

If not

Swap colors in one of the components and make a color available for v , giving a 5-coloring of G .



So, suppose $d_G(v) = 5$ and each nbr of v has different color.


Planar embed G and label nbrs of v clockwise v_1, v_2, v_3, v_4, v_5



Let $G_{i,j}$ be subgraph induced by vertices colored i and j

e.g. $G_{2,4}$

Are $v_2 + v_4$ in same component?



If not




Swap colors in one of the components and make a color available for v , giving a 5-coloring of G .

But what if v_i connected to v_j in every $G_{i,j}$?


Let $G_{i,j}$ be subgraph induced by vertices colored i and j

Suppose in every $G_{i,j}$ there is path $P_{i,j}$ joining v_i and v_j



Let $G_{i,j}$ be subgraph induced by vertices colored i and j


Suppose in every $G_{i,j}$ there is path $P_{i,j}$ joining v_i and v_j



$P_{1,3} + v_1 v_2 + v_2 v_3$
forms a cycle C

v_2 and v_4 are on opposite sides (int/ext)

Let $G_{i,j}$ be subgraph induced by vertices colored i and j



Suppose in every G_{ij} there is path P_{ij} joining v_i and v_j


$P_{13} + N_{N_1} + N_{N_3}$
forms a cycle C

N_2 and N_4 are on opposite sides (int/ext)

1. 
2. 
3. 
4. 
5. 

Let $G_{i,j}$ be subgraph induced by vertices colored i and j

Suppose in every $G_{i,j}$ there is path $P_{i,j}$ joining v_i and v_j



$P_{1,3} + v_1 v_2 + v_2 v_3$
forms a cycle C

v_2 and v_4 are on opposite sides (int/ext)

Let $G_{i,j}$ be subgraph induced by vertices colored i and j


Suppose in every $G_{i,j}$ there is path $P_{i,j}$ joining v_i and v_j




$P_{1,3} + v_1 v_2 + v_2 v_3$
forms a cycle C

v_2 and v_4 are on opposite sides (int/ext)

So $P_{2,4}$ must cross C (not at a vertex) contradicting that the embedding is planar.



Let $G_{i,j}$ be subgraph induced by vertices colored i and j



Suppose in every $G_{i,j}$ there is path P_{ij} joining v_i and v_j



$P_{13} + v_1 v_5 + v_5 v_3$
forms a cycle C

v_2 and v_4 are on opposite sides (int/ext)

So P_{24} must cross C (not at a vertex) contradicting that the embedding is planar.

Thm 6.3.6 [1977 Appel & Haken]

For a planar graph G ,

Thm 6.3.6 [1977 Appel & Haken]

For a planar graph G ,

