Synthesis, Biological Evaluation of Novel Compounds 2-Substituted-6-(furan-2-yl)-6H-chromeno-(4,3-b)quinoline and their Derivatives

M. Komal Reddy*, Samarla Surender# and Baithy Mallesham

Department of Chemistry, Satavahana University, Karimnagar -505 002, Telangana, India

komalreddy@yahoo.com, surendersamarla89@gmail.com

Quinolines are significantly important compounds due to their wide range of pharmacological activities behaving like anti-malarial, anti-inflammatory, anti-asthmatic, anti-hypertensive and tyrosine kinase inhibiting agents. In addition to medicinal applications, quinoline derivatives are found to undergo hierarchical self-assembly into a variety of nano-structures and meso-structures with enhanced electronic and photonic functions. 2-Substituted-6-(furan-2-yl)-6H-chromeno-(4,3-b)quinoline were synthesized by condensation between E-3-(furan-2-yl)-1-(2-hydroxy-5-substituted phenyl) prop-2-en-1-one and orthoaminobenzaldehyde in presence of acedic medium to give good yield percentage.

All the synthesized compounds were characterised by ^{1}H-NMR, ^{13}C-NMR, IR, MASS-spectral studies and elemental analysis. All the synthesized compounds were subjected to biological evaluation studies. More information on this examination will be described in this presentation.



R= H, NO_2 ,OH, Br,CH_3,CH_3O

X= O,S.N