

Week 7: The fundamental theorem of Galois theory (14.1, 14.2)

On this homework, you may assume without proof that $\mathbb{Q}(\zeta_n)/\mathbb{Q}$ is Galois with $(\mathbb{Z}/n\mathbb{Z})^\times \cong \text{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$, with the isomorphism given by $a \mapsto (\zeta_n \mapsto \zeta_n^a)$. We will prove this result next week.

Practice Problems

1. Let $\tau: \mathbb{C} \rightarrow \mathbb{C}$ be given by complex conjugation. Show directly that τ is a field automorphism of \mathbb{C} . What is the fixed field of τ ?
2. Determine the lattice of subfields of $\mathbb{Q}(\zeta_5)$. *Hint:* $\sqrt{5} = \zeta_5 - \zeta_5^2 - \zeta_5^3 + \zeta_5^4$.
3. Determine the lattice of subfields of $\mathbb{Q}(\zeta_8)$ and $\mathbb{Q}(\zeta_{12})$.

Presentation Problems

1. Let $\alpha = \sqrt{2 + \sqrt{2}}$. Compute $\text{Gal}(\mathbb{Q}(\alpha)/\mathbb{Q})$.
2. Let $\tau: \mathbb{Q}(\zeta_n) \rightarrow \mathbb{Q}(\zeta_n)$ be given by complex conjugation. What is a primitive element for the fixed field of τ ? *Hint:* For the hard direction, use the fact that ζ_n satisfies the quadratic $x^2 - (\zeta_n + \zeta_n^{-1})x + 1 = 0$.
3. Determine the lattice of subfields of $\mathbb{Q}(\zeta_{20})$.
4. Let $L/K/F$ be a tower of field extension with L/F Galois. Let $G = \text{Gal}(L/F)$, let $H = \text{Gal}(L/K)$, and let G/H denote the collection of left cosets of H in G .
 - (a) Show that for each left coset $C \in G/H$ and each $\alpha \in K$, the value of $C(\alpha)$ does not depend on the choice of an element of C .

For $\alpha \in K$, the trace and norm of α from K to F are defined by

$$\text{Tr}_{K/F}(\alpha) = \sum_{C \in G/H} C(\alpha), \quad \text{N}_{K/F}(\alpha) = \prod_{C \in G/H} C(\alpha).$$

- (b) Let $\alpha \in K$. Show that $\text{Tr}_{K/F}(\alpha) \in F$ and $\text{N}_{K/F}(\alpha) \in F$.
- (c) Show that $\text{Tr}_{K/F}$ is additive and $\text{N}_{K/F}$ is multiplicative.

Tricky Problems

1. Existence of abelian Galois groups.
2. Let $n \geq 1$, let $K = \mathbb{Q}(\sqrt[n]{2})$, let $L = \mathbb{Q}(\zeta_n)$, and let $M = KL = \mathbb{Q}(\sqrt[n]{2}, \zeta_n)$.
 - (a) Show that $f(x) = x^n - 2$ is irreducible and that $[K : \mathbb{Q}] = n$.
 - (b) Show that M is a splitting field of $f(x)$ over \mathbb{Q} .
 - (c) Show that if F is a subfield of K then $F = \mathbb{Q}(\sqrt[d]{2})$ for some $d \mid n$. *Hint:* Let $d = [F : \mathbb{Q}]$ and show that $\text{N}_{K/F}(\sqrt[n]{2}) = \sqrt[d]{2}$.
 - (d) Show that if F is a subfield of L then F is Galois over \mathbb{Q} .
 - (e) Show that

$$K \cap L = \begin{cases} \mathbb{Q} & 8 \nmid n, \\ \mathbb{Q}(\sqrt{2}) & 8 \mid n. \end{cases}$$

(f) Use the isomorphism $\text{Gal}(KL/K) \cong \text{Gal}(L/(K \cap L))$ to show that

$$[M : \mathbb{Q}] = \begin{cases} n\varphi(n) & 8 \nmid n, \\ n\varphi(n)/2 & 8 \mid n. \end{cases}$$

(g) Show that if $8 \nmid n$ then $\text{Gal}(M/\mathbb{Q}) \cong \mathbb{Z}/n\mathbb{Z} \rtimes_{\varphi} (\mathbb{Z}/n\mathbb{Z})^{\times}$ where $\varphi: (\mathbb{Z}/n\mathbb{Z})^{\times} \rightarrow \text{Aut}(\mathbb{Z}/n\mathbb{Z})$ is the multiplication isomorphism. *Hint:* Produce an injective group homomorphism from $\text{Gal}(M/\mathbb{Q})$ to $\mathbb{Z}/n\mathbb{Z} \rtimes_{\varphi} (\mathbb{Z}/n\mathbb{Z})^{\times}$ by considering the action on $\sqrt[n]{2}$ and ζ_n .

3. Let $\alpha = \sqrt{(2 + \sqrt{2})(3 + \sqrt{3})}$. Compute $\text{Gal}(\mathbb{Q}(\alpha)/\mathbb{Q})$.