

Week 1: Ring homomorphisms, quotient rings, and ideals

Let R be a commutative ring with identity.

Practice Problems

1. Let R be the collection of continuous functions on $[0, 1]$. Show that R is a commutative ring with identity. Let $x \in [0, 1]$ and let $I = \{f \in R : f(x) = 0\}$. Show that I is a prime ideal of R .
2. Let I and J be ideals of R .
 - (a) Show that $I + J$ and $I \cap J$ and IJ are ideals of R .
 - (b) Show that $IJ \subseteq I \cap J \subseteq I + J$.
 - (c) Give an example where $IJ \neq I \cap J$.
3. Let $R = \mathbb{C}[x]$ and let I be the collection of polynomials in R with no constant or linear term. Show that I is an ideal of R . Show that R is an integral domain but R/I is not an integral domain.

Presentation Problems

1.
 - (a) Let I, J , and K be ideals of R . Show that if $I \subseteq J \cup K$ then $I \subseteq J$ or $I \subseteq K$.
 - (b) Let P be a prime ideal of R and I, J be ideals of R . Show that if $IJ \subseteq P$ then $I \subseteq P$ or $J \subseteq P$.
2.
 - (a) Show that if I is an ideal of R then $\sqrt{I} = \{x \in R : x^m \in I \text{ for some } m \geq 1\}$ is an ideal of R .
 - (b) The *nilradical* of R is defined by $\mathfrak{N}(R) = \sqrt{0}$. A ring S satisfying $\mathfrak{N}(S) = 0$ is called *reduced*. Show that $R/\mathfrak{N}(R)$ is reduced.

We now show that $\mathfrak{N}(R)$ is the intersection of all prime ideals of R .

- (c) Show that $\mathfrak{N}(R)$ is contained in the intersection of the prime ideals of R .
 - (d) Let $x \in R \setminus \mathfrak{N}(R)$ and let $S = \{x^n : n \geq 1\}$. Apply Zorn's Lemma to the collection of ideals of R disjoint from S to obtain a prime ideal P of R with $x \notin P$.
 - (e) Deduce that $\mathfrak{N}(R)$ is the intersection of the prime ideals of R .
 - (f) Let I be an ideal of R . Show that \sqrt{I} is the intersection of the prime ideals of R containing I .
3. Let J, I_1, \dots, I_n be ideals of R . Suppose that I_k is a prime ideal of R for all $k \geq 3$. The prime avoidance lemma states that if $J \subseteq I_1 \cup \dots \cup I_n$ then $J \subseteq I_k$ for some k .
 - (a) Show that the prime avoidance lemma holds when $n = 2$.

Now suppose that $n \geq 3$ and inductively assume that the prime avoidance lemma holds for all smaller values of n . Suppose for contradiction that $J \subseteq I_1 \cup \dots \cup I_n$ but $J \not\subseteq I_k$ for all k .

- (b) Use the inductive hypothesis to find $x_k \in J \setminus (I_1 \cup \dots \cup I_{k-1} \cup I_{k+1} \cup \dots \cup I_n)$.
- (c) Show that $x_k \in I_k$.
- (d) Use the primality of I_n to show that $x_1 \cdots x_{n-1} + x_n \in J \setminus (I_1 \cup \dots \cup I_n)$. Derive a contradiction.

This proves the prime avoidance lemma.

4. Let R be the ring of continuous functions on $[0, 1]$. For each $x \in [0, 1]$, let $I_x = \{f \in R : f(x) = 0\}$.
 - (a) Prove that each I_x is a maximal ideal of R .
 - (b) Prove that every maximal ideal of R is of the form I_x for some $x \in I$.

Module Theory Problem

1. A sequence of R -modules is a diagram of R -modules of the form

$$\cdots \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0 \xrightarrow{f_0} M_{-1} \xrightarrow{f_{-1}} \cdots$$

If only finitely many M_n are nonzero, we only include one of the zero terms on each side. For example, if $M_n = 0$ for $n > 2$ and $n < -2$, then we would draw the above as

$$0 \longrightarrow M_2 \xrightarrow{f_2} M_1 \xrightarrow{f_1} M_0 \xrightarrow{f_0} M_{-1} \xrightarrow{f_{-1}} M_{-2} \longrightarrow 0.$$

A sequence of R -modules is called a chain complex if $f_{n-1} \circ f_n = 0$ for all n , i.e. $\text{im } f_n \subseteq \ker f_{n-1}$. A sequence is called exact if $\text{im } f_n = \ker f_{n-1}$.

Let $R = \mathbb{R}$, and let $C^\infty(\mathbb{R}^n, \mathbb{R}^k)$ denote the vector space (R -module) of infinitely differentiable functions from \mathbb{R}^n to \mathbb{R}^k . Come up with an exact sequence of R -modules

$$0 \longrightarrow \mathbb{R} \longrightarrow C^\infty(\mathbb{R}^3, \mathbb{R}) \longrightarrow C^\infty(\mathbb{R}^3, \mathbb{R}^3) \longrightarrow C^\infty(\mathbb{R}^3, \mathbb{R}^3) \longrightarrow C^\infty(\mathbb{R}^3, \mathbb{R}) \longrightarrow 0.$$

Hint: Where have you heard the term “exact” before?

2. A short exact sequence is an exact sequence of the form

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

What property will the map f have? What property will the map g have? Show that A is isomorphic to a submodule of B and that C is isomorphic to a quotient of B .

3. We say a short exact sequence (as above) left-splits if there is a map $r: B \rightarrow A$ such that $r \circ f = \text{id}_A$. We say it right-splits if there is a map $s: C \rightarrow B$ such that $g \circ s = \text{id}_C$. Prove the following are equivalent:

- (a) The short exact sequence left-splits.
- (b) The short exact sequence right-splits.
- (c) There is an isomorphism $\varphi: B \rightarrow A \oplus C$ and a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{g} & C \longrightarrow 0 \\ & & \downarrow \text{id} & & \downarrow \varphi & & \downarrow \text{id} \\ 0 & \longrightarrow & A & \xrightarrow{\iota} & A \oplus C & \xrightarrow{\pi} & C \longrightarrow 0 \end{array}$$

where ι and π are the inclusion and projection maps.

This is known as the splitting lemma.

4. Let k be a field and let V and W be finite dimensional vector spaces over k . Let $L: V \rightarrow W$ be a linear transformation. Prove that $\dim V = \text{rk } L + \text{null } L$.

This is known as the rank-nullity theorem.

Tricky Problems

1. Ideals I and J of R are called coprime if $I + J = R$.

(a) Show that if I and J are coprime, then $I \cap J = IJ$.

(b) Show that if I and J are coprime, then $R/(I \cap J) \cong R/I \times R/J$.

(c) Show that if I_1, \dots, I_n are pairwise coprime then $R/(I_1 \cap \dots \cap I_k) \cong R/I_1 \times \dots \times R/I_k$.

This is known as the Chinese remainder theorem (2/2).

Now let n be a positive integer and let $n = p_1^{a_1} \dots p_k^{a_k}$ be the prime factorization of n .

(d) Show that $\mathbb{Z}/n\mathbb{Z} \cong \mathbb{Z}/p_1^{a_1}\mathbb{Z} \times \dots \times \mathbb{Z}/p_k^{a_k}\mathbb{Z}$.

(e) Show that $(\mathbb{Z}/n\mathbb{Z})^\times \cong (\mathbb{Z}/p_1^{a_1}\mathbb{Z})^\times \times \dots \times (\mathbb{Z}/p_k^{a_k}\mathbb{Z})^\times$.

2. Let $\text{Spec } R$ denote the set of prime ideals of R . Given an ideal I of R , let $V(I) = \{P \in \text{Spec } R : I \subseteq P\}$. Prove that the following are equivalent.

(a) There are ideals I and J of R such that $V(I)$ and $V(J)$ are disjoint and $V(I) \cup V(J) = \text{Spec } R$.

(b) There exist nonzero idempotents $e_1, e_2 \in R$ such that $e_1 + e_2 = 1$ and $e_1 e_2 = 0$.

(c) R is isomorphic to a direct product $R_1 \times R_2$ of two nonzero rings (both commutative with identity).

If you know about topological spaces, show that we can put a topology on $\text{Spec } R$ by defining closed sets to be sets of the form $V(I)$. What does condition (a) say topologically?