

# CS 611: Theory of Computation

Hongmin Li

Department of Computer Science  
California State University, East Bay

# Finite Languages

## Definition

A language is finite if it has finitely many strings.

# Finite Languages

## Definition

A language is finite if it has finitely many strings.

## Example

$\{0, 1, 00, 10\}$  is a finite language

# Finite Languages

## Definition

A language is finite if it has finitely many strings.

## Example

$\{0, 1, 00, 10\}$  is a finite language, however,  $(00 \cup 11)^*$  is not.

# Finiteness and Regularity

## Proposition

*If  $L$  is finite then  $L$  is regular.*

# Finiteness and Regularity

## Proposition

*If  $L$  is finite then  $L$  is regular.*

## Proof.

Let  $L = \{w_1, w_2, \dots, w_n\}$ . Then  $R = w_1 \cup w_2 \cup \dots \cup w_n$  is a regular expression defining  $L$ . □

# Are all languages regular?

# Are all languages regular?

## Proposition

*The language*

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  *is not regular.*



# Are all languages regular?

## Proposition

*The language*

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  *is not regular.*

## Proof

No DFA has enough states to keep track of the number of 0s and 1s it might see. □

# Are all languages regular?

## Proposition

*The language*

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  *is not regular.*

## Proof?

No DFA has enough states to keep track of the number of 0s and 1s it might see. □

Above is a weak argument because  $E = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 01 and 10 substrings}\}$  is regular!

# Proving Non-Regularity

## Proposition

*The language*

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  *is not regular.*

# Proving Non-Regularity

## Proposition

*The language*

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  *is not regular.*

## Proof.

Suppose (for contradiction)  $L_{\text{eq}}$  is recognized by DFA  $M = (Q, \{0,1\}, \delta, q_0, F)$ , where  $|Q| = n$ .

# Proving Non-Regularity

## Proposition

*The language*

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  *is not regular.*

## Proof.

Suppose (for contradiction)  $L_{\text{eq}}$  is recognized by DFA

$M = (Q, \{0,1\}, \delta, q_0, F)$ , where  $|Q| = n$ .

- There must be  $j < k \leq n$  such that  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k)$  ( $= q$  say).

# Proving Non-Regularity

## Proposition

*The language*

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  *is not regular.*


## Proof.

Suppose (for contradiction)  $L_{\text{eq}}$  is recognized by DFA  $M = (Q, \{0,1\}, \delta, q_0, F)$ , where  $|Q| = n$ .

- There must be  $j < k \leq n$  such that  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k)$  ( $= q$  say).
- Let  $x = 0^j$ ,  $y = 0^{k-j}$ , and  $z = 0^{n-k}1^n$ ; so  $xyz = 0^n1^n$ .  $\rightarrow$

# Proving Non-Regularity


Proof (contd).



- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$

# Proving Non-Regularity

Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .



# Proving Non-Regularity

Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\hat{\delta}(q_0, 0^n 1^n) = \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n)$$

# Proving Non-Regularity

Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\hat{\delta}(q_0, 0^n 1^n) = \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n) \quad (\text{since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v))$$

# Proving Non-Regularity

Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\begin{aligned}\hat{\delta}(q_0, 0^n 1^n) &= \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n) && \text{(since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v) \text{)} \\ &= \hat{\delta}(\hat{\delta}(q_0, 0^j), 0^{n-k} 1^n)\end{aligned}$$

# Proving Non-Regularity

Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\begin{aligned}\hat{\delta}(q_0, 0^n 1^n) &= \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n) && \text{(since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v) \text{)} \\ &= \hat{\delta}(\hat{\delta}(q_0, 0^j), 0^{n-k} 1^n) && (\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k))\end{aligned}$$

# Proving Non-Regularity

## Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\begin{aligned}\hat{\delta}(q_0, 0^n 1^n) &= \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n) && \text{(since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v) \text{)} \\ &= \hat{\delta}(\hat{\delta}(q_0, 0^j), 0^{n-k} 1^n) && (\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k)) \\ &= \hat{\delta}(q_0, 0^{n-k+j} 1^n)\end{aligned}$$

# Proving Non-Regularity

## Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\begin{aligned}
 \hat{\delta}(q_0, 0^n 1^n) &= \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n) && (\text{since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v)) \\
 &= \hat{\delta}(\hat{\delta}(q_0, 0^j), 0^{n-k} 1^n) && (\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k)) \\
 &= \hat{\delta}(q_0, 0^{n-k+j} 1^n) && (\text{since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v))
 \end{aligned}$$

# Proving Non-Regularity

## Proof (contd).




- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\begin{aligned}
 \hat{\delta}(q_0, 0^n 1^n) &= \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n) && (\text{since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v)) \\
 &= \hat{\delta}(\hat{\delta}(q_0, 0^j), 0^{n-k} 1^n) && (\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k)) \\
 &= \hat{\delta}(q_0, 0^{n-k+j} 1^n) && (\text{since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v))
 \end{aligned}$$

- So  $M$  accepts  $0^{n-k+j} 1^n$  as well.

# Proving Non-Regularity

## Proof (contd).



- We have  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^n 1^n \in L_{\text{eq}}$ ,  $\hat{\delta}(q_0, 0^n 1^n) \in F$ .

$$\begin{aligned}
 \hat{\delta}(q_0, 0^n 1^n) &= \hat{\delta}(\hat{\delta}(q_0, 0^k), 0^{n-k} 1^n) && (\text{since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v)) \\
 &= \hat{\delta}(\hat{\delta}(q_0, 0^j), 0^{n-k} 1^n) && (\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k)) \\
 &= \hat{\delta}(q_0, 0^{n-k+j} 1^n) && (\text{since } \hat{\delta}(q, uv) = \hat{\delta}(\hat{\delta}(q, u), v))
 \end{aligned}$$

- So  $M$  accepts  $0^{n-k+j} 1^n$  as well. But,  $0^{n-k+j} 1^n \notin L_{\text{eq}}$ ! □



# Pumping Lemma: Overview

## Pumping Lemma

The lemma generalizes this argument. Gives the template of an argument that can be used to easily prove that many languages are non-regular.

# Pumping Lemma

## The Statement

### Lemma

*If  $L$  is regular then there is a number  $p$  (the pumping length) such that  $\forall w \in L$  with  $|w| \geq p$ ,  $\exists x, y, z \in \Sigma^*$  such that  $w = xyz$  and*

$\exists p,$

$\forall w \in L$  with  $|w| \geq p,$

$\exists x, y, z \in \Sigma^*$  such that  $w = xyz$ ,  $|y| > 0$  and  $|xy| \leq p,$

$\forall i \geq 0. xy^iz \in L.$

# Pumping Lemma

## The Statement

### Lemma

*If  $L$  is regular then there is a number  $p$  (the pumping length) such that  $\forall w \in L$  with  $|w| \geq p$ ,  $\exists x, y, z \in \Sigma^*$  such that  $w = xyz$  and*

$$\textcircled{1} \quad |y| > 0$$

$$\exists p,$$

$$\forall w \in L \text{ with } |w| \geq p,$$

$$\exists x, y, z \in \Sigma^* \text{ such that } w = xyz, |y| > 0 \text{ and } |xy| \leq p,$$

$$\forall i \geq 0. xy^iz \in L.$$

# Pumping Lemma

## The Statement

### Lemma

*If  $L$  is regular then there is a number  $p$  (the pumping length) such that  $\forall w \in L$  with  $|w| \geq p$ ,  $\exists x, y, z \in \Sigma^*$  such that  $w = xyz$  and*

- ①  $|y| > 0$
- ②  $|xy| \leq p$

$\exists p,$

$\forall w \in L$  with  $|w| \geq p,$

$\exists x, y, z \in \Sigma^*$  such that  $w = xyz$ ,  $|y| > 0$  and  $|xy| \leq p,$

$\forall i \geq 0. xy^iz \in L.$

# Pumping Lemma

## The Statement

### Lemma

*If  $L$  is regular then there is a number  $p$  (the pumping length) such that  $\forall w \in L$  with  $|w| \geq p$ ,  $\exists x, y, z \in \Sigma^*$  such that  $w = xyz$  and*

- ①  $|y| > 0$
- ②  $|xy| \leq p$
- ③  $\forall i \geq 0. xy^iz \in L$

$\exists p,$

$\forall w \in L$  with  $|w| \geq p,$

$\exists x, y, z \in \Sigma^*$  such that  $w = xyz$ ,  $|y| > 0$  and  $|xy| \leq p,$

$\forall i \geq 0. xy^iz \in L.$

# Proving the Pumping Lemma

Proof.

Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA such that  $L(M) = L$  and let  $p = |Q|$ .

# Proving the Pumping Lemma

## Proof.

Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA such that  $L(M) = L$  and let  $p = |Q|$ . Let  $w = w_1 w_2 \cdots w_n \in L$  be such that  $n \geq p$ .

# Proving the Pumping Lemma

## Proof.

Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA such that  $L(M) = L$  and let  $p = |Q|$ . Let  $w = w_1 w_2 \cdots w_n \in L$  be such that  $n \geq p$ . For  $1 \leq i \leq n$ , let  $s_i = \hat{\delta}(q_0, w_1 \cdots w_i)$ ; define  $s_0 = q_0$ .



# Proving the Pumping Lemma

## Proof.

Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA such that  $L(M) = L$  and let  $p = |Q|$ . Let  $w = w_1 w_2 \cdots w_n \in L$  be such that  $n \geq p$ . For  $1 \leq i \leq n$ , let  $s_i = \hat{\delta}(q_0, w_1 \cdots w_i)$ ; define  $s_0 = q_0$ .

- Since  $s_0, s_1, \dots, s_i, \dots, s_p$  are  $p + 1$  states, there must be  $j, k$ ,  $0 \leq j < k \leq p$  such that  $s_j = s_k$  ( $= q$  say).

# Proving the Pumping Lemma

## Proof.

Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA such that  $L(M) = L$  and let  $p = |Q|$ . Let  $w = w_1 w_2 \cdots w_n \in L$  be such that  $n \geq p$ . For  $1 \leq i \leq n$ , let  $s_i = \hat{\delta}(q_0, w_1 \cdots w_i)$ ; define  $s_0 = q_0$ .

- Since  $s_0, s_1, \dots, s_i, \dots, s_p$  are  $p + 1$  states, there must be  $j, k$ ,  $0 \leq j < k \leq p$  such that  $s_j = s_k$  ( $= q$  say).
- Take  $x = w_1 \cdots w_j$ ,  $y = w_{j+1} \cdots w_k$ , and  $z = w_{k+1} \cdots w_n$

# Proving the Pumping Lemma

## Proof.

Let  $M = (Q, \Sigma, \delta, q_0, F)$  be a DFA such that  $L(M) = L$  and let  $p = |Q|$ . Let  $w = w_1 w_2 \cdots w_n \in L$  be such that  $n \geq p$ . For  $1 \leq i \leq n$ , let  $s_i = \hat{\delta}(q_0, w_1 \cdots w_i)$ ; define  $s_0 = q_0$ .

- Since  $s_0, s_1, \dots, s_i, \dots, s_p$  are  $p + 1$  states, there must be  $j, k$ ,  $0 \leq j < k \leq p$  such that  $s_j = s_k$  ( $= q$  say).
- Take  $x = w_1 \cdots w_j$ ,  $y = w_{j+1} \cdots w_k$ , and  $z = w_{k+1} \cdots w_n$
- Observe that since  $j < k \leq p$ , we have  $|xy| \leq p$  and  $|y| > 0$ .



# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

### Proof.

We will prove it by induction on  $i$ .

# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

### Proof.

We will prove it by induction on  $i$ .

- **Base Case:** By our assumption that  $s_j = s_k$  and the definition of  $x$  and  $y$ , we have  $\hat{\delta}(q_0, xy) = s_k = s_j = \hat{\delta}(q_0, x)$ .

# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

### Proof.

We will prove it by induction on  $i$ .

- **Base Case:** By our assumption that  $s_j = s_k$  and the definition of  $x$  and  $y$ , we have  $\hat{\delta}(q_0, xy) = s_k = s_j = \hat{\delta}(q_0, x)$ .
- **Induction Step:** We have





# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

### Proof.

We will prove it by induction on  $i$ .

- **Base Case:** By our assumption that  $s_j = s_k$  and the definition of  $x$  and  $y$ , we have  $\hat{\delta}(q_0, xy) = s_k = s_j = \hat{\delta}(q_0, x)$ .
- **Induction Step:** We have

$$\hat{\delta}(q_0, xy^{\ell+1}) = \hat{\delta}(\hat{\delta}(q_0, xy^{\ell}), y)$$



# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

### Proof.

We will prove it by induction on  $i$ .

- **Base Case:** By our assumption that  $s_j = s_k$  and the definition of  $x$  and  $y$ , we have  $\hat{\delta}(q_0, xy) = s_k = s_j = \hat{\delta}(q_0, x)$ .
- **Induction Step:** We have

$$\begin{aligned}\hat{\delta}(q_0, xy^{\ell+1}) &= \hat{\delta}(\hat{\delta}(q_0, xy^{\ell}), y) \\ &= \hat{\delta}(\hat{\delta}(q_0, x), y)\end{aligned}$$



# Proof ...

## Technical Claim

### Claim

For all  $i \geq 1$ ,  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$ .

### Proof.

We will prove it by induction on  $i$ .


- **Base Case:** By our assumption that  $s_j = s_k$  and the definition of  $x$  and  $y$ , we have  $\hat{\delta}(q_0, xy) = s_k = s_j = \hat{\delta}(q_0, x)$ .
- **Induction Step:** We have

$$\begin{aligned}\hat{\delta}(q_0, xy^{\ell+1}) &= \hat{\delta}(\hat{\delta}(q_0, xy^{\ell}), y) \\ &= \hat{\delta}(\hat{\delta}(q_0, x), y) \\ &= \hat{\delta}(q_0, xy) = \hat{\delta}(q_0, x)\end{aligned}$$




# Completing the Proof

Proof (contd).



# Completing the Proof


Proof (contd).



- We have  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$  for all  $i \geq 1$

# Completing the Proof


Proof (contd).



- We have  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$  for all  $i \geq 1$
- Since  $w \in L$ , we have  $\hat{\delta}(q_0, w) = \hat{\delta}(q_0, xyz) \in F$

# Completing the Proof


## Proof (contd).



- We have  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$  for all  $i \geq 1$
- Since  $w \in L$ , we have  $\hat{\delta}(q_0, w) = \hat{\delta}(q_0, xyz) \in F$
- Observe,  
 $\hat{\delta}(q_0, xz) = \hat{\delta}(\hat{\delta}(q_0, x), z) = \hat{\delta}(\hat{\delta}(q_0, xy), z) = \hat{\delta}(q_0, w)$ . So  
 $xz \in L$

# Completing the Proof

## Proof (contd).



- We have  $\hat{\delta}(q_0, xy^i) = \hat{\delta}(q_0, x)$  for all  $i \geq 1$
- Since  $w \in L$ , we have  $\hat{\delta}(q_0, w) = \hat{\delta}(q_0, xyz) \in F$
- Observe,  
 $\hat{\delta}(q_0, xz) = \hat{\delta}(\hat{\delta}(q_0, x), z) = \hat{\delta}(\hat{\delta}(q_0, xy), z) = \hat{\delta}(q_0, w)$ . So  $xz \in L$
- Similarly,  $\hat{\delta}(q_0, xy^i z) = \hat{\delta}(q_0, xyz) \in F$  and so  $xy^i z \in L$  □



# Finite Languages and Pumping Lemma

## Question

Do finite languages really satisfy the condition in the pumping lemma?

# Finite Languages and Pumping Lemma

## Question

Do finite languages really satisfy the condition in the pumping lemma?

**Recall Pumping Lemma:** If  $L$  is regular then there is a number  $p$  (the pumping length) such that  $\forall w \in L$  with  $|w| \geq p$ ,

$\exists x, y, z \in \Sigma^*$  such that  $w = xyz$  and

- 1  $|y| > 0$
- 2  $|xy| \leq p$
- 3  $\forall i \geq 0. xy^iz \in L$

# Finite Languages and Pumping Lemma

## Question

Do finite languages really satisfy the condition in the pumping lemma?

**Recall Pumping Lemma:** If  $L$  is regular then there is a number  $p$  (the pumping length) such that  $\forall w \in L$  with  $|w| \geq p$ ,  $\exists x, y, z \in \Sigma^*$  such that  $w = xyz$  and

- 1  $|y| > 0$
- 2  $|xy| \leq p$
- 3  $\forall i \geq 0. xy^iz \in L$

## Answer

Yes, they do. Let  $p$  be larger than the longest string in the language. Then the condition " $\forall w \in L$  with  $|w| \geq p, \dots$ " is **vaccuously** satisfied as there are no strings in the language longer than  $p$ !

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

## Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

## Pumping Condition

$$\begin{array}{l} \exists p. \quad \forall w \in L. \text{ with } |w| \geq p \quad \exists x, y, z \in \Sigma^*. w = xyz \\ \left. \begin{array}{l} (1) \quad |y| > 0 \\ (2) \quad |xy| \leq p \\ (3) \quad \forall i \geq 0. xy^i z \in L \end{array} \right\} \end{array}$$



# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

## Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

## Negation of the Pumping Condition

$$\neg p. \quad \forall w \in L. \text{ with } |w| \geq p \quad \exists x, y, z \in \Sigma^*. w = xyz$$
$$\left. \begin{array}{l} (1) \quad |y| > 0 \\ (2) \quad |xy| \leq p \\ (3) \quad \forall i \geq 0. xy^i z \in L \end{array} \right\}$$

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

## Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

## Negation of the Pumping Condition

$$\forall p. \quad \nexists w \in L. \text{ with } |w| \geq p \quad \exists x, y, z \in \Sigma^*. w = xyz$$
$$\left. \begin{array}{l} (1) \quad |y| > 0 \\ (2) \quad |xy| \leq p \\ (3) \quad \forall i \geq 0. xy^i z \in L \end{array} \right\}$$

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

## Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

## Negation of the Pumping Condition

$$\forall p. \quad \exists w \in L. \text{ with } |w| \geq p \quad \neg \exists x, y, z \in \Sigma^*. w = xyz$$
$$\left. \begin{array}{l} (1) \quad |y| > 0 \\ (2) \quad |xy| \leq p \\ (3) \quad \forall i \geq 0. xy^i z \in L \end{array} \right\}$$

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

## Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

## Negation of the Pumping Condition

$$\forall p. \quad \exists w \in L. \text{ with } |w| \geq p \quad \forall x, y, z \in \Sigma^*. w = xyz$$

$\left. \begin{array}{l} (1) \quad  y  > 0 \\ (2) \quad  xy  \leq p \\ (3) \quad \forall i \geq 0. xy^i z \in L \end{array} \right\}$	not all of them hold
---	----------------------

# Using the Pumping Lemma

$L$  regular implies that  $L$  satisfies the condition in the pumping lemma. If  $L$  regular  $\Rightarrow$  pumping Lemma. Then, Not pumping Lemma  $\Rightarrow L$  not regular.

## Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

## Negation of the Pumping Condition

$$\forall p. \quad \exists w \in L. \text{ with } |w| \geq p \quad \forall x, y, z \in \Sigma^*. w = xyz$$

$\left. \begin{array}{l} (1) \quad  y  > 0 \\ (2) \quad  xy  \leq p \\ (3) \quad \forall i \geq 0. xy^i z \in L \end{array} \right\}$	not all of them hold
---	----------------------

Equivalent to showing that if (1), (2) then (3) does not. In other words, we can find  $i$  such that  $xy^i z \notin L$

# Using the Pumping Lemma

## Pumping Lemma, in contrapositive

If  $L$  does not satisfy the pumping condition, then  $L$  not regular.

## Pumping Lemma, in contrapositive

$\forall p \geq 0,$

$\exists w \in L$  with  $|w| \geq p,$

$\forall x, y, z \in \Sigma^*$  such that  $w = xyz$ ,  $|y| > 0$  and  $|xy| \leq p,$

$\exists i \geq 0. xy^i z \notin L.$

Proving something like  $\forall \text{ bla } \exists \text{ bla } \forall \text{ bla } \exists \text{ bla}$  means winning a game.

Example: Two players: You, Adversary. Rules: First Adversary says a number. Then You say a number. You win if your number is bigger.

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

$L$       Task: To show that  $L$  is not regular



# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

$L$                       Task: To show that  $L$  is not regular  
 $\forall p.$                     Opponent picks  $p$

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

$L$	Task: To show that $L$ is not regular
$\forall p.$	<b>Opponent picks <math>p</math></b>
$\exists w.$	Pick $w$ that is of length at least $p$

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

- |                   |  |
|-------------------|--|
| $L$               | Task: To show that $L$ is not regular  |
| $\forall p.$      | Opponent picks $p$   |
| $\exists w.$      | Pick $w$ that is of length at least $p$  |
| $\forall x, y, z$ | Opponent divides $w$ into $x, y$ , and $z$ such that $ y  > 0$ , and $ xy  \leq p$ |

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

- $L$       Task: To show that  $L$  is not regular
- $\forall p.$       Opponent picks  $p$
- $\exists w.$       Pick  $w$  that is of length at least  $p$
- $\forall x, y, z$       Opponent divides  $w$  into  $x, y$ , and  $z$  such that  
                   $|y| > 0$ , and  $|xy| \leq p$
- $\exists k.$       You pick  $k$  and win if  $xy^kz \notin L$

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

- $L$  Task: To show that  $L$  is not regular
- $\forall p.$  Opponent picks  $p$
- $\exists w.$  Pick  $w$  that is of length at least  $p$
- $\forall x, y, z$  Opponent divides  $w$  into  $x, y$ , and  $z$  such that  $|y| > 0$ , and  $|xy| \leq p$
- $\exists k.$  You pick  $k$  and win if  $xy^kz \notin L$

**Pumping Lemma:** If  $L$  is regular, **opponent** has a winning strategy (no matter what you do).

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

- $L$       Task: To show that  $L$  is not regular
- $\forall p.$       **Opponent picks  $p$**
- $\exists w.$       **Pick  $w$  that is of length at least  $p$**
- $\forall x, y, z$       **Opponent divides  $w$  into  $x, y$ , and  $z$  such that  $|y| > 0$ , and  $|xy| \leq p$**
- $\exists k.$       **You pick  $k$  and win if  $xy^kz \notin L$**

**Pumping Lemma:** If  $L$  is regular, **opponent** has a winning strategy (no matter what you do).

**Contrapositive:** If **you** can beat the opponent,  $L$  not regular.

# Game View

Think of using the Pumping Lemma as a game between **you** and an **opponent**.

- $L$       Task: To show that  $L$  is not regular
- $\forall p.$       **Opponent picks  $p$**
- $\exists w.$       Pick  $w$  that is of length at least  $p$
- $\forall x, y, z$       **Opponent divides  $w$  into  $x, y$ , and  $z$  such that  $|y| > 0$ , and  $|xy| \leq p$**
- $\exists k.$       You pick  $k$  and win if  $xy^kz \notin L$

**Pumping Lemma:** If  $L$  is regular, **opponent** has a winning strategy (no matter what you do).

**Contrapositive:** If **you** can beat the opponent,  $L$  not regular.

Your strategy should work for any  $p$  and any subdivision that the opponent may come up with.

# Example I

## Proposition

$L_{0^n 1^n} = \{0^n 1^n \mid n \geq 0\}$  is not regular.



# Example I

## Proposition

$L_{0^n 1^n} = \{0^n 1^n \mid n \geq 0\}$  is not regular.

## Proof.

Suppose  $L_{0^n 1^n}$  is regular. Let  $p$  be the pumping length for  $L_{0^n 1^n}$ .

# Example I

## Proposition

$L_{0^n1^n} = \{0^n1^n \mid n \geq 0\}$  is not regular.

## Proof.

Suppose  $L_{0^n1^n}$  is regular. Let  $p$  be the pumping length for  $L_{0^n1^n}$ .

- Consider  $w = 0^p1^p$

# Example I

## Proposition

$L_{0^n1^n} = \{0^n1^n \mid n \geq 0\}$  is not regular.

## Proof.

Suppose  $L_{0^n1^n}$  is regular. Let  $p$  be the pumping length for  $L_{0^n1^n}$ .

- Consider  $w = 0^p1^p$
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^iz \in L_{0^n1^n}$ , for all  $i$ .

# Example I

## Proposition

$L_{0^n1^n} = \{0^n1^n \mid n \geq 0\}$  is not regular.

## Proof.

Suppose  $L_{0^n1^n}$  is regular. Let  $p$  be the pumping length for  $L_{0^n1^n}$ .

- Consider  $w = 0^p1^p$
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^iz \in L_{0^n1^n}$ , for all  $i$ .
- Since  $|xy| \leq p$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t1^p$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

# Example I

## Proposition

$L_{0^n1^n} = \{0^n1^n \mid n \geq 0\}$  is not regular.

## Proof.

Suppose  $L_{0^n1^n}$  is regular. Let  $p$  be the pumping length for  $L_{0^n1^n}$ .

- Consider  $w = 0^p1^p$
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^iz \in L_{0^n1^n}$ , for all  $i$ .
- Since  $|xy| \leq p$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t1^p$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

$$xy^0z = 0^r \epsilon 0^t1^p = 0^{r+t}1^p$$

# Example I

## Proposition

$L_{0^n1^n} = \{0^n1^n \mid n \geq 0\}$  is not regular.

## Proof.

Suppose  $L_{0^n1^n}$  is regular. Let  $p$  be the pumping length for  $L_{0^n1^n}$ .

- Consider  $w = 0^p1^p$
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^iz \in L_{0^n1^n}$ , for all  $i$ .
- Since  $|xy| \leq p$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t1^p$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

$$xy^0z = 0^r \epsilon 0^t1^p = 0^{r+t}1^p$$

Since  $r + t < p$ ,  $xy^0z \notin L_{0^n1^n}$ . Contradiction!



## Example II

### Proposition

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  is not regular.

## Example II

### Proposition

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  is not regular.

### Proof.

Suppose  $L_{\text{eq}}$  is regular. Let  $p$  be the pumping length for  $L_{\text{eq}}$ .



## Example II

### Proposition

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  is not regular.

### Proof.

Suppose  $L_{\text{eq}}$  is regular. Let  $p$  be the pumping length for  $L_{\text{eq}}$ .

- Consider  $w = 0^p 1^p$

## Example II

### Proposition

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  is not regular.

### Proof.

Suppose  $L_{\text{eq}}$  is regular. Let  $p$  be the pumping length for  $L_{\text{eq}}$ .

- Consider  $w = 0^p 1^p$
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_{\text{eq}}$ , for all  $i$ .

## Example II

### Proposition

$L_{\text{eq}} = \{w \in \{0,1\}^* \mid w \text{ has an equal number of 0s and 1s}\}$  is not regular.

### Proof.

Suppose  $L_{\text{eq}}$  is regular. Let  $p$  be the pumping length for  $L_{\text{eq}}$ .

- Consider  $w = 0^p 1^p$
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_{\text{eq}}$ , for all  $i$ .
- Since  $|xy| \leq p$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t 1^p$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

$$xy^0 z = 0^r 0^t 1^p = 0^{r+t} 1^p$$

Since  $r + t < p$ ,  $xy^0 z \notin L_{\text{eq}}$ . Contradiction!



# A Tale of two Proofs

## Non Pumping Lemma

Suppose  $L_{\text{eq}}$  is recognized by DFA  $M$  with  $p$  states. Consider the input  $0^p 1^p$ .

## Pumping Lemma

Suppose  $L_{\text{eq}}$  is regular. Let  $p$  be pumping length for  $L_{\text{eq}}$ . Consider  $w = 0^p 1^p$ .

# A Tale of two Proofs

## Non Pumping Lemma

Suppose  $L_{eq}$  is recognized by DFA  $M$  with  $p$  states. Consider the input  $0^p 1^p$ . There exist  $j, k$  and state  $q$  such that

## Pumping Lemma

Suppose  $L_{eq}$  is regular. Let  $p$  be pumping length for  $L_{eq}$ . Consider  $w = 0^p 1^p$ . There exist  $x, y, z$  such that

# A Tale of two Proofs

## Non Pumping Lemma

Suppose  $L_{eq}$  is recognized by DFA  $M$  with  $p$  states. Consider the input  $0^p 1^p$ . There exist  $j, k$  and state  $q$  such that

- $j < k$  and  
 $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$

## Pumping Lemma

Suppose  $L_{eq}$  is regular. Let  $p$  be pumping length for  $L_{eq}$ . Consider  $w = 0^p 1^p$ . There exist  $x, y, z$  such that

- $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ :  
so for some  $r, s, t$ ,  $x = 0^r$ ,  
 $y = 0^s$  and  $z = 0^t 1^p$ , with  
 $s > 0$ .

# A Tale of two Proofs

## Non Pumping Lemma

Suppose  $L_{eq}$  is recognized by DFA  $M$  with  $p$  states. Consider the input  $0^p 1^p$ . There exist  $j, k$  and state  $q$  such that

- $j < k$  and  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^p 1^p \in L_{eq}$ ,  $0^k 0^{(p-k)} 1^p$  is accepted by  $M$  and so is  $0^j 0^{(p-k)} 1^p$ .

## Pumping Lemma

Suppose  $L_{eq}$  is regular. Let  $p$  be pumping length for  $L_{eq}$ . Consider  $w = 0^p 1^p$ . There exist  $x, y, z$  such that

- $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ :  
so for some  $r, s, t$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t 1^p$ , with  $s > 0$ .
- $xy^i z \in L_{eq}$  for all  $i$ : so  $xy^0 z \in L_{eq}$ .

# A Tale of two Proofs

## Non Pumping Lemma

Suppose  $L_{\text{eq}}$  is recognized by DFA  $M$  with  $p$  states. Consider the input  $0^p 1^p$ . There exist  $j, k$  and state  $q$  such that

- $j < k$  and  $\hat{\delta}(q_0, 0^j) = \hat{\delta}(q_0, 0^k) = q$
- Since  $0^p 1^p \in L_{\text{eq}}$ ,  $0^k 0^{(p-k)} 1^p$  is accepted by  $M$  and so is  $0^j 0^{(p-k)} 1^p$ .
- But  $0^j 0^{(p-k)} 1^p \notin L_{\text{eq}}$ .

## Pumping Lemma

Suppose  $L_{\text{eq}}$  is regular. Let  $p$  be pumping length for  $L_{\text{eq}}$ . Consider  $w = 0^p 1^p$ . There exist  $x, y, z$  such that

- $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ :  
so for some  $r, s, t$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t 1^p$ , with  $s > 0$ .
- $xy^i z \in L_{\text{eq}}$  for all  $i$ : so  $xy^0 z \in L_{\text{eq}}$ .
- But  $xy^0 z = 0^{p-s} 1^p \notin L_{\text{eq}}$



# Example III

## Proposition

$L_p = \{0^i \mid i \text{ prime}\}$  is not regular

# Example III

## Proposition

$L_p = \{0^i \mid i \text{ prime}\}$  is not regular

## Proof.

Suppose  $L_p$  is regular. Let  $p$  be the pumping length for  $L_p$ .

# Example III

## Proposition

$L_p = \{0^i \mid i \text{ prime}\}$  is not regular

## Proof.

Suppose  $L_p$  is regular. Let  $p$  be the pumping length for  $L_p$ .

- Consider  $w = 0^m$ , where  $m \geq p + 2$  and  $m$  is prime.

## Example III

### Proposition

$L_p = \{0^i \mid i \text{ prime}\}$  is not regular

### Proof.

Suppose  $L_p$  is regular. Let  $p$  be the pumping length for  $L_p$ .

- Consider  $w = 0^m$ , where  $m \geq p + 2$  and  $m$  is prime.
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^iz \in L_p$ , for all  $i$ .

## Example III

### Proposition

$L_p = \{0^i \mid i \text{ prime}\}$  is not regular

### Proof.

Suppose  $L_p$  is regular. Let  $p$  be the pumping length for  $L_p$ .

- Consider  $w = 0^m$ , where  $m \geq p + 2$  and  $m$  is prime.
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_p$ , for all  $i$ .
- Thus,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

# Example III

## Proposition

$L_p = \{0^i \mid i \text{ prime}\}$  is not regular

## Proof.

Suppose  $L_p$  is regular. Let  $p$  be the pumping length for  $L_p$ .

- Consider  $w = 0^m$ , where  $m \geq p + 2$  and  $m$  is prime.
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_p$ , for all  $i$ .
- Thus,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t$ . Further, as  $|y| > 0$ , we have  $s > 0$ .  $xy^{r+t}z = 0^r(0^s)^{(r+t)}0^t = 0^{r+s(r+t)+t}$ .

# Example III

## Proposition

$L_p = \{0^i \mid i \text{ prime}\}$  is not regular

## Proof.

Suppose  $L_p$  is regular. Let  $p$  be the pumping length for  $L_p$ .

- Consider  $w = 0^m$ , where  $m \geq p + 2$  and  $m$  is prime.
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_p$ , for all  $i$ .
- Thus,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t$ . Further, as  $|y| > 0$ , we have  $s > 0$ .  $xy^{r+t}z = 0^r(0^s)^{(r+t)}0^t = 0^{r+s(r+t)+t}$ . Now  $r + s(r + t) + t = (r + t)(s + 1)$ . Further  $m = r + s + t \geq p + 2$  and  $s > 0$  mean that  $t \geq 2$  and  $s + 1 \geq 2$ . Thus,  $xy^{r+t}z \notin L_p$ . Contradiction!



# Example IV

## Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?



# Example IV

## Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?

Suppose  $L_{xx}$  is regular, and let  $p$  be the pumping length of  $L_{xx}$ .

# Example IV

## Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?

Suppose  $L_{xx}$  is regular, and let  $p$  be the pumping length of  $L_{xx}$ .

- Consider  $w = 0^p 0^p \in L$ .

## Example IV

### Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?

Suppose  $L_{xx}$  is regular, and let  $p$  be the pumping length of  $L_{xx}$ .

- Consider  $w = 0^p 0^p \in L$ .
- Can we find substrings  $x, y, z$  satisfying the conditions in the pumping lemma?

## Example IV

### Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?

Suppose  $L_{xx}$  is regular, and let  $p$  be the pumping length of  $L_{xx}$ .

- Consider  $w = 0^p 0^p \in L$ .
- Can we find substrings  $x, y, z$  satisfying the conditions in the pumping lemma? Yes! Consider  $x = \epsilon, y = 00, z = 0^{2p-2}$ .

## Example IV

### Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?

Suppose  $L_{xx}$  is regular, and let  $p$  be the pumping length of  $L_{xx}$ .

- Consider  $w = 0^p 0^p \in L$ .
- Can we find substrings  $x, y, z$  satisfying the conditions in the pumping lemma? Yes! Consider  $x = \epsilon, y = 00, z = 0^{2p-2}$ .
- Does this mean  $L_{xx}$  satisfies the pumping lemma? Does it mean it is regular?

## Example IV

### Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?

Suppose  $L_{xx}$  is regular, and let  $p$  be the pumping length of  $L_{xx}$ .

- Consider  $w = 0^p 0^p \in L$ .
- Can we find substrings  $x, y, z$  satisfying the conditions in the pumping lemma? Yes! Consider  $x = \epsilon, y = 00, z = 0^{2p-2}$ .
- Does this mean  $L_{xx}$  satisfies the pumping lemma? Does it mean it is regular?
  - No! We have chosen a bad  $w$ . To prove that the pumping lemma is violated, we only need to exhibit **some**  $w$  that cannot be pumped.

## Example IV

### Question

Is  $L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  regular?

Suppose  $L_{xx}$  is regular, and let  $p$  be the pumping length of  $L_{xx}$ .

- Consider  $w = 0^p 0^p \in L$ .
- Can we find substrings  $x, y, z$  satisfying the conditions in the pumping lemma? Yes! Consider  $x = \epsilon, y = 00, z = 0^{2p-2}$ .
- Does this mean  $L_{xx}$  satisfies the pumping lemma? Does it mean it is regular?
  - No! We have chosen a bad  $w$ . To prove that the pumping lemma is violated, we only need to exhibit **some**  $w$  that cannot be pumped.
- Another bad choice  $(01)^p(01)^p$ .

# Example IV

Reloaded

## Proposition

$L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  *is not regular.*



# Example IV

Reloaded

## Proposition

$L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  *is not regular.*

## Proof.

Suppose  $L_{xx}$  is regular. Let  $p$  be the pumping length for  $L_{xx}$ .

# Example IV

Reloaded

## Proposition

$L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  is not regular.

## Proof.

Suppose  $L_{xx}$  is regular. Let  $p$  be the pumping length for  $L_{xx}$ .

- Consider  $w = 0^p 10^p 1$ .

# Example IV

Reloaded

## Proposition

$L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  is not regular.

## Proof.

Suppose  $L_{xx}$  is regular. Let  $p$  be the pumping length for  $L_{xx}$ .

- Consider  $w = 0^p 1 0^p 1$ .
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_p$ , for all  $i$ .

# Example IV

Reloaded

## Proposition

$L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  is not regular.

## Proof.

Suppose  $L_{xx}$  is regular. Let  $p$  be the pumping length for  $L_{xx}$ .

- Consider  $w = 0^p 10^p 1$ .
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_p$ , for all  $i$ .
- Since  $|xy| \leq p$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t 10^p 1$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

# Example IV

Reloaded

## Proposition

$L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  is not regular.

## Proof.

Suppose  $L_{xx}$  is regular. Let  $p$  be the pumping length for  $L_{xx}$ .

- Consider  $w = 0^p 1 0^p 1$ .
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_p$ , for all  $i$ .
- Since  $|xy| \leq p$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t 1 0^p 1$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

$$xy^0 z = 0^r \epsilon 0^t 1 0^p 1 = 0^{r+t} 1 0^p 1$$

# Example IV

Reloaded

## Proposition

$L_{xx} = \{xx \mid x \in \{0,1\}^*\}$  is not regular.

## Proof.

Suppose  $L_{xx}$  is regular. Let  $p$  be the pumping length for  $L_{xx}$ .

- Consider  $w = 0^p 1 0^p 1$ .
- Since  $|w| > p$ , there are  $x, y, z$  such that  $w = xyz$ ,  $|xy| \leq p$ ,  $|y| > 0$ , and  $xy^i z \in L_p$ , for all  $i$ .
- Since  $|xy| \leq p$ ,  $x = 0^r$ ,  $y = 0^s$  and  $z = 0^t 1 0^p 1$ . Further, as  $|y| > 0$ , we have  $s > 0$ .

$$xy^0 z = 0^r \epsilon 0^t 1 0^p 1 = 0^{r+t} 1 0^p 1$$

Since  $r + t < p$ ,  $xy^0 z \notin L_{xx}$ . Contradiction!



# Lessons on Expressivity

## Limits of Finite Memory

Finite automata cannot

- “keep track of counts”: e.g.,  $L_{0n1n}$  not regular.
- “compare far apart pieces” of the input: e.g.  $L_{xx}$  not regular.
- do “computations that require it to look at global properties” of the input. e.g.  $L_{\text{prime}}$  not regular.

# Lessons on Expressivity

## Limits of Finite Memory

Finite automata cannot

- “keep track of counts”: e.g.,  $L_{0n1n}$  not regular.
- “compare far apart pieces” of the input: e.g.  $L_{xx}$  not regular.
- do “computations that require it to look at global properties” of the input. e.g.  $L_{\text{prime}}$  not regular.

... and pumping lemma provides **one way** to find out some of these limitations.