

CS 611: Theory of Computation

Hongmin Li

Department of Computer Science
California State University, East Bay

Emptiness Problem

Given a CFG G with start symbol S , is $L(G)$ empty?

Emptiness Problem

Given a CFG G with start symbol S , is $L(G)$ empty?

Solution: Check if the start symbol S is generating.

Emptiness Problem

Given a CFG G with start symbol S , is $L(G)$ empty?

Solution: Check if the start symbol S is generating. How long does that take?

Determining generating symbols

Algorithm

```
Gen = {}  
for every rule  $A \rightarrow x$  where  $x \in \Sigma^*$   
    Gen = Gen  $\cup$  {A}  
repeat  
    for every rule  $A \rightarrow \gamma$   
        if all variables in  $\gamma$  are generating then  
            Gen = Gen  $\cup$  {A}  
until Gen does not change
```

Determining generating symbols

Algorithm

```
Gen = {}  
for every rule  $A \rightarrow x$  where  $x \in \Sigma^*$   
    Gen = Gen  $\cup$  {A}  
repeat  
    for every rule  $A \rightarrow \gamma$   
        if all variables in  $\gamma$  are generating then  
            Gen = Gen  $\cup$  {A}  
until Gen does not change
```

- Both for-loops take $O(n)$ time where $n = |G|$.

Determining generating symbols

Algorithm

```
Gen = {}  
for every rule  $A \rightarrow x$  where  $x \in \Sigma^*$   
    Gen = Gen  $\cup$  {A}  
repeat  
    for every rule  $A \rightarrow \gamma$   
        if all variables in  $\gamma$  are generating then  
            Gen = Gen  $\cup$  {A}  
until Gen does not change
```

- Both for-loops take $O(n)$ time where $n = |G|$.
- Each iteration of repeat-until loop discovers a new variable. So number of iterations is $O(n)$. And total is $O(n^2)$.

Membership Problem

Given a CFG $G = (V, \Sigma, R, S)$ in **Chomsky Normal Form**, and a string $w \in \Sigma^*$, is $w \in L(G)$?

Membership Problem

Given a CFG $G = (V, \Sigma, R, S)$ in **Chomsky Normal Form**, and a string $w \in \Sigma^*$, is $w \in L(G)$?

Central question in parsing.

“Simple” Solution

“Simple” Solution

- Let $|w| = n$. Since G is in Chomsky Normal Form, w has a parse tree of size $2n - 1$ iff $w \in L(G)$

“Simple” Solution

- Let $|w| = n$. Since G is in Chomsky Normal Form, w has a parse tree of size $2n - 1$ iff $w \in L(G)$
- Construct all possible parse (binary) trees and check if any of them is a valid parse tree for w

“Simple” Solution

- Let $|w| = n$. Since G is in Chomsky Normal Form, w has a parse tree of size $2n - 1$ iff $w \in L(G)$
- Construct all possible parse (binary) trees and check if any of them is a valid parse tree for w
- Number of parse trees of size $2n - 1$ is k^{2n-1} where k is the number of variables in G . So algorithm is exponential in n !

“Simple” Solution

- Let $|w| = n$. Since G is in Chomsky Normal Form, w has a parse tree of size $2n - 1$ iff $w \in L(G)$
- Construct all possible parse (binary) trees and check if any of them is a valid parse tree for w
- Number of parse trees of size $2n - 1$ is k^{2n-1} where k is the number of variables in G . So algorithm is exponential in n !
- We will see an algorithm that runs in $O(n^3)$ time (the constant will depend on k).

First Ideas

Notation

Suppose $w = w_1 w_2 \cdots w_n$, where $w_i \in \Sigma$. Let $w_{i,j}$ denote the substring of w starting at position i of length j . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

First Ideas

Notation

Suppose $w = w_1 w_2 \cdots w_n$, where $w_i \in \Sigma$. Let $w_{i,j}$ denote the substring of w starting at position i of length j . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

Main Idea

For every $A \in V$, and every $i \leq n$, $j \leq n + 1 - i$, we will determine if $A \xRightarrow{*} w_{i,j}$.

First Ideas

Notation

Suppose $w = w_1 w_2 \cdots w_n$, where $w_i \in \Sigma$. Let $w_{i,j}$ denote the substring of w starting at position i of length j . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

Main Idea

For every $A \in V$, and every $i \leq n$, $j \leq n + 1 - i$, we will determine if $A \xRightarrow{*} w_{i,j}$.

Now, $w \in L(G)$ iff $S \xRightarrow{*} w_{1,n} = w$; thus, we will solve the membership problem.

First Ideas

Notation

Suppose $w = w_1 w_2 \cdots w_n$, where $w_i \in \Sigma$. Let $w_{i,j}$ denote the substring of w starting at position i of length j . Thus,

$$w_{i,j} = w_i w_{i+1} \cdots w_{i+j-1}$$

Main Idea

For every $A \in V$, and every $i \leq n$, $j \leq n + 1 - i$, we will determine if $A \xRightarrow{*} w_{i,j}$.

Now, $w \in L(G)$ iff $S \xRightarrow{*} w_{1,n} = w$; thus, we will solve the membership problem.

How do we determine if $A \xRightarrow{*} w_{i,j}$ for every A, i, j ?

Base Case

Substrings of length 1

Observation

For any A, i , $A \xRightarrow{*} w_{i,1}$ iff $A \rightarrow w_{i,1}$ is a rule.

Base Case

Substrings of length 1

Observation

For any A, i , $A \xRightarrow{*} w_{i,1}$ iff $A \rightarrow w_{i,1}$ is a rule.

- Since G is in Chomsky Normal Form, G does not have any ϵ -rules, nor any unit rules.

Base Case

Substrings of length 1

Observation

For any A, i , $A \xRightarrow{*} w_{i,1}$ iff $A \rightarrow w_{i,1}$ is a rule.

- Since G is in Chomsky Normal Form, G does not have any ϵ -rules, nor any unit rules.

Thus, for each A and i , one can determine if $A \xRightarrow{*} w_{i,1}$.


Inductive Step

Longer substrings

Suppose for every variable X and every $w_{i,\ell}$ ($\ell < j$) we have determined if $X \xRightarrow{*} w_{i,\ell}$

Inductive Step

Longer substrings




Suppose for every variable X and every $w_{i,\ell}$ ($\ell < j$) we have determined if $X \xRightarrow{*} w_{i,\ell}$

- $A \xRightarrow{*} w_{i,j}$ iff there are variables B and C and some $k < j$ such that $A \rightarrow BC$ is a rule, and $B \xRightarrow{*} w_{i,k}$ and $C \xRightarrow{*} w_{i+k,j-k}$

Inductive Step

Longer substrings



Suppose for every variable X and every $w_{i,\ell}$ ($\ell < j$) we have determined if $X \xRightarrow{*} w_{i,\ell}$

- $A \xRightarrow{*} w_{i,j}$ iff there are variables B and C and some $k < j$ such that $A \rightarrow BC$ is a rule, and $B \xRightarrow{*} w_{i,k}$ and $C \xRightarrow{*} w_{i+k,j-k}$
- Since k and $j - k$ are both less than j , we can inductively determine if $A \xRightarrow{*} w_{i,j}$.

Cocke-Younger-Kasami (CYK) Algorithm

Algorithm maintains $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$.

Initialize: $X_{i,1} = \{A \mid A \rightarrow w_{i,1}\}$

for $j = 2$ **to** n **do**

for $i = 1$ **to** $n - j + 1$ **do**

$X_{i,j} = \emptyset$

for $k = 1$ **to** $j - 1$ **do**

$X_{i,j} = X_{i,j} \cup \{A \mid A \rightarrow BC, B \in X_{i,k}, C \in X_{i+k,j-k}\}$

Cocke-Younger-Kasami (CYK) Algorithm

Algorithm maintains $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$.

Initialize: $X_{i,1} = \{A \mid A \rightarrow w_{i,1}\}$

for $j = 2$ to n **do**

for $i = 1$ to $n - j + 1$ **do**

$X_{i,j} = \emptyset$

for $k = 1$ to $j - 1$ **do**

$X_{i,j} = X_{i,j} \cup \{A \mid A \rightarrow BC, B \in X_{i,k}, C \in X_{i+k,j-k}\}$

Correctness: After each iteration of the outermost loop, $X_{i,j}$ contains exactly the set of variables A that can derive $w_{i,j}$, for each i .

Cocke-Younger-Kasami (CYK) Algorithm

Algorithm maintains $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$.

Initialize: $X_{i,1} = \{A \mid A \rightarrow w_{i,1}\}$

for $j = 2$ to n **do**

for $i = 1$ to $n - j + 1$ **do**

$X_{i,j} = \emptyset$

for $k = 1$ to $j - 1$ **do**

$X_{i,j} = X_{i,j} \cup \{A \mid A \rightarrow BC, B \in X_{i,k}, C \in X_{i+k,j-k}\}$

Correctness: After each iteration of the outermost loop, $X_{i,j}$ contains exactly the set of variables A that can derive $w_{i,j}$, for each i . Time = $O(n^3)$.

Example

Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$ Let

$w = baaba$. The sets $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$:

Example

Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$ Let

$w = baaba$. The sets $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$:

j/i	1	2	3	4	5
5					
4					
3					
2					
1	$\{B\}$	$\{A, C\}$	$\{A, C\}$	$\{B\}$	$\{A, C\}$
	b	a	a	b	a

Example

Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$ Let

$w = baaba$. The sets $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$:

j/i	1	2	3	4	5
5					
4					
3					
2	$\{S, A\}$	$\{B\}$	$\{S, C\}$	$\{S, A\}$	
1	$\{B\}$	$\{A, C\}$	$\{A, C\}$	$\{B\}$	$\{A, C\}$
	b	a	a	b	a

Example

Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$ Let

$w = baaba$. The sets $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$:

j/i	1	2	3	4	5
5					
4					
3	\emptyset	$\{B\}$	$\{B\}$		
2	$\{S, A\}$	$\{B\}$	$\{S, C\}$	$\{S, A\}$	
1	$\{B\}$	$\{A, C\}$	$\{A, C\}$	$\{B\}$	$\{A, C\}$
	b	a	a	b	a

Example

Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$ Let

$w = baaba$. The sets $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$:

j/i	1	2	3	4	5
5					
4	\emptyset	$\{S, A, C\}$			
3	\emptyset	$\{B\}$	$\{B\}$		
2	$\{S, A\}$	$\{B\}$	$\{S, C\}$	$\{S, A\}$	
1	$\{B\}$	$\{A, C\}$	$\{A, C\}$	$\{B\}$	$\{A, C\}$
	b	a	a	b	a

Example

Example

Consider grammar

$S \rightarrow AB \mid BC, A \rightarrow BA \mid a, B \rightarrow CC \mid b, C \rightarrow AB \mid a$ Let

$w = baaba$. The sets $X_{i,j} = \{A \mid A \xRightarrow{*} w_{i,j}\}$:

j/i	1	2	3	4	5
5	{S, A, C}				
4	\emptyset	{S, A, C}			
3	\emptyset	{B}	{B}		
2	{S, A}	{B}	{S, C}	{S, A}	
1	{B}	{A, C}	{A, C}	{B}	{A, C}
	b	a	a	b	a

More Decision Problems

Given a CFGs G_1 and G_2

More Decision Problems

Given a CFGs G_1 and G_2

- Is $L(G_1) = \Sigma^*$?

More Decision Problems

Given a CFGs G_1 and G_2

- Is $L(G_1) = \Sigma^*$?
- Is $L(G_1) \cap L(G_2) = \emptyset$?

More Decision Problems

Given a CFGs G_1 and G_2

- Is $L(G_1) = \Sigma^*$?
- Is $L(G_1) \cap L(G_2) = \emptyset$?
- Is $L(G_1) = L(G_2)$?

More Decision Problems

Given a CFGs G_1 and G_2

- Is $L(G_1) = \Sigma^*$?
- Is $L(G_1) \cap L(G_2) = \emptyset$?
- Is $L(G_1) = L(G_2)$?
- Is G_1 ambiguous?

More Decision Problems

Given a CFGs G_1 and G_2

- Is $L(G_1) = \Sigma^*$?
- Is $L(G_1) \cap L(G_2) = \emptyset$?
- Is $L(G_1) = L(G_2)$?
- Is G_1 ambiguous?
- Is $L(G_1)$ inherently ambiguous?

More Decision Problems

Given a CFGs G_1 and G_2

- Is $L(G_1) = \Sigma^*$?
- Is $L(G_1) \cap L(G_2) = \emptyset$?
- Is $L(G_1) = L(G_2)$?
- Is G_1 ambiguous?
- Is $L(G_1)$ inherently ambiguous?

All these problems are undecidable.