VECTORS

12^{th} Math - Chapter 10

This is Problem-16 from Exercise 10.5

- 1. If θ is the angle between two vectors $\overrightarrow{\mathbf{a}}$ and $\overrightarrow{\mathbf{b}}$, then $\overrightarrow{\mathbf{a}} \cdot \overrightarrow{\mathbf{b}} \geq 0$.
 - (a) $0 < \theta < \frac{\pi}{2}$
 - (b) $0 \le \theta \le \frac{\pi}{2}$
 - (c) $0 < \theta < \pi$
 - (d) $0 \le \theta \le \pi$

Solution: Given

$$\mathbf{a}, \mathbf{b}$$
 are two vectors (1)

$$\mathbf{a}^{\top}\mathbf{b} \ge 0 \tag{2}$$

Assume that **a**,**b** are,

$$\mathbf{a} = \begin{pmatrix} 4\\3 \end{pmatrix} \tag{3}$$

$$\mathbf{b} = \begin{pmatrix} 5\\12 \end{pmatrix} \tag{4}$$

We know that

$$\theta = \cos^{-1} \left(\frac{\mathbf{a}^{\mathsf{T}} \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|} \right) \tag{5}$$

$$\implies \mathbf{a}^{\mathsf{T}}\mathbf{b} = \cos\theta \|\mathbf{a}\| \|\mathbf{b}\| \tag{6}$$

$$\|\mathbf{a}\| = \sqrt{\mathbf{a}_1^2 + \mathbf{a}_2^2} \tag{7}$$

Verification:

$$\|\mathbf{a}\| = \sqrt{4^2 + 3^2}$$

$$\Rightarrow = 5$$

$$\|\mathbf{b}\| = \sqrt{5^2 + 12^2}$$

$$\Rightarrow = 13$$

$$\text{for } \theta = 0$$

$$\mathbf{a}^{\mathsf{T}} \mathbf{b} = \cos(0)(5)13)$$

$$\Rightarrow = 65$$

$$\text{for } \theta = \frac{\pi}{2}$$

$$\mathbf{a}^{\mathsf{T}} \mathbf{b} \ge 0$$

$$\text{for } \theta = \frac{\pi}{2}$$

$$\mathbf{a}^{\mathsf{T}} \mathbf{b} = \cos(\frac{\pi}{2})(5)(13)$$

$$\Rightarrow = 0$$

$$\text{for } \theta = \pi$$

$$\mathbf{a}^{\mathsf{T}} \mathbf{b} = \cos(\pi)(5)(13)$$

$$\Rightarrow = -65$$

$$\Rightarrow \mathbf{a}^{\mathsf{T}} \mathbf{b} < 0$$

$$(16)$$

$$\Rightarrow \mathbf{a}^{\mathsf{T}} \mathbf{b} < 0$$

Therefore, the θ is $0 \leq \theta \leq \frac{\pi}{2}$. So, option (b) is correct.