

# Quadrature

JUDD MEHR

## Introduction

In general, quadrature is a process by which the calculus problem of determining the integral of a curve is cast as a linear algebra problem whereby a weighted sum of intelligently chosen samples approximates the true value of the integral.

$$\int_a^b f(x)dx \approx \sum_{i=1}^n w_i f(x_i)$$

Undergraduate calculus and numerical methods courses often cover rectangle, trapezoid, and Simpson's quadrature schemes, and will therefore not be covered here. We will discuss the basics of Gauss-Legendre Quadrature first, a powerful method allowing the exact approximation of polynomial functions.

## Basic Theory of Gauss-Legendre Quadrature

In Gauss-Legendre quadrature, we obtain the abscissae at which we sample the curve in question using points derived from the Legendre polynomials, and weights chosen wisely to get the best possible approximation given a certain number of degrees of freedom.

### Legendre Polynomials

Legendre polynomials are a set of polynomials forming a basis of the polynomial space up to degree  $n$  where  $n$  is the order of the highest order poly-

nomial in the set. To obtain the Legendre polynomials we start with the orthogonal set

$$\{1, x, x^2, \dots, x^n\}$$

and through the Gram-Schmidt orthogonalization process, with respect to the inner product

$$\langle p, q \rangle = \int_{-1}^1 p(x)q(x)dx$$

we obtain a new set of orthogonal polynomials that spans the same function space. We then apply constant scalings to the polynomials so that  $P_n(1) = 1$

These polynomials (scaling included) can also be calculated using Rodrigues' formula

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$$

Legendre polynomials have several attractive properties. Here I mention just a few that will be shown to be important momentarily. First, Legendre polynomials of degree  $n$  have exactly  $n$  real roots. Second, all the roots of Legendre polynomials fall in the range  $(-1, 1)$ .

### Gaussian Weights

Given a polynomial  $p(x)$  of degree  $2n - 1$ , we express  $p$  in terms of a polynomial division .

$$p(x) = q(x)L_n(x) + r(x)$$

where  $L_n(x)$  is the  $n$ th degree Legendre polynomial, and  $q(x)$  is some polynomial of degree  $\leq n - 1$ , and  $r(x)$  is the remainder of the division of  $q$  and  $L$  and is also of degree  $\leq n - 1$ .

Integrating, we have

$$\int_{-1}^1 p(x)dx = \int_{-1}^1 q(x)L_n(x)dx + \int_{-1}^1 r(x)dx$$

First note that the definition of our inner product mentioned earlier has appeared, because  $q$  and  $L$  are, by definition, orthonganal, this first term on the right side equals zero. We can therefore approximate this part of the integral *exactly* by choosing the abscissae at which to sample the curve to be the roots of the Legendre polynomial of degree  $n$ , which are all real and

conveniently fall between the bounds of intergration. Doing this will cause the approximation of that term to equate to zero as is its true value.

Next, we can also approximate the second term on the right side *exactly* by intelligently choosing weights. With  $n$  degrees of freedom, we can approximate exactly any polynomial of order  $\leq n - 1$  which is the order of  $r$ . We do this by applying the Vandermonde matrix in this manner

$$\begin{bmatrix} 1 & 1 & \dots & 1 \\ x_1 & x_2 & \dots & x_n \\ x_1^2 & x_2^2 & \dots & x_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_1^{n-1} & x_2^{n-1} & \dots & x_n^{n-1} \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ \vdots \\ w_n \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ \frac{2}{3} \\ \vdots \\ \int_{-1}^1 x^{n-1} \end{bmatrix}$$

where  $x_n$  are the  $n$ th roots of the  $n$ th degree Legendre polynomial,  $w_n$  is the  $n$ th weighting value, and the right hand side are the true values of the integrals from with bounds  $[-1, 1]$  from degree zero up to  $n - 1$ .

We can also calculate the weights using the following formula

$$w_i = \frac{2(1 - x_i^2)}{(n + 1)^2 [P_{n+1}(x_i)]^2}$$

where again  $x_i$  are the roots of the  $n$ th degree Legendre polynomial.

Thus with the intelligent choice of both abscissae and weights, we can approximate *exactly* any polynomial of degree  $2n - 1$  with only  $n$  points. This is the power of Gaussian quadrature.

## Transforming the Bounds of Integration

Above we have shown that the integral of any polynomial of degree  $2n - 1$  with bounds  $[-1, 1]$  can be approximated exactly using  $n$  points. With a simple transformation, we can evaluate the integral with arbitrary bounds  $[a, b]$  in the following manner

$$\int_a^b f(x)dx \approx \frac{b-a}{2} \sum_{i=1}^n w_n f(\xi_n)$$

where

$$\xi_n = \frac{b-a}{2} x_n + \frac{a+b}{2}$$

## Code Implementation of Gauss-Legendre Quadrature

After understanding the basic theory, the implementation becomes relatively simple. In the Julia language, there is a FastGaussQuadrature package that efficiently solves for the abscissae and weights for an  $n$ th degree approximation, making any efforts of mine to do so redundant. To find the polynomial roots, the package uses analytic expressions up to degree 5, Newton's method up to degree 60, and asymptotic expansion for degree greater than 60. Using that package, an implementation of this quadrature scheme is as follows.

```
"""
gaussquadrature(fun, n; lb=-1, ub=1)

Evaluate the numeric integral of the function, fun, from the
lowerbound, lb, to the upper bound, ub, using an n-th
degree Gauss-Legendre quadrature rule.
"""
function gaussquadrature(fun, n; lb=-1, ub=1)
#compute abscissae and weights using FastGaussQuadrature
t, w = FastGaussQuadrature.gausslegendre(n)

#transform bounds
a = (ub-lb).*t./2.0 .+ (lb+ub)/2.0
d = (ub-lb)/2

#initialize integral
integral = 0.0

#calculate numeric integral:
#Evaluate the function (or spline object) at the abscissae and
multiply by the weights; sum the values together to obtain
an estimate of the integral.
for i=1:n
integral += w[i]*d*fun(a[i])
end

return integral
end
```

## Application to Strongly Singular Integrals

In Boundary Element Methods (BEM), we often find ourselves evaluating singular integrals. Presented here is one method for doing so, sometimes

referred to as the Subtraction of Singularity Method (SST) as described by Guiggiani and Casalini.[\[1\]](#)

# Bibliography

- [1] Guiggiani, M. and Casalini, P., “Direct computation of Cauchy principal value integrals in advanced boundary elements,” *International Journal for Numerical Methods in Engineering*, Vol. 24, No. 9, Sep 1987, pp. 1711–1720.