

Leave
blank

5. [In this question \mathbf{i} and \mathbf{j} are perpendicular horizontal unit vectors.]

A particle P is moving with constant acceleration. At 2pm, the velocity of P is $(3\mathbf{i} + 5\mathbf{j}) \text{ km h}^{-1}$ and at 2.30pm the velocity of P is $(\mathbf{i} + 7\mathbf{j}) \text{ km h}^{-1}$

At time T hours after 2pm, P is moving in the direction of the vector $(-\mathbf{i} + 2\mathbf{j})$

- (a) Find the value of T .

(6)

Another particle, Q , has velocity $\mathbf{v}_Q \text{ km h}^{-1}$ at time t hours after 2pm, where

$$\mathbf{v}_Q = (-4 - 2t)\mathbf{i} + (\mu + 3t)\mathbf{j}$$

and μ is a constant.

Given that there is an instant when the velocity of P is equal to the velocity of Q ,

- (b) find the value of μ .

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank**Question 5 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 6 A 0 1 7 3 2

Leave
blank**Question 5 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 5 continued

Leave
blank

Q5

(Total 9 marks)



P 6 5 2 6 6 A 0 1 9 3 2