

Question Number	Scheme	Marks
2(a)	$M(D), \frac{3a}{5} Xg = \frac{2a}{5} Mg$ <p>Other possible equations:</p> $(\uparrow)T_D = Mg + Xg$ $M(A), Mga + Xg 2a = T_D \frac{7a}{5}$ $M(B), Mga = T_D \frac{3a}{5} \quad T_D \text{ would then need to be eliminated}$ $M(C), Mg \frac{3a}{5} + Xg \frac{8a}{5} = T_D a$ $M(G), Xga = T_D \frac{2a}{5}$	M1A1
	$X = \frac{2M}{3}, 0.67 M \text{ or better}$	A1 (3)
2(b)	$M(D), T_C a + \frac{1}{2} Mg \frac{3a}{5} = \frac{2a}{5} Mg$ <p>Other possible equations:</p> $(\uparrow)T_C + T_D = Mg + \frac{1}{2} Mg$ $M(A), Mga + \frac{1}{2} Mg 2a = T_C \frac{2a}{5} + T_D \frac{7a}{5}$ $M(B), Mga = T_C \frac{8a}{5} + T_D \frac{3a}{5} \quad T_D \text{ would need eliminating}$ $M(C), Mg \frac{3a}{5} + \frac{1}{2} Mg \frac{8a}{5} = T_D a$ $M(G), T_C \frac{3a}{5} + \frac{1}{2} Mga = T_D \frac{2a}{5}$	M1A1
	$T_C = \frac{1}{10} Mg \text{ oe}$	A1 (3)
		(6)
	Notes for question 2	
2(a)	<p>M1 For an equation (or inequality,,) in X, M and a only (allow consistent missing a's) with correct no. of terms. Allow if one g is missing. N.B. M0 if T_C appears and never becomes zero</p>	
	A1 Correct equation or inequality	
	A1 cao	
2(b)	<p>M1 For an equation in T_C, M, g and a only (allow consistent missing a's or if g's) missing) with correct no. of terms M0 if they assume that $T_C = T_D$ or if they assume their X value from (a).</p>	
	A1 Correct equation	
	A1 cao	