Please check the examination details belo	w before entering your candidate information
Candidate surname	Other names
Pearson Edexcel Intern	
Friday 26 May 2023	
Afternoon (Time: 2 hours)	Paper reference 4PM1/01
Further Pure Math PAPER 1	nematics
Calculators may be used.	Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- You must NOT write anything on the formulae page.
 Anything you write on the formulae page will gain NO credit.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶





International GCSE in Further Pure Mathematics Formulae sheet

Mensuration

Surface area of sphere = $4\pi r^2$

Curved surface area of cone = $\pi r \times \text{slant height}$

Volume of sphere =
$$\frac{4}{3}\pi r^3$$

Series

Arithmetic series

Sum to *n* terms,
$$S_n = \frac{n}{2} [2a + (n-1)d]$$

Geometric series

Sum to *n* terms,
$$S_n = \frac{a(1-r^n)}{(1-r)}$$

Sum to infinity,
$$S_{\infty} = \frac{a}{1-r} |r| < 1$$

Binomial series

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots$$
 for $|x| < 1, n \in \mathbb{Q}$

Calculus

Quotient rule (differentiation)

$$\frac{\mathrm{d}}{\mathrm{d}x} \left(\frac{\mathrm{f}(x)}{\mathrm{g}(x)} \right) = \frac{\mathrm{f}'(x)\mathrm{g}(x) - \mathrm{f}(x)\mathrm{g}'(x)}{\left[\mathrm{g}(x)\right]^2}$$

Trigonometry

Cosine rule

In triangle ABC: $a^2 = b^2 + c^2 - 2bc \cos A$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

Logarithms

$$\log_a x = \frac{\log_b x}{\log_b a}$$



Answer all ELEVEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

(a) Show that $\sum_{r=1}^{n} (3r+2) = \frac{n}{2} (3n+7)$

(3)

(b) Hence, or otherwise, evaluate $\sum_{r=10}^{40} (3r+2)$

(2)

(Total for Question 1 is 5 marks)

2			$y = (\sin 2x)\sqrt{3 + 2x}$		
	Show that	$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\sin 2x + (A + Bx)c}{\sqrt{3 + 2x}}$	$\frac{\cos 2x}{\text{where } A \text{ and } B \text{ are}}$	e integers to be found.	(5)



Question 2 continued
(Total for Question 2 is 5 marks)



(5)

3




Figure 1

Figure 1 shows part of the curve with equation $y = \frac{x}{2} + \frac{4}{x^2}$ in the interval 0.8 < x < 7

By drawing a suitable straight line on the grid, obtain an estimate, to one decimal place, of the roots of the equation $3x^3 - 12x^2 + 8 = 0$ in the interval 0.8 < x < 7

P 7	2 8 6	4 A 0	7 3 2
' '	2 0 0	4 / 0	1 3 2

4	A particle <i>P</i> is moving along the <i>x</i> -axis. At time <i>t</i> seconds, $t \ge 0$, the velocity, $v \text{ m/s}$, of <i>P</i> is given by $v = 2t^2 - 16t + 30$	
	(a) Find the acceleration, in m/s^2 , of P when $t = 5$	(2)
	P comes to instantaneous rest at the points M and N at times t_1 seconds and t_2 seconds where $t_2 > t_1$	
	(b) Find the exact distance MN	(8)
••••		

Question 4 continued
(Total for Question 4 is 10 marks)



5 A solid cuboid has width x cm, length 4x cm and height h cm.

The volume of the cuboid is 75 cm³ and the surface area of the cuboid is S cm²

(a) Show that $S = 8x^2 + \frac{375}{2x}$

(4)

Given that x can vary, using calculus,

- (b) (i) find to 3 significant figures, the value of x for which S is a minimum,
 - (ii) justify that this value of x gives a minimum value of S

(5)

(c) Find, to 3 significant figures, the minimum value of S

(2)

Question 5 continued	

Question 5 continued	
(T)	otal for Question 5 is 11 marks)



6 Solve the equation	n
-----------------------------	---

$$\log_2 x^3 + \log_4 x^2 - 3\log_x 2 = 0$$

giving your	answers	to 3	significant	figures


(8)		
(0)		
(0)		

(0)



Question 6 continued
/T-4-1 f
(Total for Question 6 is 8 marks)





When x is increased to $(x + \delta x)$, y increases to $(y + \delta y)$ where δx and δy are small.

(a) Show that
$$\delta y \approx \frac{e^{2x}(4x-7)}{(2x-3)^{\frac{3}{2}}} \delta x$$

(7)

Given that x = 2.5

(b) find an estimate, to 2 significant figures, of the value of δy when the value of x increases by 0.2%

(3)

|
 |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | |
|
 |

>>> 	
	Question 7 continued
	Question / continued
A	
E	
S S	
至	
DO NOT WRITE IN THIS AREA	
ō	
A A	
AR	
THIS AREA	
NOTWRITEIN	
5	
Ž	
5	
₹	
5	
<u> </u>	
乙	
X	
DO NOT WRITE IN THIS AREA	
0	
XXX '	



Question 7 continued	

Question 7 continued	
(Tatal	for Question 7 is 10 marks)
(Total	ZOZ ZUCSHOII / IS IV IIIdI NS)



8	$f'(x) = 18x^2 - 2x + 13$	
	Given that $(2x-1)$ is a factor of $f(x)$	
	show that the curve with equation $y = f(x)$ has only one intersection with the x-axis.	(9)

Question 8 continued
(Total for Question 8 is 9 marks)



9 (a) Using the formulae on page 2, show that

$$(i) \cos^2 A = \frac{\cos 2A + 1}{2}$$

(ii)
$$\sin^2 A = \frac{1 - \cos 2A}{2}$$

(4)

(b) Show that

$$(2\sin x - \cos x)(\sin x - 3\cos x) = \frac{1}{2}(\cos 2x - 7\sin 2x + 5)$$

(5)

$$y = (2\sin x - \cos x)(\sin x - 3\cos x)$$

(c) Solve, for $0^{\circ} \leqslant x \leqslant 180^{\circ}$ the equation, $\frac{dy}{dx} = 0$

Give your answers to the nearest whole number.

(4)

	$\Pi\Pi\Pi$	ШП						ШП				П
						Ш					Ш	Ш
Ρ	7	2	8	6	4 /	Α .	0 :	2 :	3 :	3	2	

Question 9 continued	

Question 9 continued	
	(Total for Question 9 is 13 marks)



10 O, A and B are fixed points such that

$$\overrightarrow{OA} = (b+1)\mathbf{i} + b\mathbf{j}$$

$$AB = 3i$$

 $\overrightarrow{OA} = (b+1)\mathbf{i} + b\mathbf{j}$ $\overrightarrow{AB} = 3\mathbf{i}$ The unit vector parallel to \overrightarrow{OB} is $\frac{\sqrt{17}}{34} [(3a+2)\mathbf{i} + b\mathbf{j}]$

Given that a and b are constants where a > 0 and b > 0

find the exact value of

- (i) a
- (ii) *b*

- /	1	1 /	"
- (Ĺ	LV	נע

• • • •

Question 10 continued	

Question 10 continued	
(To	tal for Question 10 is 10 marks)



11 $f(x) = 10 + 6x - x^2$

Given that f(x) can be written in the form $A(x + B)^2 + C$ where A, B and C are constants,

(a) find the value of A, the value of B and the value of C

(4)

- (b) Hence, or otherwise, find
 - (i) the value of x for which f(x) has its greatest value
 - (ii) the greatest value of f(x)

(2)

The curve C has equation y = f(x)

The curve S with equation $y = x^2 - x + 13$ intersects curve C at two points.

(c) Find the *x* coordinate of each of these two points.

(3)

(d) Use algebraic integration to find the exact area of the finite region bounded by the curve C and the curve S.

(5)

|
 | |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--|
|
 | |
|
 | |
|
 | |



DO NOT WRITE IN THIS AREA

Question 11 continued	
	(Total for Question 11 is 14 marks) TOTAL FOR PAPER IS 100 MARKS