

Question Number	Scheme	Marks
2(a)	$(-10\mathbf{i} + a\mathbf{j}) + (b\mathbf{i} - 5\mathbf{j}) + (2a\mathbf{i} + 7\mathbf{j}) = 3(3\mathbf{i} + 4\mathbf{j})$ $a - 5 + 7 = 12 \Rightarrow a = 10$ $-10 + b + 2a = 9 \Rightarrow b = -1$	M1 M1 A1 M1 A1 (5)
(b)	$20\mathbf{i} + 20\mathbf{j} = \mathbf{u} + 4(3\mathbf{i} + 4\mathbf{j})$ $\mathbf{u} = (8\mathbf{i} + 4\mathbf{j})$ $u = \sqrt{8^2 + 4^2} = \sqrt{80} = 8.9 \text{ (or better)}$	M1 A1 M1 A1 (4) 9
	Notes	
2(a)	First M1 for applying $\mathbf{F} = m\mathbf{a}$; need all terms but allow slips and allow m instead of 3 Second M1 (independent but M0 if they have 0 instead of $m\mathbf{a}$) for equating <i>coefficients</i> of j First A1 for $a = 10$ Third M1 (independent but M0 if they have 0 instead of $m\mathbf{a}$) for equating <i>coefficients</i> of i Second A1 for $b = -1$	
(b)	First M1 for applying $\mathbf{v} = \mathbf{u} + t\mathbf{a}$; need all terms and must be vector u First A1 for $8\mathbf{i} + 4\mathbf{j}$ Second M1 (independent) for finding magnitude of their vector u Second A1 for $\sqrt{80}$ or 8.9 or better	