

- 9 The finite region R enclosed by the y -axis, the straight line with equation $y + 2x = 13$ and the curve with equation $y = x^2 - 2$, is defined for points with coordinates (x, y) with $x \geq 0$

The region R is rotated through 360° about the y -axis.

Use algebraic integration to find the volume of the solid generated.
Give your answer in terms of π .

(9)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 9 continued

Handwriting practice area with horizontal dotted lines.



P 6 1 8 8 3 A 0 2 7 3 6

Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 9 continued

Handwriting practice area with horizontal dotted lines.

(Total for Question 9 is 9 marks)



10 (a) Use the formula for $\cos(A + B)$ to show that $\cos 2A = 2\cos^2 A - 1$ (2)

(b) Show that $\cos 4A = 8\cos^4 A - 8\cos^2 A + 1$ (4)

(c) Solve the equation $\cos^2\left(\frac{\theta}{4} + \frac{\pi}{24}\right)\left[\cos^2\left(\frac{\theta}{4} + \frac{\pi}{24}\right) - 1\right] = -\frac{1}{16}$ $0 \leq \theta < 2\pi$

Give your answers in terms of π . (5)

$$f(A) = 4\cos^4 A - 4\cos^2 A + 1$$

(d) Using calculus, find the exact value of $\int_{\frac{\pi}{6}}^{\frac{\pi}{2}} f(A) dA$

Give your answer in the form $a\pi - b\sqrt{c}$ where a and b are fractions in their lowest terms and c is a prime number.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 10 continued

Handwriting practice area with horizontal dotted lines.



P 6 1 8 8 3 A 0 3 1 3 6

Question 10 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 10 continued**(Total for Question 10 is 15 marks)**

P 6 1 8 8 3 A 0 3 3 3 6

11 The quadratic equation $x^2 - px + q = 0$ where $p > 0$, has roots α and β .

Given that $2\alpha\beta = 3$ and that $4(\alpha^2 + \beta^2) = k^2 - 6k - 3$ where $k > 3$

(a) (i) write down the value of q ,

(ii) find an expression, in terms of k , for p .

(5)

Given also that $7\alpha\beta = 3(\alpha + \beta)$

(b) find the value of k .

(2)

(c) Hence form an equation, with integer coefficients, which has roots

$$\frac{\alpha}{\alpha + \beta} \text{ and } \frac{\beta}{\alpha + \beta}$$

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 11 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 1 8 8 3 A 0 3 5 3 6

Question 11 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 11 is 12 marks)**TOTAL FOR PAPER IS 100 MARKS**