

Leave
blank

7. A helicopter is hovering at rest above horizontal ground at the point H . A parachutist steps out of the helicopter and immediately falls vertically and freely under gravity from rest for 2.5 s. His parachute then opens and causes him to immediately decelerate at a constant rate of 3.9 m s^{-2} for T seconds ($T < 6$), until his speed is reduced to $V \text{ m s}^{-1}$. He then moves with this constant speed $V \text{ m s}^{-1}$ until he hits the ground. While he is decelerating, he falls a distance of 73.75 m. The total time between the instant when he leaves H and the instant when he hits the ground is 20 s.

The parachutist is modelled as a particle.

- (a) Find the speed of the parachutist at the instant when his parachute opens. (1)
- (b) Sketch a speed-time graph for the motion of the parachutist from the instant when he leaves H to the instant when he hits the ground. (2)
- (c) Find the value of T . (5)
- (d) Find, to the nearest metre, the height of the point H above the ground. (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 7 continued

DO NOT WRITE IN THIS AREA



Leave
blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued

Handwriting practice area with horizontal lines.



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave
blank

Question 7 continued

Q7

(Total 12 marks)



P 6 1 9 0 2 A 0 1 9 2 4