Please check the examination details bel	ow before entering your candidate information
Candidate surname	Other names
Centre Number Candidate No Pearson Edexcel Inter	
<b>Time</b> 2 hours	Paper reference 4PM1/02R
Further Pure Mat	hematics
Calculators may be used.	Total Marks

#### **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- You must NOT write anything on the formulae page.
   Anything you write on the formulae page will gain NO credit.

#### **Information**

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

#### **Advice**

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶







#### **International GCSE in Further Pure Mathematics Formulae sheet**

#### Mensuration

**Surface area of sphere** =  $4\pi r^2$ 

**Curved surface area of cone** =  $\pi r \times \text{slant height}$ 

Volume of sphere =  $\frac{4}{3}\pi r^3$ 

#### Series

#### **Arithmetic series**

Sum to *n* terms,  $S_n = \frac{n}{2} [2a + (n-1)d]$ 

#### Geometric series

Sum to *n* terms, 
$$S_n = \frac{a(1-r^n)}{(1-r)}$$

Sum to infinity, 
$$S_{\infty} = \frac{a}{1-r} |r| < 1$$

#### **Binomial series**

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots$$
 for  $|x| < 1, n \in \mathbb{Q}$ 

#### **Calculus**

#### **Quotient rule (differentiation)**

$$\frac{\mathrm{d}}{\mathrm{d}x} \left( \frac{\mathrm{f}(x)}{\mathrm{g}(x)} \right) = \frac{\mathrm{f}'(x)\mathrm{g}(x) - \mathrm{f}(x)\mathrm{g}'(x)}{\left[\mathrm{g}(x)\right]^2}$$

#### **Trigonometry**

#### Cosine rule

In triangle *ABC*:  $a^2 = b^2 + c^2 - 2bc \cos A$ 

$$\tan\theta = \frac{\sin\theta}{\cos\theta}$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

### Logarithms

$$\log_a x = \frac{\log_b x}{\log_b a}$$



There are no questions on this page.



## Answer all ELEVEN questions.

## Write your answers in the spaces provided.

## You must write down all the stages in your working.

1 The position vector of the point A is  $(3\mathbf{i} - 2\mathbf{j})$ , referred to a fixed origin O.

The point B is such that  $\overrightarrow{AB} = (6\mathbf{i} + 8\mathbf{j})$ 

(a) Find the position vector of B as a simplified expression in terms of  $\mathbf{i}$  and  $\mathbf{j}$ 

(2)

(b) Find the magnitude of vector  $\overrightarrow{AB}$ 

(1)

(c) Find a unit vector, in terms of **i** and **j**, that is parallel to  $\overrightarrow{AB}$ 

(2)




Question 1 continued	
	(Total for Organian 1 is 5
	(Total for Question 1 is 5 marks)



The volume of the cuboid increases at a constant rate of $8 \mathrm{m}^3/\mathrm{s}$	
Find the rate of increase, in m/s, of x when $x = 2$ metres.	
That the face of merease, in m/s, of w when w = 2 metres.	(6)

	Question 2 continued
AREA	
(THIS	
WRITE IN	
NOT W	
DON	
AREA	
Ž T T	
WRITEIN	
W TO	
DO	
A	
DO NOT WRITE IN THIS AREA	
五二五	
VRITE	
NOTV	
DO	
	(Total for Question 2 is 6 marks)



3	A geometric series has first term $a$ and common ratio $r$ , where $r > 0$	
	Given that the 3rd term of the series is 5 and that the 5th term of the series is $\frac{5}{2}$	
	(a) find	
	(i) the exact value of $r$	
	(ii) the value of a	(4)
	(b) Find the sum to infinity of this series. Give your answer in the form $p + q\sqrt{2}$ where $p$ and $q$ are integers.	
		(2)

Question 3 continued			
			,
	 		,
	(TE) 1 1 0	or Question 3 is 6 r	1 \



4	$f(x) = x^3 + px^2 + qx + 7$ where p and q are integers.	
	(x + 1) is a factor of $f(x)The remainder when f(x) is divided by (x + 2) is -5$	
	(a) Find the value of $p$ and the value of $q$	
		(5)
	(b) Hence, show that $f(x) = 0$ has only one real root.	(3)
••••		
••••		
••••		
••••		
••••		
••••		

(Total	for	Question	4	is	8	marks

5 (a) Complete the table of values for  $y = e^{3x-2}$  giving your answers to 2 decimal places.

x	0	0.25	0.5	0.75	1
y	0.14				2.72


(2)

(b) On the grid opposite, draw the graph of  $y = e^{3x-2}$  for  $0 \le x \le 1$ 

(2)

(c) By drawing a suitable straight line on the grid, obtain an estimate, to one decimal place, of the root of the equation  $3x = 2 + \ln(3 - x)$ 

**(3)** 






Turn over for a spare grid if you need to redraw your graph.

DO NOT WRITE IN THIS AREA

Question 5 continued	

# **Question 5 continued**

Only use this grid if you need to redraw your graph.



(Total for Question 5 is 7 marks)






Diagram **NOT** accurately drawn

**(6)** 

Figure 1

Figure 1 shows a right pyramid VABCD with vertex V and square base ABCD.

Each of the edges of the pyramid has the same length.

Find the size, in degrees to one decimal place, of the angle between the plane CVD and the base ABCD.

	Question 6 continued
AREA	
HIS	
NOT WRITE	
NOT	
DO	
AREA	
HIS A	
EINT	
WRITE	
OTV	
DO	
W.	
IS AR	
N.	
RITE	
DO NOT WRITE IN THIS AREA	
DO N	
	(Total for Question 6 is 6 marks)



7 (a) Solve the equation

$$\cos(3x - 15)^\circ = \frac{\sqrt{3}}{2}$$
 for  $0 \le x < 180$ 

(4)

(b) Solve, giving your solutions to one decimal place where appropriate,

$$3 \tan y^{\circ} + 4 \sin y^{\circ} = 0$$
 for  $-180 \le y < 180$ 

(4)

(c) Solve, giving your solutions to one decimal place where appropriate,

$$\cos \theta^{\circ} = 3 \sin^2 \theta^{\circ} - 1$$
 for  $-180 \leqslant \theta < 180$ 

**(4)** 







Question 7 continued		



DO NOT WRITE IN THIS AREA

$\times$	ς.
$\times f \ominus \times$	5
	5
$\times$	
0	ς.
××××	
$\times \times \times \times$	
$\times$	ς.
×O×	ς.
ğ	
	<
$\leftrightarrow$	
$\times \times \times$	
<b>3</b>	
	2
H	
XXXXX	2
$\times$	
$\times$	>
$\times$	>
	>
$\times = \times$	
$\times \times \times \times$	>
$\times$	S
	5
$\times \times \times$	
$\times$	S
	S
$\sim\sim\sim$	
UA.	
$\langle \times \times \times \times \rangle$	
<b>&gt;</b>	<
$\circ$	
	<
$\langle \mathbf{n} \rangle$	2
	2
$\times \times \times \times$	2
$\times\!\!\times\!\!\times\!\!\times$	
$\times\!\!\times\!\!\times\!\!\times$	>
$\times\!\!\times\!\!\times\!\!\times$	
$\times\!\!\times\!\!\times\!\!\times$	>
$\times\!\!\times\!\!\times\!\!\times$	
$\times\!\!\times\!\!\times\!\!\times\!\!\times$	
$\times\!\!\times\!\!\times\!\!\times$	
	S
	5
$\times\times\times$	ς.
	ς.
	ς.
$\sim\sim$	
$\times\!\!\times\!\!\times\!\!\times$	
	}
	}
	}
	}
	}
	}
D	
D	
DO	
WTONO	
WTONO	
O NOT WR	
O NOT WR	
O NOT WR	
O NOT WRIT	
O NOT WRIT	
O NOT WRIT	
O NOT WR	
O NOT WRIT	
O NOT WRITE IN TH	
O NOT WRITE IN TH	
O NOT WRITE IN TH	
O NOT WRIT	
O NOT WRITE IN THIS /	
O NOT WRITE IN THIS /	
O NOT WRITE IN THIS /	
O NOT WRITE IN THIS /	
O NOT WRITE IN THIS AR	
O NOT WRITE IN THIS ARE	
O NOT WRITE IN THIS ARE	
O NOT WRITE IN THIS AR	

Question 7 continued	

*****
****
****
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
EA
RITE IN THIS ARE
$\times$
$\times$
$\times$
$\times\!$
$\otimes \Xi \otimes$
$\times\!\!\!+\!\!\!+\!\!\!\times$
$\times\!$
$\times$
$\times$
$\times$
$\otimes \mapsto \otimes$
imes
$\propto \sim \times \times$
$\times \times \times \times$
$\times$
$\times$
$\times\!\!\times\!\!\times$
$\times \odot \times$
$\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
XXXX
$\times\!\!\times\!\!\times\!\!\times$
*****
$\times\!\!\times\!\!\times\!\!\times$
*****
$\times\!\!\times\!\!\times\!\!\times$
XXXX
XXXX
XXXX
$\times\!\!\times\!\!\times\!\!\times$
$\times$ MM $\times$
$\times$
XX-XX
$\times$
$\times\!\!\!\times\!\!\!\times\!\!\!\times$
$\times$
ITE IN THIS AREA
$\times$
$\times \times \times \times$
$\times$
<u></u>
O NOT WRI
DO NOT WR
A DO NOT WR
EA DO NOT WR
EA DO NOT WR
EA DO NOT WR
EA DO NOT WR
AREA DO NOT WR
AREA DO NOT WR
AREA DO NOT WR
AREA DO NOT WR
AREA DO NOT WR
NTHIS AREA DO NOT WR
NIHIS AREA DO NOT WR
NIHIS AREA DO NOT WR
EIN THIS AREA DO NOT WR
EIN THIS AREA DO NOT WR
EIN THIS AREA DO NOT WR
RITE IN THIS AREA DO NOT WR
RITE IN THIS AREA DO NOT WR
WRITE IN THIS AREA DO NOT WR
WRITE IN THIS AREA DO NOT WR
WRITE IN THIS AREA DO NOT WR
OT WRITE IN THIS AREA DO NOT WR
OT WRITE IN THIS AREA DO NOT WR
NOT WRITE IN THIS AREA DO NOT WR
OT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR
O NOT WRITE IN THIS AREA DO NOT WR

Question 7 continued
(Total for Question 7 is 12 marks)






Figure 2

Figure 2 shows part of the curve C with equation  $y = e^{3x} - 1$  and part of the curve D with equation  $y = 9 - 9e^{-3x}$ 

The curves intersect at the origin O and the point A.

(a) (i) Show that the x coordinate of the point A satisfies the equation

$$(e^{3x})^2 - 10e^{3x} + 9 = 0$$

(ii) Hence, show that the x coordinate of the point A is  $\frac{1}{3} \ln 9$ 

(5)

The finite region bounded by C and by D is shown shaded in Figure 2.

(b) Use calculus to find the exact area of this region.

**(6)** 

Question 8 continue	ed		



DO NOT WRITE IN THIS AREA

X	$\rangle$	$\Diamond$	♦	Ö	Ŏ.
X	2	♦	$\bigcirc$	Ŷ	♦.
X	d	Ř	Ď	X	X
Ŏ	♂	8	y.	X	X
0	ğ	K	3	X	Χ
♦	₫	×	χ	X	X
Ó	ζ	X	×	X	X
Ó	Š	S	6		X
$\Diamond$	9	Z	7	X	X
Q	A	6	a	X	Χ.
X			4	Ö	Ŏ.
X	×	X	d	$\Diamond$	Ο.
×	$\geq$		C	$\Diamond$	♦.
$\times$	ì		S	$\Diamond$	♦
X	à	ë	v	$\Diamond$	♦.
X	à	0	C		♡.
Ö	4	Я	U	×	2
Ö	ě	H		×	X
♦	G		ч	×	X
$\Diamond$	ς	$\times$			X
♦	4	И	и	×	X
$\Diamond$	ς		X	X	X
	Ŷ	8	Z	X	Χ.
×	S	4	8	Ö	Ŏ.
X				Ö	Õ.
×	Ž	٥	û	$\Diamond$	٥.
×	y	$\overline{}$	Ų.	$\Diamond$	٥.
X	×	Ġ	Ŕ	$\Diamond$	♦.
X	ĕ	Ä		Ø	$\Diamond$
ŏ	ð	Ħ	Ź	X	
Ŏ	4	И	Я		X
Ó	Ĉ	X	×	X	X
♦				X	X
0	Ġ	ø	7	X	Χ
Q	Ŷ	Ħ			X
9	ģ	S	썯	8	X.
X	g	ø	ø	ŏ	Ŏ.
X	Ś	š	Ö	Ó	Ŏ
X	S	ä	⋛	Ŏ	Ó.
X	Ź	Ó	<b>&lt;</b>	ø	Q
X	×	Q	0	Ŷ	♦.
	Ż	0	$\Diamond$	Q	Ω.
X	2		S	X	$\otimes$
Ö	Š			X	X
Ŏ	Ĉ,	X	X		X
♦	Ċ.	X	X		X
0	Ç		Х	X	X
Q	ς		X	×	X.
	S		X		X
Q	S	8	ŏ	Ö	8.
×	S	Ŏ	Ŏ	Ö	Ŏ.
×	×		٥	$\Diamond$	◌.
	$\geq$	0	♦	$\Diamond$	♦.
		$\vee$	V	$^{\vee}$	$\vee$
×	۶		X	Ş	X.
8	ξ	Š	X	×	Š
8	8	X	X	×	Š
	2	X	X	× ×	X
	\{ \{ \}			>	X
				× ×	X
	Ş	ž			
	Ş	ž			
	Ş	ž			
	Ş	ž			
	Ş	ž			
	Ş				
	Ş				
	Ş				

Question 8 continued

	Question 8 continued
AREA	
HIS	
NOT WRITE IN	
NOT	
DQ	
AREA	
THIS	
WRITEIN	
T WR	
ONO.	
۵	
A	
ARE	
THE	
ITE II	
DO NOT WRITE IN THIS AREA	
OO NK	
	(Total for Question 8 is 11 marks)



(a) Write  $\frac{3}{(3-x)^3}$  in the form  $a(1-bx)^{-3}$ 

where a and b are fractions in their lowest terms.

- (2)
- (b) Expand  $\frac{3}{(3-x)^3}$  in ascending powers of x up to and including the term in  $x^3$ Express each coefficient as a fraction in its lowest terms.

(3)

- (c) (i) Use a suitable value of x with your expansion in part (b), to obtain an approximation for  $\frac{24}{125}$  to 5 decimal places.
  - (ii) Find the percentage error, to 2 decimal places, of your approximation from the actual value.

(4)

| <br> |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| <br> |
|      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
|      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| <br> |
|      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
|      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| <br> |
|      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| <br> |
|      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
|      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| <br> |

$\otimes$ 1	
$\otimes$	
$\otimes$	Question 9 continued
$\otimes$	
$\otimes$	
$\otimes$	
$\otimes$	
$\otimes$	
$\otimes$	
×	
$\otimes$	
$\otimes$	
$\otimes$	
$\otimes$	
$\otimes$ $ $	
$\otimes$	
$\stackrel{>}{\sim}$	
$\stackrel{>}{\sim}$	
§	
$\otimes$	
$\otimes$	
<b>%</b>	
$\otimes$	
$\otimes$	
$\otimes$ 1	
⊗ 1	
$\stackrel{\circ}{\mathbb{R}}$	



DO NOT WRITE IN THIS AREA

$\times\!\!\times\!\!\times$	$\times$
$\times \times \times$	XX.
$\sim$	XX.
$\times \otimes$	KX.
	XX.
X	XX.
	22
$\propto \times \times$	ZX.
	XX.
$\vee_{\sim}$	XX.
X	$\times \times$
	$\times \times$
	XX.
	$\times \times$
$\leftrightarrow$	iXX.
	XX.
$\sim$	$\times \times$
	XX.
CON.	$\times \times$
	ixx.
$\sim\sim$	KX.
$\times \times$	KX.
$\times m$	KXZ.
	$\langle \chi \chi \rangle$
$\times \times \times$	
$\times \times \times$	$\otimes$
$\times$	$\infty$
$\times \times $	KO.
$\times \times \times$	
X	68
$\times$	ĸΧ.
$\sim$	KK.
	88.
VA	KX.
$\times\!\!\times\!\!\times$	XX.
	$\times \times$
	$\times \times$
	XX.
	XX.
XIII	$\bowtie$
	XX.
	iXX.
0.00X	XX.
$\times\!\!\times\!\!\times$	××.
$\times\!\!\times\!\!\times$	~~
$\times\!\times\!\times$	~~
$\times\!\!\times\!\!\times$	$\sim$
$\times\!\!\times\!\!\times$	$\sim$
$\times\!\!\times\!\!\!\times$	$\Diamond \Diamond$
$\times\!\!\times\!\!\times$	$\sim$
$\times\!\!\times\!\!\times$	$\sim$
	~~
XXX	~~
	~~
$\sim$	88
$\leftrightarrow\!$	~~
	$\otimes$
	$\otimes$
	$\otimes$
	$\overset{\times}{\otimes}$
	*
	×
D(	
00	
MALONO	
MALONO	
O NOT WE	
O NOT WE	
O NOT WE	
O NOT WRIT	
O NOT WRIT	
MALONO	
O NOT WRIT	
O NOT WRITEIN TH	
O NOT WRIT	
O NOT WRITE IN THIS Y	
O NOT WRITE IN THIS Y	
O NOT WRITE IN THIS Y	
O NOT WRITE IN THIS AR	
O NOT WRITE IN THIS AR	
O NOT WRITE IN THIS ARE	
O NOT WRITE IN THIS AR	

Question 9 continued	

Question 9 continued	
	(Total for Question 9 is 9 marks)



## **10** A curve *C* has equation

$$y = \frac{7x - 2}{2x - 3} \qquad x \neq \frac{3}{2}$$

- (a) Write down an equation of the asymptote to C that is
  - (i) parallel to the y-axis,
  - (ii) parallel to the *x*-axis.

(2)

(b) Find the coordinates of the points of intersection of C with the coordinate axes.

(2)

(c) Using calculus, show that at every point on the curve, the gradient of C is negative.

(4)

(d) Using the axes on the opposite page, sketch *C*. Show clearly and label with their equation any asymptotes and the coordinates of the points of intersection of *C* with the coordinate axes.

(3)

The straight line l is the normal to C at the point A. The x coordinate of A is positive and the gradient of l is 17


The line l also intersects C at the point B.

(e) Find the exact coordinates of B.

(7)









DO NOTWRITE IN THIS AREA

X	XX	$\otimes \otimes$
Ö	$\approx$	$\times$
♦		$\bowtie$
X		
X		$\bowtie$
Ó		$\times\!\!\times$
X	Ż	XX.
X		$\otimes \rangle$
Ö	W	$\bowtie$
Q		$\otimes$
X		$\otimes \otimes$
X		$\otimes$
Ş	-	
2	策	iXX.
X		XX.
8	XX	iXX.
Ŏ	$\infty$	$\bowtie$
Q		$\otimes$
X	$\times\!\!\times\!\!\times$	
X	**	$\otimes$
0		XX.
Ø	$\otimes$	XX
X	$\nabla$	KØ.
X		500
Ö	**	XX.
ø	97	88
X	M	$\Diamond \Diamond$
X	8	$\otimes$
Ó		:XX
S	20	$\bowtie$
X		
X		$\otimes$
Ŏ	1	iXX.
$\Diamond$		XX.
X	XX	$\langle \rangle \rangle$
X	88	$\times\!\!\times\!\!\times$
Ŏ	$\!$	$\times\!\!\times$
ø	$\stackrel{\Diamond}{\otimes}$	XX.
X	$\times\!\!\times\!\!\times$	$\langle \rangle$
X	$\times\!\!\times\!\!\times$	$\times\!$
Ö	$\Diamond \Diamond$	$\times\!\!\times$
Ø	$\otimes$	$\times\!\!\times$
X	$\times\!\!\times\!\!\times$	$\times\!$
X	88	$\times$
Ŏ	$\Diamond \Diamond$	$\times\!\!\times$
$\Diamond$	$\times\!\!\!\times\!\!\!\!\times$	$\times\!\!\times$
X	XX	××.
ŏ	$\langle \rangle \langle$	$\otimes$
Ó	$\langle \rangle \langle$	$\otimes \! \times$
$\Diamond$	98	XX.
X	$\times\!\!\times\!\!$	$\langle \rangle \rangle$
X	$\propto \propto$	$\times$
Ó	$\Diamond\Diamond$	$\times\!\!\times$
X	XX.	XX.
X		$\bowtie$
X	ĕ	$\otimes$
Ó	⟨⊘	$\bowtie$
2	×	$\times\!\!\times$
X		$\Diamond \Diamond$
ŏ	-	KX.
Ó	0	$\bowtie$
X		XX.
X		K⊗.
ŏ	×	$\otimes$
Ó	2	XX.
X		XX.
X		$\otimes$
ŏ	₩	$\otimes$
Ó	剱	$\bowtie$
X		iXX.
X	**!	
X	$\approx$	$\otimes$
Ó		<u> </u>
2	XX.	XX.
	$\times\!\!\times\!\!\!\times$	KØ.
Ø	$\times \times$	$\times$
А		$\wedge \wedge 2$
		KOO.
X		$\otimes$
Š	T	$\otimes$
8	Z	$\otimes$
\ \ \	XX	
	HS ARD	
	XX	

Question 10 continued

Question 10 continued	
(Tatal	for Question 10 is 18 marks)
(Total	TOT QUESTION TO IS TO MAINS)






Diagram **NOT** accurately drawn

Figure 3

Figure 3 shows a solid metal right circular cylinder of radius rcm and height hcm.

The total surface area of the cylinder is 600 cm<sup>2</sup>

The volume of the cylinder is  $V \text{cm}^3$ 

(a) Show that  $V = 300r - \pi r^3$ 

(4)

Given that r can vary,

(b) (i) use calculus to show that the exact value of r for which V is a maximum is

$$r = \sqrt{\frac{100}{}}$$

(ii) justify that this value of r gives a maximum value of V

(5)

The cylinder is melted down and reformed into a sphere of radius  $p \, \text{cm}$ .

(c) Find, to one decimal place, the greatest possible value of p

(3)

 	 •••••	 •••••	 	 

<b>Question 11 continued</b>		



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 11 continued	
	(Total for Question 11 is 12 marks)
	TOTAL FOR PAPER IS 100 MARKS

