


M1 October 2016 (IAL) (MA)

Q(a)



$$\begin{aligned} \text{C.L.M : } 2m(2u) + 3m(-u) &= 2m(V_p) + 3m(V_q) \\ \Rightarrow u &= 2V_p + 3V_q \end{aligned}$$

Impulse on P acts towards the left.

$$\therefore -5mu = 2m(V - u) = -5mu = 2m(V_p - 2u)$$

$$\Rightarrow -5u = 2V_p - 4u$$

$$\Rightarrow 2V_p = -u \quad \therefore V_p = -\frac{u}{2}$$

$$\text{so speed of P} = \boxed{\frac{u}{2}}$$

- b)  $V_p < 0$  so P is travelling in the opposite direction to that which we assumed in the diagram.  
So yes the dir. of P has been reversed.

c) from (a),  $u = 2V_p + 3V_q$

$$\Rightarrow u = 2\left(-\frac{u}{2}\right) + 3V_q$$

$$\Rightarrow 3V_q = 2u \quad \therefore V_q = \boxed{\frac{2u}{3}} = \text{speed}_q.$$

(Q2a)  $\sum \vec{F} = m\vec{a}$  :

$$\begin{pmatrix} -10 \\ a \end{pmatrix} + \begin{pmatrix} b \\ -5 \end{pmatrix} + \begin{pmatrix} 2a \\ 7 \end{pmatrix} = \underbrace{3}_m \begin{pmatrix} 3 \\ 4 \end{pmatrix}^{\vec{a}}$$

$$\Rightarrow \begin{pmatrix} 2a + b - 10 \\ 2 + a \end{pmatrix} = \begin{pmatrix} 9 \\ 12 \end{pmatrix}$$

equating  $\underline{i}$  :  $2a + b - 10 = 9$   
 $\Rightarrow 2a + b = 19 \quad \sim \textcircled{1}$

equating  $\underline{j}$  :  $2 + a = 12$   
 $\Rightarrow \boxed{a = 10} //$

$\downarrow$   
 $\textcircled{1} : 2(10) + b = 19$   
 so  $\boxed{b = -1} //$

b)  $a = \frac{v-u}{t}$  :  $3\underline{i} + 4\underline{j} = \frac{(20\underline{i} + 20\underline{j}) - (\underline{u})}{(4)}$


$\times 4$   
 $\Rightarrow 12\underline{i} + 16\underline{j} = 20\underline{i} + 20\underline{j} - (\underline{u})$

$\Rightarrow \underline{u} = 20\underline{i} - 12\underline{i} + 20\underline{j} - 16\underline{j}$

$\Rightarrow \underline{u} = 8\underline{i} + 4\underline{j} //$

hence  $u = |\underline{u}| = \sqrt{8^2 + 4^2} = \boxed{4\sqrt{5}} \text{ ms}^{-1}$

Q3)



We are told:  $R_D = 2R_C$

$$\begin{aligned}
 R(\uparrow): \quad R_C + R_D &= 12g + 3g \\
 R_C + 2R_C &= 15g \\
 \therefore R_C &= \frac{15g}{3} = 5g // \rightarrow R_D = 10g //
 \end{aligned}$$

$$M(A): R_C(3) + R_D(x) = 12g(4) + 3g(8)$$

$$5g(3) + 10g(x) = 48g + 24g$$

$$10x + 15 = 72$$

$$10x = 57 \quad \therefore x = \frac{57}{10} = \boxed{5.7\text{m}}$$

Q4a)  $p = r_0 + vt$

$$p = \begin{pmatrix} -5 \\ 9 \end{pmatrix} + t \begin{pmatrix} 1 \\ -2 \end{pmatrix} \Rightarrow \boxed{p = (-5+t)\underline{i} + (9-2t)\underline{j}}$$

b)  $\underline{j} = 2$  when P is due west of A.

$$\therefore 9 - 2t = 2$$

$$\Rightarrow 7 = 2t \quad \therefore t = \frac{7}{2} //$$

$$\begin{aligned}
 \text{at } t = \frac{7}{2}, \quad p &= (-5 + \frac{7}{2})\underline{i} + (9 - 7)\underline{j} \\
 \boxed{p &= -\frac{3}{2}\underline{i} + 2\underline{j}}
 \end{aligned}$$

c) if they move along parallel lines then their velocity vectors are parallel.

$$\Rightarrow \begin{pmatrix} 2b-1 \\ 5-2b \end{pmatrix} = c \begin{pmatrix} 1 \\ -2 \end{pmatrix} \quad \text{for some } c.$$

$$\therefore 2b-1 = c \quad \text{--- (1) (equating i)}$$

$$5-2b = -2c \quad \text{--- (2) (equating j)}$$


$$\underline{\textcircled{1} + \textcircled{2}} : \cancel{2b} - 1 + 5 - \cancel{2b} = c - 2c$$

$$4 = -c \quad \therefore c = -4 //$$

$$\text{on d } 2b-1 = -4$$

$$\Rightarrow b = \frac{-4+1}{2} = \boxed{-\frac{3}{2}}$$

Q5)



$$R(\uparrow): R = (8\sin 40 + 0.5g\cos 30)$$

$$R(\nwarrow): 8\cos 40 = F + 0.5g\sin 30$$

but  $F = \mu R$  as P is on the point of sliding up the plane.

$$\therefore 8\cos 40 = \mu R + 0.5g\sin 30$$

$$8\cos 40 = \mu (8\sin 40 + 0.5g\cos 30) + 0.5g\sin 30$$

$$\text{So } \mu = \frac{8\cos 40 - 0.5g\sin 30}{8\sin 40 + 0.5g\cos 30}$$

$$= \boxed{0.392}$$

Q6)

For A :

$$S = S_A$$

$$U = 35$$

$$V = V_A$$

$$a = 0.4$$

$$t =$$

For B :

$$S = S_B$$

$$U = 44$$

$$V = V_B$$

$$a = 0.5$$

$$t =$$

When B overtakes A,  $S_B - 200 = S_A$  // (since B starts 200m behind,

$$S_A = 35t + 0.2t^2$$

$$\text{and } S_B = 44t + 0.25t^2$$

$$\text{but using } S_B - 200 = S_A,$$

$$\Rightarrow 44t + 0.25t^2 - 200 = 35t + 0.2t^2$$

$$\Rightarrow 0.05t^2 + 9t - 200 = 0$$

$$\times 20 \Rightarrow t^2 + 180t - 4000 = 0$$

By Quadratic Formula :

$$t = 20$$

or

$$t = -200$$


//

t > 0 !

So at  $t = 20$  B overtakes A.

using  $v = u + at$  with B =  $V_B = 44 + 0.5(20)$   
 $= \boxed{54}$

(Q7a)



b)  $a = \frac{v-u}{t}$  for ① :  $1 = \frac{V-0}{a}$   
 $\therefore V = a$

$a = \frac{v-u}{t}$  for ③ :  $-0.5 = \frac{0-V}{180-(a+T)}$

$$-90 + 0.5(T+V) = -V$$

$$90 - \frac{1}{2}(T+V) = V$$


$$180 - T - V = 2V$$

$$\therefore T = 180 - 3V$$



c) Total area under graph = 4800 //

$$\text{Area of trapezium} = \frac{(a+b)h}{2} (= 4800)$$



$$\therefore 4800 = \frac{1}{2} (180 + 180 - 3V)V$$

$$9600 = 360V - 3V^2$$

$$\div 3 : 3200 = 120V - V^2$$

$$V^2 - 120V + 3200 = 0$$


$$(V-40)(V-80) = 0$$

$$\text{so } \boxed{V=40} \text{ or } V=80$$

can't be  $V=80$  as then area of ① would be greater than 4800m which isn't possible.

Q8a)

$$\begin{aligned}\cos \alpha &= \frac{4}{5} \\ \sin \alpha &= \frac{3}{5} \\ \tan \alpha &= \frac{3}{4}\end{aligned}$$



$$\downarrow \text{N2L(Q)}: 3g - T = 3a \quad \sim (1)$$

$$\text{N2L(P)} \nearrow: T - 2g \sin \alpha - F = 2a \quad \sim (2)$$

$$(Q) \left\{ \begin{array}{l} s = 0.5 \\ u = 0 \\ v = 1.4 \\ a = a \\ t = \end{array} \right. \quad \left. \begin{array}{l} v^2 = u^2 + 2as \\ 1.4^2 = 0^2 + 2a(0.5) \\ 1.4^2 = a \quad // \end{array} \right.$$

$$\begin{aligned}\text{from (1): } T &= 3g - 3a = 3g - 3(1.4^2) \\ &= \boxed{23.5N}\end{aligned}$$

$$b) (2): T - 2g \sin \alpha - \mu R = 2a$$

$$23.5 - 2g\left(\frac{3}{5}\right) - \mu(2g \cos \alpha) = 2(1.4^2)$$

$$23.5 - \frac{6g}{5} - \mu\left(2g \times \frac{4}{5}\right) = 2(1.4^2)$$

$$\therefore \mu = \frac{23.5 - \frac{6g}{5} - 2(1.4^2)}{2g \times \frac{4}{5}} = \boxed{0.5}$$