



Question		Working	Answer	Mark	Notes
1	(a)	$\frac{42}{7} \times 2$ oe or $\frac{42}{7} \times 9[-12]$ or eg $\frac{2}{7} = \frac{t}{42}$		2	M1 for a correct method to find number of tables or total number of tables and chairs or setting up a correct equation to find the number of chairs May be seen within a ratio ie 12 : 42
			12		A1 cao
	(b)	$\frac{276 - 240}{240} [= 0.15]$ or $\frac{276 - 240}{240} \times 100$ or $\frac{276}{240} [= 1.15]$ or $\frac{276}{240} \times 100 [= 115]$		2	M1 for a correct method to find the % profit May use multiple number of tables eg $\frac{12 \times 276 - 12 \times 240}{12 \times 240}$
			15(%)		A1 cao
	(c)	$\frac{513}{1.35}$ oe		2	M1 for a correct method to find the price they were bought at Allow $x + 0.35x = 513$ oe eg $\frac{513 - x}{x} \times 100 = 35$ or $135\% = 513$ or $135\%_x = 513$ or $x + 35\%_x = 513$
			(\$) 380		A1 cao
	(d)	$732 \times 0.95 [= 695.4(0)]$ or $\frac{684}{0.95} [= 720]$ oe		3	M1 for a correct conversion
		“695.40” – 684 [= 11.4(0)] or 732 – “720” [= 12] oe			M1 for a correct subtraction “695.4(0)” and “720” must come from a correct calculation
			11.4(0) euros or \$12		A1 oe eg €11.4(0) or 12 dollars For a correct value with the correct units ISW
		<i>cas for each part</i>			Total 9 marks

Question		Working	Answer	Mark	Notes
2	(a)	$\frac{4 \times 180 - 4 \times 125}{2} [= 110]$ oe eg $\frac{(2 \times 6 - 4) \times 90 - 4 \times 125}{2} [= 110]$ or $360 - 2 \times 125 [= 110]$ oe eg $2 \times (180 - 125) [= 110]$ or $\frac{360 - 4 \times (180 - 125)}{2} [= 70]$ oe		5	M2 for correct method to find one of the unknown interior angles of the hexagon could use sum of interior angles or symmetry in the hexagon (eg line drawn from <i>C</i> to <i>L</i>) or for correct method to find the exterior angle (at <i>C</i> or <i>L</i>) of the hexagon can be implied by a correct calculation for <i>y</i> (If not M2 then M1 for: finding the angle sum of the hexagon eg $4 \times 180 [= 720]$ or $(2 \times 6 - 4) \times 90 [= 720]$ oe or finding one exterior angle of the hexagon eg $180 - 125 [= 55]$ or use of symmetry in the hexagon (eg line drawn from <i>C</i> to <i>L</i>) and finding <i>BCL</i> or <i>DCL</i> or <i>CLA</i> or <i>CLE</i> eg $\frac{360 - 2 \times 125}{2} [= 55]$ or $180 - 125 [= 55]$
		eg $\frac{(8 - 2) \times 180}{8} [= 135]$ oe eg $\frac{(2 \times 8 - 4) \times 90}{8} [= 135]$ or $\frac{360}{8} [= 45]$			M1 for finding one interior angle of the octagon or for finding one exterior angle of the octagon can be implied by a correct calculation for <i>y</i>
		eg $360 - \text{“110”} - \text{“135”}$ or $\text{“70”} + (180 - \text{“135”})$ or $\text{“70”} + \text{“45”}$ or $\text{“45”} + (180 - \text{“110”})$ oe			M1 (dep on all previous method marks awarded)
			115		A1 cao
	(b)	$10x + 25 + 7x - 83 = 180$ oe eg $17x - 58 = 180$		4	M1 for a correct equation in <i>x</i>
		$[x =] \frac{180 + 58}{17} [= 14]$			M1 (dep on first method mark) for a correct expression for <i>x</i>
		$\frac{360}{7 \times \text{“14”} - 83}$ or $\frac{360}{180 - (10 \times \text{“14”} + 25)}$			M1 ft dep on M1 ft their value of <i>x</i> provided <i>x</i> is from $\frac{180 + 58}{17}$ or clearly labelled as <i>x</i>
			24		A1 cao

ALT	(b)	eg $T(7x - 83) = 360$ and $T(10x + 25) = 90(2T - 4)$ oe eg $T(7x - 83) = 360$ and $T(10x + 25) = 180(T - 2)$ or $7x - 83 = \frac{360}{T}$ and $10x + 25 = \frac{180(T - 2)}{T}$			M1 for setting up two equations in x and T Allow the use of any letters for T and x provided they are not the same
		eg $7Tx = 360 + 83T$ oe and $10Tx = 90(2T - 4) - 25T$ oe eg $x = \frac{360 + 83T}{7T}$ oe and $x = \frac{90(2T - 4) - 25T}{10T}$ oe eg $70Tx = 3600 + 830T$ oe and $70Tx = 630(2T - 4) - 175T$ oe			M1 for making $7xT$ or x the subject of each equation or for coefficient of xT the same in both equations or for correct rearrangement of one equation followed by correct substitution into the other Allow the use of any letters for T and x provided they are not the same
		eg $1260T - 175T - 830T = 3600 + 2520$ oe			M1 for collecting T terms on 1 side and numbers on the other in a correct equation Allow the use of any letters for T and x provided they are not the same
			24		A1 cao
		<i>cas for each part</i>			<i>Total 9 marks</i>

Question	Working	Answer	Mark	Notes
In this question ignore any (incorrect) labelling of triangles				
3 (a)		Rotation	3	B1 allow rotate, rotated, rotation do not accept turn B0 if multiple transformations stated. Multiple transformations are when more than one of reflection (mirrored), rotation (turn), translation (move), enlargement (stretch / squash) is stated eg a vector or SF or equation of a line do not imply multiple transformations
		90[°] [anticlockwise]		B1 oe eg 270 clockwise or −270 Do not allow 90 clockwise or −90 or −90 anticlockwise
		(−1, −1)		B1 must be a coordinate and not a vector Do not allow if another coordinate is given as well
(b)		Correct triangle <i>C</i> at (1, −3), (2, −3), (3, −5)	2	B2 Fully correct triangle. Award 2 marks for a correct triangle drawn, irrespective of working in the working space. (B1 for correct line drawn or a triangle of the correct size and orientation, or for 2 correct vertices plotted correctly or 3 correct vertices listed) SCB1 for triangle <i>B</i> reflected in the line $y = -1$ (vertices (−3,−3) (−3,−4) (−5,−5)) or for triangle <i>A</i> reflected in the line $x = -1$ (vertices (−4, 1) (−3, 1) (−5, 3))
(c)		Correct triangle <i>D</i> at (2, −1), (4, −1), (6, 3)	2	B2 Fully correct triangle. Award 2 marks for a correct triangle drawn, irrespective of working in the working space. (B1 for a triangle of the correct size and orientation, or for 2 correct vertices plotted correctly or 3 correct vertices listed)
(d)	Points can be in any order $\begin{pmatrix} -2 & 1 \\ 0 & -2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 1 & 1 & 3 \end{pmatrix}$		3	M1 for intention to multiply the correct way, can be implied by writing the matrices in the correct order or correctly stating or plotting one point.
	Points can be in any order $\begin{pmatrix} -1 & -3 & -3 \\ -2 & -2 & -6 \end{pmatrix}$			M1 for at least two correct columns or correctly stating or plotting two points.
		Correct triangle <i>E</i> at (−1, −2), (−3, −2), (−3, −6)		A1 Fully correct triangle Award 3 marks for a correct triangle drawn, irrespective of working in the working space.
cas for each part				Total 10 marks



Question	Working	Answer	Mark	Notes
4(a)	$4x \leq 31 - 3$ oe or $x + \frac{3}{4} \leq \frac{31}{4}$		2	M1 for isolating the x term or reducing for x May be the wrong inequality sign or an = sign Also allow for a critical value of 7 with incorrect sign
		$x \leq 7$		A1 oe eg $(-\infty, 7]$ or $[-\infty, 7]$ do not isw $x = 7$
(b)		$-1 < x < 9$	2	B2 or other correct notation eg $(-1, 9)$ or $-1 < x, x < 9$ or $-1 < x$ and $x < 9$ or $-1 < x \cap x < 9$ (B1 for one end correct or for “ $-1 < x$ or $x < 9$ ”)
(c)	A single line joining $x = -1$ and $x = 7$		2	M1 Do not allow lines with arrows at end points or identification of any other values
				A1 Both end points identified using the correct symbols and one correct line drawn between the two correct points
	<i>cas for each part</i>			Total 6 marks

Question	Working	Answer	Mark	Notes
5(a)		$-3, 0, 9$	2	B2 all values correct (B1 for 2 values correct)
(b)			2	<p>M1 for at least 5 points plotted correctly ± 1 small square or for at least 4 points plotted correctly ± 1 small square and a smooth curve drawn through all their plotted points</p> <p>If you are unable to see the points plotted then allow if curve goes through them within/on the circles.</p>
		Correct graph		A1 for fully correct graph ± 1 small square. Do not allow straight line segments
(c)			2	M1 for the line $y = 2$ drawn on the grid that goes from at least $x = -1.5$ to $x = 1.3$ and intersects the curve at all points at which $y = 2$
		$-1.7, 0.2, 1.5$		<p>A1 dep on M1 and M1 being awarded in part (b) (condone straight line segments for this part of the question) For all 3 values and no extras Allow -1.9 to -1.5 and 0 to 0.4 and 1.3 to 1.7</p> <p>A0 if values are given to >2dp or if answers given as coordinates or y values given as well as x values or if answer is given as an inequality</p> <p>Do not isw so if $-1.7, 0.2, 1.5$ then $-1.7 < x < 1.5$ A0</p> <p>For reference values from calc are $x = -1.672981648, 1.469617434, 0.2033642138$</p>
	(a) <i>cas</i> (b) <i>cas</i> (c) <i>wr</i>			Total 6 marks

Question	Working	Answer	Mark	Notes
6 (a)		Correctly completed Venn diagram	3	B3 All 8 regions completed correctly (B2 for 5, 6 or 7 regions completed correctly (B1 for 3 or 4 regions completed correctly))
(b)		25	1	B1 ft from a diagram where values other than 0 are present in the required regions
(c)		$\frac{23}{48}$	2	B2 oe eg 0.47(91...) rounded or truncated to 2sf (B1 ft from a diagram where values other than 0 are present in the required regions for $\frac{23}{a}$ where $a > 23$ or $\frac{"9+14"}{a}$ or $\frac{"9+14"}{80} \div \frac{a}{80}$ (ft their 23) where $a > "9 + 14"$ or $\frac{b}{48}$ or $\frac{b}{80} \div \frac{48}{80}$ oe eg $\frac{b}{80} \div \frac{3}{5}$ where $b < 48$ or 23 : 48) B1 only for $\frac{9+14}{48}$
	<i>cas for each part</i>			Total 6 marks

Question	Working	Answer	Mark	Notes
7 (a)			2	M1 for at least 2 correct values in the correct places
		$\begin{pmatrix} 10 & -4 \\ -4 & 10 \end{pmatrix}$		A1 cao Do not isw
(b)			3	M2 for a matrix of the correct order and at least 4 correct values in the correct places (M1 for a matrix of the correct order and at least 2 correct values in the correct places)
		$\begin{pmatrix} -2 & -6 & 22 \\ 2 & 4 & -18 \end{pmatrix}$		A1 cao Do not isw
(c)	$10 \times 5 - (-6 \times -8) [= 2]$		4	M1 a correct method to find the determinant of A can be implied by multiplying matrix A by $\frac{1}{2}$
	$\frac{1}{2} \begin{pmatrix} 5 & 6 \\ 8 & 10 \end{pmatrix}$			M2 for 4 values in the correct places in their A ⁻¹ (M1 for at least 2 correct values in the correct places in their A ⁻¹)
	$\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} - \begin{pmatrix} 2.5 & 3 \\ 4 & 5 \end{pmatrix}$	$\begin{pmatrix} -0.5 & -3 \\ -4 & -3 \end{pmatrix}$		A1 cao
	<i>cas for each part</i>			Total 9 marks

Question	Working	Answer	Mark	Notes
8 (a)		15, 15, 16, x, 22	3	M1 for a list of 5 numbers where the mode is 15 or the median is 16 or the range is 7
				M1 for a list of 5 numbers with two of: the mode is 15 the median is 16 the range is 7
				A1 where x is 17, 18, 19, 20 or 21 (numbers can be in any order)
(b)	$8 \times 104 [= 832]$ or $5 \times 89 [= 445]$		3	M1 for the total weight of the 8 letters or the total weight of the 5 letters
	$8 \times 104 - 5 \times 89$ oe eg “832” – “445” [= 387]			M1 for the total weight of the 3 letters
		129		A1 cao
(c)	<u>cm square (5ss by 5ss)</u> $1 \times 3 + 3 \times 4$ cm squares = 75 or 15 cm squares = 75 or 1 cm square = 5 parcels oe <u>small squares (ss)</u> 75 lines of 5 ss = 75 or $5 \times 15 + 15 \times 20$ ss = 75 or 375 ss = 75 or 5 ss = 1 parcel oe		5	M1 for showing frequency is related to area by a correct calculation or a correct value for area and frequency or a correct value on FD axis (1 cm vertically is FD 5) this may come from $1 \times 15x + 3 \times 20x = 75$ oe eg $1 \times 3x + 3 \times 4x = 75$ Implied by a frequency of 20 or 15 or 60 or 30 or 20 seen Any fd / frequency may be seen in the correct place on the histogram
	<u>using FD</u> eg $2 \times 10 + 75 + 2 \times 15 + 4 \times 5 [= 145]$ $2 \times 20 + 2 \times 15 + 4 \times 5 [= 90]$ <u>cm square (5ss by 5ss)</u> eg $14 \times “5” + 75 [= 145]$ or $29 \times “5” [= 145]$ oe or $18 \times “5” [= 90]$ <u>small squares (ss)</u> eg $\frac{100 + 75 + 300 + 150 + 100}{5} [= 145]$ $\frac{200 + 150 + 100}{5} [= 90]$			M1 implies the previous method mark for a correct method to find the total number of parcels or a correct method to find the number of parcels greater than 4 kg Also allow correct method to find the number of parcels less than 4 kg eg $2 \times 10 + 1 \times 15 + 1 \times 20 [= 55]$ oe
	$\frac{90}{145}$ or $\frac{18}{29}$ or “0.62(06...)” oe			M1 implies both previous method marks for a correct first probability Only ft numbers from correct working Also allow the correct probability for a parcel being less than 4 kg eg $\frac{55}{145}$ or $\frac{11}{29}$ or 0.37(93...)

	$\frac{90}{145} \times \frac{90-1}{145-1}$ oe			M1 A correct product Allow $\left(\frac{90}{145}\right)^2 = \frac{324}{841} \approx 0.385$ Condone $2 \times \frac{90}{145} \times \frac{90-1}{145-1}$ and $2 \times \left(\frac{90}{145}\right)^2$
		$\frac{89}{232}$		A1 oe eg 0.38(36...) [decimal or % 2sf or better] Allow 0.38 – 0.3853
	<i>cas for each part</i>			<i>Total 11 marks</i>

Question	Working	Answer	Mark	Notes
9 (a) (i)		80	1	B1 cao
(a) (ii)		correct reason	1	B1 dep on B1 in (i) for <u>Angle</u> at the <u>centre</u> is <u>2 ×</u> (double) angle at <u>circumference</u> / <u>angle at circumference</u> is <u>½</u> angle at <u>centre</u> allow the symbol for the word ‘angle’
(b)	eg “80” – 37 or 360 – “80” – 37 – (360 – 160) or 180 – “80” – 37 – (180 – 160) oe		2	M1 ft their answer to part (a)(i) A correct method to find angle <i>TPO</i>
		43		A1 cao
(c)	Throughout part (c) allow 3.1, 3.14, etc or $\frac{22}{7}$ for π and allow any letter or symbol for r			
(c)	$\frac{160}{360} \times \pi \times r^2 = \frac{196}{25} \pi$ oe eg $\frac{160}{360} \times r^2 = \frac{196}{25}$		6	M1 for an equation in the form $\frac{160}{360} \times \pi \times (...r)^2 = \frac{196}{25} \pi$ or $\frac{160}{360} \times \pi \times ...r^2 = \frac{196}{25} \pi$
	$[r =] \sqrt{\frac{196\pi}{25} \div \frac{160\pi}{360}}$ oe eg $[r =] \sqrt{\frac{196}{25} \times \frac{9}{4}} \left[= \frac{21}{5} = 4.2 \right]$			M1 implies previous M a correct calculation for the radius of the sector (slant height of cone)
	[radius of cone =] eg $\frac{196}{25} \div "4.2" \left[= \frac{28}{15} = 1.866... \right]$ oe eg $\frac{\frac{160}{360} \times 2\pi \times "4.2"}{2\pi} \left[= \frac{28}{15} = 1.866... \right]$			M1 for $\frac{196}{25} \div "r"$ where “ r ” is the candidate’s value for the radius of the sector. If r is incorrect it must be clearly labelled and working for this method mark must be shown Condone radius of cone being labelled as l
	(cone height=) $\sqrt{"4.2"^2 - "1.866..."^2} \left[= \frac{7\sqrt{65}}{15} = 3.762... \right]$			M1 dep on first and third method marks, for a correct calculation for the height of the cone Follow through candidate’s values for r or r and l . These must be clearly labelled and working shown if they are incorrect Condone $\sqrt{"1.866..."^2 - "4.2"^2}$
	$\frac{1}{3} \times \pi \times "1.866..."^2 \times "3.762..." \left[= 4.36(99..) \pi \right]$			M1 dependent on previous four method marks for a correct calculation for the volume where r and h are from correct methods NB: $\frac{1}{3} \times \pi \times "4.2"^2 \times "3.762..."$ does not get this mark
		13.7 (cm ³)		
	cas			Total 10 marks