Write your name here Surname	Other n	ames
Pearson Edexcel International GCSE	Centre Number	Candidate Number
<b>Further Pu</b>	ire Math	ematics
Paper 2	are matri	Ciliatics
Paper 2 Wednesday 20 June 2018		Paper Reference
Paper 2		

### **Instructions**

- Use **black** ink or ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
  - there may be more space than you need.

#### Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

### **Advice**

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶



P53392RA ©2018 Pearson Education Ltd.



# Answer all TEN questions.

# Write your answers in the spaces provided.

### You must write down all the stages in your working

	Tou must write down an the stages in your working.	
1	In triangle $ABC$ , $AB = 9$ cm, $BC = 6$ cm and $CA = 8$ cm.	
	Find, in degrees to the nearest $0.1^{\circ}$ , the size of angle <i>BAC</i> .	(2)
		(3)

Question 1 continued	
	(Total for Question 1 is 3 marks)



2	Differentiate with respect to x	
	(a) $e^{3x} \cos 2x$	(3)
	(b) $\frac{2e^x}{(2x^2-1)}$	(3)

Question 2 continued	
	(Total for Question 2 is 6 marks)



3	The volume of liquid in a container is $V \text{ cm}^3$ when the depth of the liquid is $h \text{ cm}$ . Liquid is leaking from the container at a rate of $24 \text{ cm}^3/\text{s}$ .		
	Given that $V = 5h^3$ , find the rate, in cm/s, at which the depth of the liquid is decreasive your answer to 2 significant figures.	asing when $V = 800$ (7)	

Question 3 continued
(Total for Question 3 is 7 marks)



4	<ul> <li>(a) Find the exact value of the root of the equation e<sup>3x</sup> = 8 Give your answer in the form ln a, where a is an integer.</li> <li>The curve C<sub>1</sub> has equation y = 2e<sup>3x</sup> and the curve C<sub>2</sub> has equation y = (e<sup>3x</sup> - 4)<sup>2</sup></li> <li>The curves C<sub>1</sub> and C<sub>2</sub> intersect at the points P and Q.</li> <li>(b) Use algebra to find the exact coordinates of the points P and Q.</li> </ul>	(2)
	(c) Find, to 3 decimal places, the length of <i>PQ</i> .	(2)



Question 4 continued	

Question 4 continued
(Total for Question 4 is 9 marks)



5	The sum of the first term and the third term of a geometric series is 75  The sum of the second term and the third term is 45	
	<ul><li>(a) Find the two possible values of the common ratio of the series.</li><li>Given that the series is convergent with sum to infinity S,</li></ul>	(5)
	(b) find the value of <i>S</i> .	(3)

Question 5 continued
(Total for Question 5 is 8 marks)



6




Diagram **NOT** accurately drawn

Figure 1

Figure 1 shows a rectangular box with length 5x cm, width 2x cm and height h cm. The box has a base but no top. The volume of the box is 1000 cm<sup>3</sup> and the total external surface area of the box is S cm<sup>2</sup>

(a) Show that 
$$S = 10x^2 + \frac{1400}{x}$$

(4)

Given that x can vary,

(b) find, to 3 significant figures, the minimum value of *S*.

**(5)** 

(c) Verify that your answer to part (b) does give the minimum value of S.

(2)


Question 6 continued	



Question 6 continued

Question 6 continued
(Total for Question 6 is 11 marks)



- 7 (a) Expand  $\left(1 + \frac{2x}{5}\right)^{\frac{1}{2}}$  in ascending powers of x up to and including the term in  $x^3$ , giving each coefficient as an exact fraction in its lowest terms.
- (3)
- (b) Expand  $\left(1 \frac{2x}{5}\right)^{-\frac{1}{2}}$  in ascending powers of x up to and including the term in  $x^3$ , giving each coefficient as an exact fraction in its lowest terms.
- (3)
- (c) Write down the range of values of x for which both of your expansions are valid.
  - (1)
- (d) Expand  $\left(\frac{5+2x}{5-2x}\right)^{\frac{1}{2}}$  in ascending powers of x up to and including the term in  $x^2$ , giving each coefficient as an exact fraction in its lowest terms.
- (3)

(e) Hence use algebraic integration to obtain an estimate of

$$\int_{0.1}^{0.3} \left( \frac{5 + 2x}{5 - 2x} \right)^{\frac{1}{2}} dx$$

Give your answer to 4 significant figures.

/		á	١
(	4	Į	


	4	e	ė	g	9					
	,	í	4	4	ì					
	1	į	4		ď					
	1	Ľ	2							
	j	Ė	ė	ġ	ø					
		4	4	4	9					
	1	ι	j	ŗ	١					
	ì	ĺ	į		í					
	1	4		ſ	į					
	i	Ē	7		7					
	1	r								
	3	į	7	j	P					
	1	4	_	-						
	1									
	1	Ľ	J	ľ	ı					
	]	ŀ	٠		i					
	1	i		į						
	j	Ė	١	Ė	2					
	į	5	5	Š	-					
	1	3	5	à	ń		٠			
	1									
	ł	ľ	7		Ą					
	1	ľ		3	)					
	7		į	3	p					
	j	Ġ	É	_	ė					
	ì	à	i	ė	ķ					
	4	ζ	ė	í	)		٠			
	i	ŕ	۹	٩	ĺ					
	ļ		7		ą					
	•				•					
	:									
							٠			
									ì	
					′			/	,	
		į			ž				,	
						,	/			
/			,	/			,	/		
×	á	r	ė	ġ	ø				,	
\	3	í	ļ	9	ņ	/		,		
(	1	þ	į	É	į					
ì	Į	Ġ	٥	Ġ	ź			į	,	
>	ķ	í	í	í	ĕ			ċ	,	
(		7	9	4						
Ż	1	ì	j	P	١	١	×	×		
Ì	K	Ę	į	Ì	ě			Ì	>	
`	1 ×	9		Ē	a ×	Ż	×	`	>	
<	1	í	2	4	ŕ	>	×	Ć		
ì	k	¢	4	7	ŕ			ì	>	
3	į			þ	į	(		×	>	
(	1	d		ŕ	-			×		
Ċ	,	Ś		è	5	1	×	×		
٦,			¥	7	- - - -		×	< x		
×	ķ	ķ	Ĭ			< <	× × ×	XXXX	> > > .	
(	k	ŀ	ì			////	×	/×/×		
<	k	Ì	1			< \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	×××××××××××××××××××××××××××××××××××××××	/×/×/×		
	K		ì			/ \ / \ / \ / \ / \ / \ / \ / \ / \ / \	· × · × · × · × · ×	/×/×/×/		
	K					/ \ / \ / \ / \ / \ /	· × · × · × · × · × ·	/×/×/×/×/	/ > / > / > / >	
						/ \ / \ / \ / \ / \ / \ / \ / \ / \ / \		/×/×/×/×/×	/ > / > / > / > / > / >	
						/ \ / \ / \ / \ / \ / \ / \ / \ / \ / \	·×·×·×·×·×	/×/×/×/×/×/×/×		
< < < < < × <						< > < < < < < < < < < < < < < < < < <	· × · × · × · × · × · × · × · ×	/×/×/×/×/×/×/	/> /> /> /> /> /> /> /> /> /> /	
< < < < < × × × × × × × × × × × × × × ×						/ \ / \ / \ / \ / \ / \ / \ / \ / \ / \	· × · × · × · × · × · × · × · × ·	/×/×/×/×/×/×/×/		
< \ < \ < \ < \ < \ < \ < \ < \ < \ < \						/ \ / \ / \ / \ / \ / \ / \ / \ / \ / \		/ × / × / × / × / × / × / ×	ノン ノン ノン ノン ノン ノン ノン ノン	
< < < < < < < < < < < < < < < < < < <						/ \ / \ / \ / \ / \ / \ / \ / \ / \ / \	· · · · · · · · · · · · · · · · · · ·	/ × / × / × / × / × / × / × / × /	ノン ノン ノン ノン ノン ノン ノン ノン	
< < < < < < < < < < < < < < < < < < <							·X·XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	/ × / × / × / × / × / × / × / × /		
< < < < < < < < < < < < < < < < < < <								,	/> /> /> /> /> /> /> /> /> /> /> /> />	
< < < < < < < < < < < < < < < < < < <								/ × / × / × / × / × / × / × / × / × / ×	ノンノン ノン ノン ノン ノン ノン ノン ノン ノン	
/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\								/ × / × / × / × / × / × / × / × / × / ×	ノン	
/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\								/×/×/×/×/×/×/×/×/×/×/×/×/×/		
/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\						/ \		/×/×/×/×/×/×/×/×/×/×/×/×/×/	ノンフンファンファンファンファファファファファファファファ	
/\/\/\/\/ /\/\/ /\/ /\/								/ * / * / * / * / * / * / * / * / * / *	/> /> /> /> /> /> /> /> /> /> /> /> /> /	
<. <. <. <. <. < < < < < < < < < < < <						/ \			/> /> /> /> /> /> /> /> /> /> /> /> /> /	
<, /, /, /, /×/×/×/×/×/×/×/×/×/×/×/×/×/×/								/ × / × / × / × / × / × / × / × / × / ×		
<, /, /, /, /, /, /, /, /, /, /, /, /, /,				コノトラー ゴンコンコンド・マント マンザラ アンメンメンメンシン				/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×		
/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\/\								/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×		
				■フトア ■ は と は マスノ ア 写 と 可 は ア 高 フ 明 フ ア ン ン ン ン ン ン ン ン ン ン ン ン ン ン ン ン ン ン				/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×		
								\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\		
								/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×/×		
								,		
								,		
				■プレフ ■ 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1						

Question 7 continued



Question 7 continued	

Question 7 continued

(Total for Question 7 is 14 marks)
(Total for Question / is 17 marks)



$$\cos(A+B) = \cos A \cos B - \sin A \sin B$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

Using the above identities

- (a) show that (i)  $\cos 2\theta = 1 2\sin^2 \theta$ 
  - (ii)  $\sin 2\theta = 2\sin \theta \cos \theta$

(3)

$$f(\theta) = \cos 4\theta + 2\cos 2\theta$$

(b) Show that  $f(\theta) = 8\sin^4\theta - 12\sin^2\theta + 3$ 

(4)

(c) Solve, giving your solutions to 3 significant figures, the equation

$$4\sin^4 x^\circ - 6\sin^2 x^\circ - \cos 2x^\circ + 1.2 = 0$$

 $0 \leqslant x < 90$ 

(4)

- (d) (i) Find  $\int (2\sin^4\theta 3\sin^2\theta) d\theta$ 
  - (ii) Hence find the exact value of  $\int_0^{\frac{\pi}{3}} (2\sin^4\theta 3\sin^2\theta) d\theta$

Give your answer in the form  $a\sqrt{b}-c\pi$  where a and c are rational numbers and b is a prime number.



**(5)** 





Question 8 continued



Question 8 continued	

**Question 8 continued** 

Y	
2	
2	
Š	
1.7	
/./	
:/:>	
<b>4</b> /	
11.	
<b>X</b> ./	
≪ ⁄ .	
XX	
XX I	
$\mathbb{Z}\mathbb{X}$	
XX	
$\times$	
×× I	
<del>&gt;</del>	
<u> </u>	
<b>3</b>	
XXX	
***	
<b>***</b>	
<b>***</b>	
<b>2000</b>	
Z	
KITEIN	
WKIEETNE	
WRITEIN	
OT WELLE IN T	
NO I WISHE IN	
NOT WRITE IN	
O NOT WRITE IN	
DONOT WRITE IN	
DO NOT WRITE IN T	
DO NOT WRITE IN T	
DONOTWENT	
DONO! WKITEINT	
DONO! WKITEINT	(Total for Question 8 is 16 marks)



9	The points $A$ , $B$ and $C$ have coordinates $(-4, 4)$ , $(1, 6)$ and $(-2, -1)$ respectively.	
	(a) Show, by calculation, that $AB$ is perpendicular to $AC$ .	(4)
	(b) Find an equation for BC in the form $px + qy + r = 0$ , where p, q and r are integers.	
	The line $l$ is the perpendicular bisector of $AB$ .	(3)
	(c) Find an equation for $l$ .	
	(c) 1 mu un equation ter ti	(4)
	The line $l$ and the line $BC$ intersect at the point $E$ .	
	(d) Find the coordinates of $E$ .	(2)
	(e) Calculate the area of triangle <i>AEC</i> .	(2)
	(c) Calculate the area of triangle ALC.	(4)
******		

۰
٠
×
(
ż
١
1
Ž
٩
1
ì
١,
ì
k
١
\ \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
< / < / < / < / < / < / < / < /
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\

Question 9 continued



Question 9 continued	

Question 9 continued	
(Total	for Question 9 is 17 marks)



10	The curve C has equation $y^2 = 16x$ where $y \ge 0$	
	Given that the point A with coordinates $(a, 2a)$ where $a \neq 0$ lies on C,	
	(a) find the value of a.	
		(2)
	The line $l$ passes through $A$ and has gradient $-2$	
	Given that $l$ crosses the $x$ -axis at the point $B$ ,	
	(b) find the <i>x</i> coordinate of <i>B</i> .	(2)
		(2)
	The finite region enclosed by $C$ , $l$ and the $x$ -axis is rotated through 360° about the $x$ -axis.	
	(c) Using algebraic integration, find, to 3 significant figures, the volume of the solid generated.	
	Samuel.	(5)

Question 10 continued



DO NOT WRITE IN THIS AREA

Question 10 continued		
	(Total for Question 10 is 9 marks)	
	TOTAL FOR PAPER IS 100 MARKS	