

Leave  
blank

2. [In this question  $\mathbf{i}$  and  $\mathbf{j}$  are horizontal unit vectors due east and due north respectively and position vectors are given relative to a fixed origin  $O$ .]

At time  $t = 0$ , a bird  $A$  leaves its nest, that is located at the point with position vector  $(20\mathbf{i} - 17\mathbf{j})\text{ m}$ , and flies with constant velocity  $(-6\mathbf{i} + 7\mathbf{j})\text{ m s}^{-1}$ . At the same time a second bird  $B$  leaves its nest which is located at the point with position vector  $(-8\mathbf{i} + 9\mathbf{j})\text{ m}$  and flies with constant velocity  $(p\mathbf{i} + 2p\mathbf{j})\text{ m s}^{-1}$ , where  $p$  is a constant. At time  $t = 4\text{ s}$ , bird  $B$  is south west of bird  $A$ .

- (a) Find the direction of motion of  $A$ , giving your answer as a bearing to the nearest degree. (3)  
(b) Find the speed of  $B$ . (10)

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**Question 2 continued**Leave  
blank

P 5 4 8 2 9 A 0 5 2 4

Leave  
blank

## **Question 2 continued**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**



**DO NOT WRITE IN THIS AREA****DO NOT WRITE IN THIS AREA****DO NOT WRITE IN THIS AREA****Question 2 continued**Leave  
blank**Q2****(Total 13 marks)**