

3D tumor spheroids

This manuscript ([permalink](#)) was automatically generated from [SharikHR/review@c21caeb](#) on May 7, 2020.

Authors

- **Sharik Hernandez**

 [XXXX-XXXX-XXXX-XXXX](#) ·  [SharikHR](#)

Department of · Funded by Grant XXXXXXXX

- **None**

 [XXXX-XXXX-XXXX-XXXX](#)

Department of; Department of

Abstract

Open collaborative writing with Manubot [1](#)

Introduction

PDAC belongs to the top five of cancer-related deaths in the world and the poor prognosis is primarily due to its advanced stage at diagnosis, the progress in its treatment remains too slow as a consequence of the complex physiopathology of this tumor characterized by a heterogeneous cellular composition and the accumulation of a very dense fibrotic tissue [???]: Due to that pancreatic cancer is a heterogeneous disease, is often modelled using established cell lines in the laboratory.

In recent years, three-dimensional (3D) culture systems have gained increasing recognition as an effective tool for biological research. One widely used 3D culturing technique is the application of multicellular spheroids (MCS). Cells cultured in 3D more closely mimic the physiological environment compared to conventional monolayer culture systems. Spheroids are three-dimensional spherical cellular aggregates with high cell-density, that more closely simulate conditions existing in solid tumors where hypoxia and alterations related to intracellular metabolism occur due to poor availability of nutrients from blood vessels [2](#).

Pancreatic tumor microenvironment

In Pancreatic ductal adenocarcinoma (PDA), the major components of the tumor microenvironment are a complex population of fibroblasts forming the bulk of the stroma, vasculature, inflammatory and immune cells [3](#). PDA is associated with evolving alterations in the tumor microenvironment, including increasing fibrosis and extracellular matrix deposition (desmoplasia). Increasing desmoplasia accompanies progressive disease and creates intratumoral pressure that compresses the vasculature, resulting in limited blood flow to the tumor and consequent hypoxia and low nutrient delivery [4](#).

Interactions between the neoplastic and non-neoplastic cells and cellular matrix have been proposed to stimulate the extensive desmoplastic reaction. Stroma production is promoted by the activation of multiple cell signalling pathways [5](#); 10.1002/mc.20827 these signalling cascades lead to secretion of structural matrix components including proteoglycans, collagen and fibronectin and the activation of catalytic enzymes such as proteinases.

Matrix metalloproteinases (MMPs) are a family of zinc-dependent endopeptidases, they are involved in degradation of the extracellular matrix. MMPs support tissue remodeling and stimulate neovascularization and inflammatory response, both in physiological and in pathological conditions, for example, in tumors [6](#).

MMP-2, MMP-7, and MMP-9 expressions correlate with various morphological features of the PDAC tumor such as inflammation, necrosis, and formation of the new blood vessels [7](#) and their tissue inhibitors (TIMPs) TIMPs 1–3 compared with normal pancreas [8](#).

Pancreatic stellate cells (PSC) are responsible for desmoplastic generation [9](#). The activation of PSC from a quiescent to an activated state is an intercellular stimuli from tumor-stromal interactions. PSCs can be transformed into myofibroblast-like cells, which express α -smooth muscle actin (α -SMA) as cancer-associated fibroblasts (CAF) [10](#); 10.1053/j.gastro.2010.05.084.

CAF are important components of tumor stroma and affect cancer growth, survival, metastasis, angiogenesis and resistance to chemotherapy or radiotherapy through various cytokines [11](#), and contribute to a diminished immune function [12](#).

2D and 3D cell culture

Cell culture is a widely used *in vitro* tool to improve the understanding of cell biology, cellular mechanisms, tissue morphology, drug action, protein production and the development of tissue engineering [13](#).

Three-dimensional (3D) cell culture is well documented to regain intrinsic metabolic properties and to better mimic the *in vivo* situation than two-dimensional (2D) cell culture [14](#).

A comparative molecular analyses from 2D and 3D *in vitro* conditions revealed that cancer cell phenotype is highly influenced by its microenvironment. Through the examination of cancer cell transcriptional behavior, cellular processes were closely linked to functions promoted by different culturing conditions. Cancer cells grown in monolayers favored rapid proliferation, and this behavior was corroborated by up-regulation of cell cycle progression genes in addition to metabolic processes that synthesize DNA, RNA, and proteins. Unlike cancer cells cultured in monolayers, cancer cells in 3D down-regulate proliferative processes, while up-regulating genes involved in ECM organization and cell adhesion [15](#).

When performing 3D cell culture experiments, the cell environment can be manipulated to mimic that of a cell *in vivo* and provide more accurate data about cell-to-cell interactions, tumor characteristics, drug discovery, metabolic profiling, stem cell research, and other types of diseases [16](#).

Alterations in cancer cell behavior under different growth conditions underlie the importance of defining culturing conditions that preserve endogenous tumor behavior. While monolayer cultures promote the most non-native behavior of cancer cells, this model still maintains value due to its ease and scalability towards applications targeting cancer-growth driving pathways. However, analysis and discovery of potential therapeutics targeting stromal interactions, ECM development, or cell signaling may yield erroneous results in this system. The 3D culturing and inclusion of stromal cell types do show increased similarity to *in vivo* cancer behavior. However, improvements upon *ex vivo* culture conditions that allow all stromal components to persist will greatly enhance our ability to conduct pre-clinical screens that may more closely recapitulate the biological responses of patients [15](#).

The structure of multicellular spheroids.

Methods for the generation of MCS

MCS and PDA

References

1. Open collaborative writing with Manubot

Daniel S. Himmelstein, Vincent Rubinetti, David R. Slochower, Dongbo Hu, Venkat S. Malladi, Casey S. Greene, Anthony Gitter

PLOS Computational Biology (2019-06-24) <https://doi.org/c7np>

DOI: [10.1371/journal.pcbi.1007128](https://doi.org/10.1371/journal.pcbi.1007128) · PMID: [31233491](https://pubmed.ncbi.nlm.nih.gov/31233491/) · PMCID: [PMC6611653](https://pubmed.ncbi.nlm.nih.gov/PMC6611653/)

2. Cancer cell spheroids are a better screen for the photodynamic efficiency of glycosylated photosensitizers

Patrícia M. R. Pereira, Naxhije Berisha, N. V. S. Dinesh K. Bhupathiraju, Rosa Fernandes, João P. C. Tomé, Charles Michael Drain

PLOS ONE (2017-05-17) <https://doi.org/f986w7>

DOI: [10.1371/journal.pone.0177737](https://doi.org/10.1371/journal.pone.0177737) · PMID: [28545086](https://pubmed.ncbi.nlm.nih.gov/28545086/) · PMCID: [PMC5435229](https://pubmed.ncbi.nlm.nih.gov/PMC5435229/)

3. Stromal biology of pancreatic cancer

Gerald C. Chu, Alec C. Kimmelman, Aram F. Hezel, Ronald A. DePinho

Journal of Cellular Biochemistry (2007-07-01) <https://doi.org/b8w83q>

DOI: [10.1002/jcb.21209](https://doi.org/10.1002/jcb.21209) · PMID: [17266048](https://pubmed.ncbi.nlm.nih.gov/17266048/)

4. Pancreatic Cancer Metabolism: Breaking It Down to Build It Back Up

R. M. Perera, N. Bardeesy

Cancer Discovery (2015-11-03) <https://doi.org/ggt22b>

DOI: [10.1158/2159-8290.cd-15-0671](https://doi.org/10.1158/2159-8290.cd-15-0671) · PMID: [26534901](https://pubmed.ncbi.nlm.nih.gov/26534901/) · PMCID: [PMC4687899](https://pubmed.ncbi.nlm.nih.gov/PMC4687899/)

5. Tumor–stromal cell interaction under hypoxia increases the invasiveness of pancreatic cancer cells through the hepatocyte growth factor/c-Met pathway

Takao Ide, Yoshihiko Kitajima, Atsushi Miyoshi, Takao Ohtsuka, Mayumi Mitsuno, Kazuma Ohtaka, Yasuo Koga, Kohji Miyazaki

International Journal of Cancer (2006-12-15) <https://doi.org/bcmh42>

DOI: [10.1002/ijc.22178](https://doi.org/10.1002/ijc.22178) · PMID: [16998831](https://pubmed.ncbi.nlm.nih.gov/16998831/)

6. ALTERED MATRIX METALLOPROTEINASE EXPRESSION ASSOCIATED WITH ONCOGENE-MEDIATED CELLULAR TRANSFORMATION AND METASTASIS FORMATION

R Baruch

Cell Biology International (2001-05) <https://doi.org/fnvccv>

DOI: [10.1006/cbir.2000.0647](https://doi.org/10.1006/cbir.2000.0647) · PMID: [11401328](https://pubmed.ncbi.nlm.nih.gov/11401328/)

7. Expressions of Matrix Metalloproteinases 2, 7, and 9 in Carcinogenesis of Pancreatic Ductal Adenocarcinoma

Katarzyna Jakubowska, Anna Pryczynicz, Joanna Januszevska, Iwona Sidorkiewicz, Andrzej Kemona, Andrzej Niewiński, Łukasz Lewczuk, Bogusław Kędra, Katarzyna Guzińska-Ustymowicz

Disease Markers (2016) <https://doi.org/f9hdkw>

DOI: [10.1155/2016/9895721](https://doi.org/10.1155/2016/9895721) · PMID: [27429508](https://pubmed.ncbi.nlm.nih.gov/27429508/) · PMCID: [PMC4939209](https://pubmed.ncbi.nlm.nih.gov/PMC4939209/)

8. Comprehensive Analysis of Matrix Metalloproteinase and Tissue Inhibitor Expression in Pancreatic Cancer: Increased Expression of Matrix Metalloproteinase-7 Predicts Poor Survival

L. E. Jones

Clinical Cancer Research (2004-04-15) <https://doi.org/c7sfqg>

DOI: [10.1158/1078-0432.ccr-1157-03](https://doi.org/10.1158/1078-0432.ccr-1157-03) · PMID: [15102692](https://pubmed.ncbi.nlm.nih.gov/15102692/)

9. **Conophylline suppresses pancreatic cancer desmoplasia and cancer-promoting cytokines produced by cancer-associated fibroblasts**
Norihiro Ishii, Kenichiro Araki, Takehiko Yokobori, Kei Hagiwara, Dorgormaa Gantumur, Takahiro Yamanaka, Tadashi Handa, Mariko Tsukagoshi, Takamichi Igarashi, Akira Watanabe, ... Ken Shirabe
Cancer Science (2018-12-13) <https://doi.org/ggt57m>
DOI: [10.1111/cas.13847](https://doi.org/10.1111/cas.13847) · PMID: [30353606](https://pubmed.ncbi.nlm.nih.gov/30353606/) · PMCID: [PMC6317962](https://pubmed.ncbi.nlm.nih.gov/PMC6317962/)
10. **A Starring Role for Stellate Cells in the Pancreatic Cancer Microenvironment**
Minoti V. Apte, Jeremy S. Wilson, Aurelia Lugea, Stephen J. Pandol
Gastroenterology (2013-05) <https://doi.org/f2j2rr>
DOI: [10.1053/j.gastro.2012.11.037](https://doi.org/10.1053/j.gastro.2012.11.037) · PMID: [23622130](https://pubmed.ncbi.nlm.nih.gov/23622130/) · PMCID: [PMC3729446](https://pubmed.ncbi.nlm.nih.gov/PMC3729446/)
11. **Cancer-associated fibroblasts in pancreatic adenocarcinoma**
Boju Pan, Quan Liao, Zheyu Niu, Li Zhou, Yupei Zhao
Future Oncology (2015-09) <https://doi.org/f7sxbh>
DOI: [10.2217/fon.15.176](https://doi.org/10.2217/fon.15.176) · PMID: [26284509](https://pubmed.ncbi.nlm.nih.gov/26284509/)
12. **Human Pancreatic Carcinoma-Associated Fibroblasts Promote Expression of Co-inhibitory Markers on CD4+ and CD8+ T-Cells**
Laia Gorchs, Carlos Fernández Moro, Peter Bankhead, Katharina P. Kern, Imrul Sadeak, Qingda Meng, Elena Rangelova, Helen Kaipe
Frontiers in Immunology (2019-04-24) <https://doi.org/ggt57p>
DOI: [10.3389/fimmu.2019.00847](https://doi.org/10.3389/fimmu.2019.00847) · PMID: [31068935](https://pubmed.ncbi.nlm.nih.gov/31068935/) · PMCID: [PMC6491453](https://pubmed.ncbi.nlm.nih.gov/PMC6491453/)
13. **2D and 3D cell cultures – a comparison of different types of cancer cell cultures**
Marta Kapalczyńska, Tomasz Kolenda, Weronika Przybyła, Maria Zajączkowska, Anna Teresiak, Violetta Filas, Matthew Ibbs, Renata Bliźniak, Łukasz Łuczewski, Katarzyna Lamperska
Archives of Medical Science (2016) <https://doi.org/ggt3hn>
DOI: [10.5114/aoms.2016.63743](https://doi.org/10.5114/aoms.2016.63743) · PMID: [30002710](https://pubmed.ncbi.nlm.nih.gov/30002710/) · PMCID: [PMC6040128](https://pubmed.ncbi.nlm.nih.gov/PMC6040128/)
14. **Oncogenic human herpesvirus hijacks proline metabolism for tumorigenesis**
Un Yung Choi, Jae Jin Lee, Angela Park, Wei Zhu, Hye-Ra Lee, Youn Jung Choi, Ji-Seung Yoo, Claire Yu, Pinghui Feng, Shou-Jiang Gao, ... Jae U. Jung
Proceedings of the National Academy of Sciences (2020-04-07) <https://doi.org/ggt4fr>
DOI: [10.1073/pnas.1918607117](https://doi.org/10.1073/pnas.1918607117) · PMID: [32213586](https://pubmed.ncbi.nlm.nih.gov/32213586/) · PMCID: [PMC7149499](https://pubmed.ncbi.nlm.nih.gov/PMC7149499/)
15. **Comparative Molecular Analysis of Cancer Behavior Cultured In Vitro, In Vivo, and Ex Vivo**
Nicholas R. Hum, Aimy Sebastian, Sean F. Gilmore, Wei He, Kelly A. Martin, Aubree Hinckley, Karen R. Dubbin, Monica L. Moya, Elizabeth K. Wheeler, Matthew A. Coleman, Gabriela G. Loots
Cancers (2020-03-14) <https://doi.org/ggt4ft>
DOI: [10.3390/cancers12030690](https://doi.org/10.3390/cancers12030690) · PMID: [32183351](https://pubmed.ncbi.nlm.nih.gov/32183351/) · PMCID: [PMC7140030](https://pubmed.ncbi.nlm.nih.gov/PMC7140030/)
16. **Is It Time to Start Transitioning From 2D to 3D Cell Culture?**
Caleb Jensen, Yong Teng
Frontiers in Molecular Biosciences (2020-03-06) <https://doi.org/ggt4fs>
DOI: [10.3389/fmolb.2020.00033](https://doi.org/10.3389/fmolb.2020.00033) · PMID: [32211418](https://pubmed.ncbi.nlm.nih.gov/32211418/) · PMCID: [PMC7067892](https://pubmed.ncbi.nlm.nih.gov/PMC7067892/)